REVIEW ARTICLE Agricultural Reviews, Volume 44 Issue 2: 245-251 (June 2023)

Conservation Agriculture (CA)-Based Tillage Practices in Maize for Enhancing Crop Yield, Resource use Efficiency and Soil Health: A Review

Niraj Biswakarma[1], Bishesh Rai[2], Somanath Nayak[1], Radheshyam[1]

10.18805/ag.R-2184

ABSTRACT

The triple challenge of acute water crisis, stagnant crop yield and soil health deterioration in NW Indo Gangetic Plains (IGP) ramble a search for potential alternative crop establishment technique (CET). Hence, maize-based crop rotations under best-bet conservation agriculture (CA) practices can plays a critical role in sustainable crop production. The CA-based tillage and CET *viz*. zero tillage (ZT) and permanent raised beds (PB) hold potential to intensify crop yield per hectare, improving resource use efficiency besides bringing desirable changes in soil physico-chemical and biological properties. Therefore, it needs to be popularized in larger scale chiefly under maize-based rotation to makes farming more attractive, profitable and sustainable.

Key words: Conservation agriculture, Maize, Resource use efficiency, Soil health, Tillage, Yield.

Globally, agriculture is entering a new era confronted with problems related to climatic change and issues of food security. Maize (*Zea mays* L.), being a versatile and highly adaptive crop to diverse agro-ecologies with highest genetic yield potentials fundamental to ensure food and nutritional security to the millions (Shiferaw *et al*., 2011; Jat *et al*., 2013). In India, maize occupies an area of ~ 9.8 M ha with annual production of ~19.8 million tones (GOI, 2020). Among various maize-based rotation maize-fallow, maize-mustard, maize-chickpea, maize-maize, maize-potato, are the promising systems across the country. However, conventional maize production have several restraints, mainly due to poor crop establishment and inputs management (Yadav *et al*., 2017; Parihar *et al*., 2017) leading to lower crop productivity and resource use-efficiency. Similarly, soils of the major maize growing are as low in soil organic matter (SOM) and nitrogen (N) (Jat *et al*., 2014). In this context, CA-based maize rotation is being advocated as an alternative to conventional practices. The conservation agriculture concept is evolved in a response to the concerns of agricultural sustainability (Bhan and Behra, 2014), which is based on three fundamental inter-linked principles; (i) minimal mechanical soil disturbance, (ii) permanent soil cover and (iii) crop diversification (Fig 1). The years of intensive tillage had a remarkable negative outcome on the soil health (Parihar *et al*., 2016) and resource use efficiency (Govaerts *et al*., 2005) thereby affecting crop yields. Similarly, tillage practices also determine the content and distribution of SOM, which is a key indicator of a soil quality (Thomas *et al*., 2007). Hence, proper optimization of tillage along with crop residue retention is necessary as it provides multiple benefits *viz*. enhance soil microbial diversity, creates positive soil carbon / nitrogen stocks (Melero *et al*., 2011; Parihar *et al*., 2018), improving water use efficiency (Chauhan *et al*., 2000) and agronomic efficiency of applied fertilizer (Kamanga *et al*., 2014). Therefore, this article reviews a performance of maize in relation to crop yield, resource use efficiency and different soil properties under CA-based tillage practices.

[1]Division of Agronomy, ICAR-Indian Agricultural Research Institute (IARI), New Delhi-110 012, India.
[2]Department of Agricultural Meteorology and Physics, Bidhan Chandra Krishi Viswavidayala, Mohanpur-741 252, West Bengal, India.

Corresponding Author: Niraj Biswakarma, ICAR-Indian Agriculture Research Institute, New Delhi-110 012, India.
Email: biswaniraj666@gmail.com

How to cite this article: Biswakarma, N., Rai, B., Nayak, S. and Radheshyam (2023). Conservation Agriculture (CA)-Based Tillage Practices in Maize for Enhancing Crop Yield, Resource use Efficiency and Soil Health: A Review. Agricultural Reviews. 44(2): 245-251. doi: 10.18805/ag.R-2184.

Submitted: 05-03-2021 **Accepted:** 28-05-2021 **Online:** 19-07-2021

Principles of conservation agriculture (CA)

According to FAO conservation agriculture is based on three universally applicable and acceptable principles:

Minimum mechanical soil disturbance

This principle primarily focuses on the soil conservation. Practicing no-till system moderates SOM oxidation (Ladha *et al*., 2009; Thomas *et al*., 2007), soil water movement (Bhan and Behra, 2014), bulk density and soil resistance to penetration (Saha *et al*., 2010) besides lowering cultivation expenditure (Sharma *et al*., 2011).

Permanent soil cover through organic crop residues

In India, annual generation of crop residues is estimated

around 500 MT (GOI, 2016). Thus, the surplus residues can be recycled after being used for feeding cattle, cooking fuel and organic manure production (Devi *et al.,* 2017). Residue retention creates a favorable crop micro-climate, act as safeguard against beating action of rains (Pimentel *et al.*, 1995), reducing surface evaporation (Tolk *et al.,* 1999) and leads to improvement in soil physico-chemical and biological properties (Jat *et al.*, 2014). Recently, crop residue management using cover crops under CA is gaining much popularity across the globe (Teasdale and Sharley, 1998; Kuo *et al.*, 1997).

Crop diversification

Diversifying crops fulfills human dietary requirement (Bhan and Behra, 2014), stabilized income / crop productivity (Das *et al.,* 2014), provides protection against pest / disease / weed infestation (Howard *et al.,* 2003) and also improves soil productivity (Stagnari *et al.,* 2009). In this regard, the Government of India (GOI) had also initiated several schemes for promoting sustainable crop diversification to address the burning issue of agriculture and allied sectors.

Impact of CA-based tillage practices

Yields levels

Globally, ever–growing population and shrinking land resources has threatened sustainable crop production. Hence, sound agronomic intervention plays an important role to ensure food and nutritional security. The CA-based tillage practices has come up to achieve maximum production potential through sustained agronomic productivity. The study (six year) conducted by (Parihar *et al.* 2016), in NW-IGP of India, had documented 9.4-12.2% greater pooled grain yield and highest system productivity (11.3–12.9 Mg ha^{-1}) with CA-based tillage practices. Several other studies also conform the higher maize yield ranged from 17-19% with CA-based practices (Bachmann and Friedrich. 2002; Blanco-Canqui *et al.,* 2006; Kaschuk *et al.*, 2010; Das *et al.*, 2014; Singh *et al.*, 2014). While, Jat *et al.* (2006) reported marginally lower maize yield under zero-tillage system due to more infestation of perennials weeds. The maize planted on permanent-bed resulted in enhanced maize yields compared CT practices (Potter *et al.*, 1996). Likewise, Aquino *et al.* (1998) had also reported yield advantage of wheat, maize and soybean under raised bed planting. Which was in consistent with the investigation of Jat *et al.* (2013) in which maize on raised bed provides substantial yield advantage up to 24-40% compared to CT based practices. Similarly, Ram *et al.* (2010); Jat *et al.*, (2006); Kumar *et al.* (2004); Govaerts *et al.* (2005); Thierfelder *et al.* (2018) concluded that higher maize yield under CA based system might be attributed to better water regimes, compound effects of additional supply of nutrients and improvement in soil physical properties, improved soil carbon status resulted from residue imbedded tillage with practices.

Resource use efficiencies

The inefficient resource use in conventional practices leading to declining factor productivity and low input use efficiency. Several studies conducted in past demonstrated the effects of CA based tillage practices exclusively on input use efficiencies such as higher water use efficiency (Chauhan *et al.*, 2000), higher energy output, increased energy productivity and net energy (Parihar *et al.*, 2017; 2018). Similarly, Jain *et al.* (2007) also reported higher energy–use efficiency and energy productivity under conservation tillage relative to traditional practices. Out of total operational energy, 1/3rd of energy and 36 liters ha^{-1} of diesel (equivalent to minimizing CO_2 emission of 93 kg ha $year^{-1}$) could be saved under ZT system compared to CT based practices (Srivastava, 2003; Erenstein and Laxmi, 2008). Similarly, Jat *et al.* (2014) had reported significantly (*P<0.05*) higher carbon sustainability index (CSI) and carbon efficiency (CE) with permanent bed planting followed by zero in maize-maize-sesbania system and they concluded as did by Dubey and Lal, (2009). However, Kamanga *et al.* (2014); Six *et al.* (2004) had registered contrast results regarding nutrient use efficiency under CA-based tillage practices.



(Sources: Stagnari *et al.*, 2009).

Fig 1: The three interlinked principles of conservation agriculture (CA) and the major practices and means needed to achieve respective principle.

Soil chemical properties (pH, SOC, N, P and K)

Soil chemical environment can be modified through different agronomic practices and maintenance is equally important to achieve higher crop growth and productivity. The CA-based tillage *viz*. ZT and PNB had a remarkable positive impact on the chemical properties as it preserves organic matter content and improves status of different nutrient in the soil (Wright and Hons, 2004; Owens *et al*., 2002). Similarly, Govaerts *et al.* (2007) observed linear relationship of soil organic matter with an amount of straw retention under permanent raised bed system. Similar findings were also obtained by Lal, (1997); Hulugalle and Entwistle (1997); Utomo *et al*., (2013). Likewise, Parihar *et al.* (2016) had reported 23.6-35.3% higher soil carbon content at 0-15 cm soil depth under CA based tillage pertained to CT practices. However, several other studies confirmed higher SOC ranged from 19.4 to 67% with CA based practices at 0-30 cm soil section compared to maize planted on CT plots (Thomas *et al*., 2007; Kaiser *et al*., 2014; Six *et al*., 2004; Das *et al*., 2014) (Table 1). The adoption of CA-based tillage in conjugation with crop residues residue retention hold potential to influences soil total and mineralizable-N content (Gosai *et al*., 2009), available soil nitrogen status (Mohanty and Mishra, 2014), extractable-P content and K content in the surface soil layer (Ismail *et al*., 1994). Which was ascribed to less leaching, fixation and release of several chelating agents resulted from improvement in soil properties due to optimization of tillage and crop residue management under CA based cropping system (Betrol *et al.,* 2007; Du Preez *et al.,* 2001; Robbins and Voss 1991; Standley *et al.,* 1990).

Soil physical properties

Tillage practices have a direct impact on soil physical health. It modifies water regimes, root proliferation/penetration into the soil and controlled the absorption of soil nutrient by the plants. The ZT technology proved to be a wise choice to improve the overall soil physical health. Several findings suggested that maize when planted under no-tilled plots had shown a significant ($P<0.05$) improvement in soil bulk density, infiltration rate, hydraulic conductivity and penetration resistance and other related physical properties relative to CT based system (Wilkins *et al*., 2002; Afzalinia and Zabihi, 2014; Lampurlanes and Cantero-Martinez, 2006; Lal, 2000; Jat *et al*., 2005; Parihar *et al*., 2016). However, some researchers had reported a non-significant results with respect to soil bulk density with no-tillage system (Gwenzi *et al*., 2009; Horn, 2004; Ram *et al*., 2002; Kumar *et al*., 2002). Similarly, Yang and Wander, (1999) and Salem *et al*., (2015), concluded that changes in soil physical properties is mainly attributed to lesser trafficking of machinery along with proper crop residue management under zero tillage in comparison with CT practices.

Soil biological properties

The soil has the most diversified population of flora and fauna and it has been estimated that the 1 gram of soil contains millions of living organism like, bacteria, fungi, mycorrhiza, protozoa, nematodes, earthworms, mites, enchytraeids ants, termites, beetles and spiders etc. (Hawksworth, 1991). Higher soil microbial population and soil enzymatic activities are supposed to be key factors for nutrient cycling in an agro-ecosystem (Dick, 1994; Hungria *et al*., 2009; Salinas-Garcia *et al*., 2002). Long-term adoption of resource conservation practices along with residue retention often increase microbial populations, biomass and enzymatic activities by providing substrates for their growth (Stott *et al*., 2009; Singh *et al*., 2009), which in turn nourishes

Table 1: Impact of CA-based tillage on soil organic carbon (SOC) under maize-based rotation.

Tillage practices	Cropping system	Study durations (years)	Depth (cm)	% Increase in SOC over CT	References
PB + R	WMCS	3	0–30	19.4	Das *et al.* (2018)
ZT–DSR/M	RMCS	5	0–30	25.8	Singh *et al.* (2016)
SSM	WMCS	8	0–20	5.65	Shao *et al.* (2016)
PB	MBCS	5	0–30	19.6–37.5	Parihar *et al.* (2018)
ZT+ R	MWMS	6	0–15	67	Jat *et al.* (2019)

Note: PB + R= Permanent bed with residue; ZT–DSR/M= Zero till direct seeded rice/maize; SSM= Sub soiling mulching; ZT+ R= Zero tillage with residue; WMCS= Wheat-maize cropping system; RMCS= Rice-maize cropping system; MBCS= Maize based cropping system; MWMCS= Maize-wheat-mungbean system and MWCS= Maize-wheat cropping system; CT= Conventional tillage.

Table 2: Impact of CA-based tillage on soil microbial properties under maize-based rotation.

Tillage practices	Cropping system	Study durations (years)	Depth (cm)	% Increase in MBC over CT	% Increase in DHA over CT	References
PB	MBCS	6	0-5	46.6	57.3	Parihar *et al.* (2018)
ZT+R	MWCS	3	0-10	56	154.8	Choudhary *et al.* (2018)
NT	MCCS	4	0-10	80	-	Wright *et al.* (2005)

Note: PB= Permanent bed; ZT+R= Zero tillage with residue; NT= No till; MBCS= Maize based cropping system; MWCS= Maize-wheat cropping system; MCCS= Maize-cotton cropping system; MBC= Microbial biomass carbon and DHA= Dehydrogenase activity.

soil through nutrient recycling back to the soil system (Marinori *et al*., 2000; Singh *et al*., 2009). It has been documented that CA-based maize rotation can increase soil microbial biomass carbon (SMBC) and dehydrogenase activity up to 25-80% and greater than 100% compared to CT practices (Table 2). The undisturbed soil, more vertical distribution of organic residues, improved aeration and water content are responsible for higher enzymatic activity under zero tillage system. (Melero *et al.,* 2011; Green *et al.,* 2007; Roldan *et al*., 2007; Ladd, 1985). The long-term study conducted by Parihar *et al.,* (2016) had also reported higher (on average 45-48.9%) improvement in soil microbial biomass carbon (MBC) and soil enzymatic activities *i.e.,* fluorescine diacetate, dehydrogenase, b glucosidase and alkaline phosphatase at 0-30 cm soil depth under CA based systems. Similarly, Kladivko, (2001) suggested that, microbial diversity is negatively correlated with the intensity of tillage. Which was in similar line with the results of Ferrerus *et al*. (2000) and Hassen *et al*. (2007).

CONCLUSION

Currently, Indian agriculture is reeling under the myriad of problems such as adversary of climate change, yield stagnation, high cost of cultivation, soil health degradation etc. This situation demands a paradigm shift in farming practices, conservation agriculture concept, if followed in true spirit, could be a panacea for the said problems. Further, it has been proved that the CA-based maize rotation are the better alternative for the prevalent resource inefficient traditional cropping system. It can be concluded that conservation based tillage practices (ZT and permanent bed) in long run found to be suitable cultivation practice to obtain better maize productivity, improving use efficiencies and overall soil health.

REFERENCES

Afzalinia, S. and Zabihi, J. (2014). Soil compaction variation during corn growing season under conservation tillage. Soil and Tillage Research. 137: 1-6.

Aquino, P. (1998). The Adoption of Bed Planting of Wheat in the Yaqui Valley, Sonora, Mexico. Wheat Special Report 17A-CIMMYT, Mexico. pp. 37.

Bachmann, T.L. and Friedrich, T. (2002). Conservation Agriculture in Mongolia. In: Proceedings of International Symposium on Conservation Agriculture for Sustainable Wheat Production in Rotation with Cotton in Limited Water Resource Area. Institute of Engineers for the Irrigation and Mechanization of Agriculture, Tashkent, Uzbekistan. pp. 91-96.

Betrol, I., Engel, F.L., Mafra, A.L., Betrol, O.J. and Ritter, S.R. (2007). Phosphorus, potassium and organic carbon concentrations in runoff water and sediments under different soil tillage systems during soybean growth. Soil and Tillage Research 94: 142-150.

Bhan, S. and Behera, U.K. (2014). Conservation agriculture in India -Problems, prospects and policy issues. International Soil and Water Conservation Research. 2(4): 1-12.

Blanco-Canqui, H., Lal, R., Post, W.M., Izaurralde, R.C. and Owens, L.B. (2006). Rapid changes in soil carbon and structural properties due to stover removal from no-till corn plots. Journal of Indian Society of Soil Science. 171: 468-482.

Chauhan, D.S., Sharma, R.K., Tripathi, S.C., Kharub, A.S. and Chhokar, R.S. (2000). Wheat cultivation after rice- A paradigm shift in tillage technology. Indian Farming. 50(6): 21-23.

Das, T.K., Bhattacharyya, R., Sudhishri, S., Sharma, A.R., Saharawat, Y.S., Bandyopadhyay, K.K., *et al.* (2014). Conservation agriculture in an irrigated cotton-wheat system of the western Indo-Gangetic Plains: Crop and water productivity and economic profitability. Field Crops Research. 158: 24-33.

Das, T.K., Saharawat, Y.S., Bhattacharyya, R., Sudhishri, S., Bandyopadhyay, K.K., Sharma, A.R. and Jat, M.L. (2018). Conservation agriculture effects on crop and water productivity, profitability and soil organic carbon accumulation under a maize-wheat cropping system in the North-western Indo-Gangetic Plains. Field Crops Research. 215: 222-231.

Devi, S., Gupta, C., Jat, S.L. and Parmar, M.S. (2017). Crop residue recycling for economic and environmental sustainability: The case of India. Open Agriculture. 2: 486-494.

Dick, R.P. (1994). Soil Enzymatic Activities as Indicators of Soil Quality. Defining Soil Quality for a Sustainable Environment. [Doran, J.W., Coleman, D.C., Bezdicek, D.F. and Stewart, B.A. (Eds.)], pp. 107-124.

Du Preez, C.C., Steyn, J.T. and Kotze, E. (2001). Long-term effects of wheat residue management on some fertility indicators of a semi-arid Plinthosol. Soil and Tillage Research. 63: 25-33.

Dubey, A. and Lal, R. (2009). Carbon footprint and sustainability of agricultural production systems in Punjab, India and Ohio, USA. Journal of Crop Improvement. 23: 332-350.

Erenstein, O. and Laxmi, V. (2008). Zero tillage impacts in India's rice-wheat systems: A review. Soil and Tillage Research. 100: 1-14.

Ferreras, L.A., Costa, J.L., Garcia, F.O. and Pecorari, C. (2000). Effect of no-tillage on some soil physical properties of a structural degraded Petrocalcic Paleudoll of the southern 'Pampas' of Argentina. Soil Tillage Research. 54: 31-39.

GOI (2020-21). Directorate of Economics and Statistics, Department of Agriculture, Cooperation and Farmer's Welfare, government of India.

GOI Annual Report (2016). Ministry of New and Renewable Energy, New Delhi, (http://mnre.gov.in).http://www.erewise.com/current-affairs/biomass-resources).

Gosai, K., Arunachalam, A. and Dutta, B.K. (2009). Influence of conservation tillage on soil physico-chemical properties in a tropical rainfed agricultural system of northeast India. Soil and Tillage Research. 105: 63-71.

Govaerts, B., Sayre, K.D. and Decker, J. (2005). Stable high yields with zero tillage and permanent bed planting. Field Crops Research. 94: 33-42.

Govaerts, B., Sayre, K.D., Lichter, K., Dendooven, L. and Deckers, J. (2007). Influence of permanent bed planting and residue management on physical and chemical soil quality in rainfed maize-wheat system. Plant and Soil. 291: 39-54.

Green, V.S., Stott, D.E., Cruz, J.C. and Curi, N. (2007). Tillage impacts on soil biological activity and aggregation in a Brazilian CerradoOxisols. Soil and Tillage Research. 92: 114-121.

Gwenzi, W., Gotosa, J., Chakanetsa, S. and Mutema, Z. (2009). Effects of tillage systems on soil organic carbon dynamics, structural stability and crop yields in irrigated wheat (*Triticum aestivum*)-cotton (*Gosspium hirsutum*) rotation in semi-arid Zimbabwae. Nutrient Cycling in Agroecosystems. 83: 211-221.

Hassen, A., Gizachew, L., Rethman, N.F.G. and Van Niekerk, W.A. (2007). Influence of under sowing perennial forages in maize on grain, fodder yield and soil properties in the sub-humid region of western Ethiopia. African Journal of Range and Forage Science. 24(1): 35-41.

Hawksworth, D.L. (1991). The fungal dimension of biodiversity: Magnitude, significance and conservation. Mycological Research. 95(6): 641-655.

Horn, R. (2004). Time dependence of soil mechanical properties and pore functions for arable soils. Soil Science Society of America Journal. 68: 1131-1137.

Howard, J., Crawford, E., Kelly, V., Demeke, M. and Jeje, J. (2003). Promoting high-input maize technologies in Africa: The Sasakawa-Global 2000 experience in Ethiopia and Mozambique. Food Policy. 28(4): 335–348.

Hulugalle, N.R. and Entwistle, P. (1997). Soil properties, nutrient uptake and crop growth in an irrigated vertisol after nine years of minimum tillage. Soil and Tillage Research. 42: 15-32.

Hungria, M., Franchini, J.C., Junior, O.B., Kaschuk, G. and Souza R.A. (2009). Soil microbial activity and crop sustainability in a long-term experiment with three soil-tillage and two crop-rotation systems. Applied Soil Ecology. 42: 288-296.

Ismail, I., Blevins, R.L. and Frye, W. (1994). Long-term no-tillage effects on soil properties and continuous corn yields. Soil Science Society of America Journal. 58: 193-198.

Jain, N., Jain, V., Mishra, J.S. and Kewat, M.L. (2007). Effect of tillage packages and herbicides on energy and economics of wheat in transplanted rice (*Oryza sativa*) wheat (*Triticum aestivum*) system. Indian Journal of Agricultural Science. 77(3): 174-176.

Jat, M.L., Gathala, M.K., Saharawat, Y.S., Tetarwal, J.P., Gupta, R. and Yadvinder, S. (2013). Double no-till and permanent raised beds in maize-wheat rotation of north-western Indo-Gangetic plains of India: Effects on crop yields, water productivity, profitability and soil physical properties. Field Crops Research. 149: 291-299.

Jat, M.L., Satyanarayana, T., Majumdar, K., Parihar, CM., Jat, S.L., Tetarwal, R.K. and Saharawat, Y.S. (2013). Fertilizers best management practices for maize systems. Indian Journal of Fertilizers. 9(4): 80-94.

Jat, M.L., Sharma, S.K., Rai, H.K., Srivastava, A. and Gupta, R.K. (2006). Effect of tillage on performance of winter maize in northern India. Paper Presented in 6th Triennial Conference of Maize Association of Australia. Griffith, NSW, Australia.

Jat, M.L., Srivastva, A., Sharma, S.K., Gupta, R.K., Zaidi, P.H. and Srinivasan, G. (2005). Evaluation of maize-wheat cropping system under double-no-till practice in Indo-Gangetic plains of India. Paper presented in the 9th Asian Regional Maize Workshop, Beijing, China.

Jat, S.L., Parihar, C.M., Singh, A.K., Jat, M.L. and Jat, R.K. (2014). Carbon sustainability and productivity of maize-based cropping system under conservation agriculture practices in Indo-Gangetic plains. Paper presented in 12th Asian Conference and Expert Consultation on Maize for Food, Feed, Nutrition and Environment.

Kaiser, M., Piegholdt, C., Andruschkewitsch, R., Linsler, D., Koch, H. and Ludwig, B. (2014). Impact of tillage intensity on carbon and nitrogen pools in surface and sub-surface soils of three long-term field experiments. European Journal of Soil Science. 65(4): 499-509.

Kamanga, B.C.G., Waddington, S.R., Whitbread, A.M., Almekinders, C.J.M. and Giller, K.E. (2014). Improving the efficiency of use of small amounts of nitrogen and phosphorus fertiliser on smallholder maize in central Malawi. Experimental Agriculture. 50(2): 229-249.

Kaschuk, G., Alberton, O. and Hungria, M. (2010). Three decades of soil microbial biomass studies in Brazilian ecosystems: lessons learned about soil quality and indications for improving sustainability. Soil Biology and Biochemistry 42: 1-13.

Kladivko, E.J. (2001). Tillage systems and soil ecology. Soil Tillage Research. 61: 61-76.

Kumar, S., Pandey, D.S. and Rana, N.S. (2004). Effect of tillage, rice residue and nitrogen management practices on yield of wheat (*Triticum aestivum*) and chemical properties of soil under rice (*Oryza sativa*)-wheat system. Indian Journal of Agronomy. 49(4): 223-225.

Kuo, S., Sainju, U.M. and Jellum, E.J. (1997). Winter cover crops effects on soil organic carbon and carbohydrate in soil. Soil Science Society of America Journal. 61: 145-152.

Ladd, J.N. (1985). Soil Enzymes. In: Soil Organic Matter and Biological Activity, Dordrecht.

Ladha, J.K., Kumar, V., Alam, M., Sharma, S., Gathala, M.K., Chandna, P., Saharawat, Y.S. Balasubramanian, V. (2009). Integrating crop and resource management technologies for enhanced productivity, profitability and sustainability of the rice-wheat system in South Asia. In: [Ladha, J.K., *et al.* (Eds.)], Integrated crop and resource management in the rice-wheat system of South Asia. IRRI, Los Banos, Philippines pp. 69-108.

Lal, R. (1997). Long-term tillage and maize monoculture effects on a tropical alfisol in western Nigeria: Crop yield and soil physical properties. Soil and Tillage Research. 42: 145-160.

Lal, R. (2000). Mulching effects on soil physical quality of an Alfisol in western Nigeria. Land Degradation and Development 11: 383-392.

Lampurlanes, J. and Cantero-Martinez, C. (2006). Soil bulk density and penetration resistance under different tillage and crop management systems and their relationship with barley root growth. Agronomy Journal. 95: 526-536.

Melero, S., Lopez-Bellido, R.J., Lopez-Bellido, L., Munoz-Romero, V., Moreno, F. and Murillo, J.M. (2011). Long-term effect of tillage, rotation and nitrogen fertilizer on soil quality in a Mediterranean Vertisol. Soil and Tillage Research. 114: 97-107.

Mohanty, A. and Mishra, K.N. (2014). Influence of conservation agriculture production system on available soil nitrogen, phosphorus, potassium and maize equivalent yield on a fluventic haplustepts in the north central plateau zone of Odisha. Trends in Biosciences Journal. 7(23): 3962-3967.

Owens, L.B., Malone, R.W., Hothem, D.L., Starr, G.C. and Lal, R. (2002). Sediment carbon concentration and transport from small watersheds under various conservation tillage. Soil Tillage Research. 67: 65-73.

Parihar, C.M., Jat, S.L., Singh, A.K., Kumar, B., Singh, Y., Pradhan, S., Pooniya, V., Dhauja, A., Chaudhary, V., Jat, M.L., Jat, R.K. and Yadav, O.P. (2016). Conservation agriculture in irrigated intensive maize-based systems of north-western India: Effects on crop yields, water productivity and economic profitability. Field Crops Research 193: 104-116.

Parihar, C.M., Jat, S.L., Singh, A.K., Majumdar, K., Jat, M.L., Saharawat, Y., Pradhan, S., Kuri, B.R. (2017). Bio-energy, water use efficiency and economics of maize-wheat-mungbean system under precision-conservation agriculture in semi-arid agro-ecosystem. Energy. 119: 471-473.

Parihar, C.M., Parihar, M.D. Sapkota, T.B., Nanwal, R.K., Singh, A.K., Jat, S.L., Nayak, H.S., Mahala, D.M., Singh, L.K., Kakraliya, S.K., Stirling, C.M. and Jat, M.L. (2018). Long-term impact of conservation agriculture and diversified maize rotations on carbon pools and stocks, mineral nitrogen fractions and nitrous oxide fluxes in inceptisol of India. Science of the Total Environment. 1382-1392.

Parihar, C.M., Yadav, M.R., Jat, S.L., Singh, A.K., Kumar, B., Pooniya, V., Pradhan, S., Verma, R.K., Jat, M.L., Jat, R.K., Parihar, M.D., Nayak, H.S. and Saharawat, Y.S. (2018). Long-term conservation agriculture and intensified cropping systems: Effects on growth, yield, water and energy-use efficiency of maize in north-western India. Pedosphere 28(6): 952-963.

Pimentel, D.C., Harvey, P., Resosudarmo, K., Sinclair, D., Kurz, M., McNair, S., Crist, L., Shpritz, L., Fitton, R., Saffouri, and Blair, R. (1995). Environmental and economic costs of soil erosion and conservation benefits. Science. 267: 1117-1123.

Potter, K.N., Morrison, J.E. and Torbent, H. A. (1996). Tillage intensity effects on corn and grain sorghum growth and productivity on a vertisols. Journal of Production Agriculture. 9: 385-390.

Ram, H., Kler, D.S., Singh, Y. and Kumar, K. (2010). Productivity of maize (*Zea mays*)-wheat (*Triticum aestivum*) system under different tillage and crop establishment practices. Indian Journal of Agronomy. 55(3): 185-190.

Ram, H., Singh, Y., Saini, K.S., Kler, D.S., Timsina, J. and Humphreys, E.J. (2002). Agronomic and economic evaluation of permanent raised beds, no tillage and straw mulching for an irrigated maize-wheat system in north-west India. Experimental Agriculture. 48(1): 21-38.

Robbins, S.G. and Voss, R.D. (1991). Phosphorus and potassium stratification in conservation tillage systems. Journal of Soil and Water Conservation. 46: 298-300.

Roldan, A., Salinas-Garcia, J.R., Alguacil, M. and Caravaca, F. (2007). Soil sustainability indicators following conservation tillage practices under subtropical maize and bean crops. Soil and Tillage Research. 93: 273-282.

Saha, S., Chakraborty, D., Purakayastha, T. and Kalra, N. (2010). Effect of tillage and residue management on soil physical properties and crop productivity in maize (*Zea mays*)-Indian mustard (*Brassica juncea*) system. Indian Journal of Agricultural Sciences. 80(8): 679-685.

Salem, H.M., Valero, C., Munoz, M.A. Rodriguez, M.G. and Silva, L.L. (2015). Short–term effects of four tillage practices on soil physical properties, soil water potential and maize yield. Geoderma. 237: 60-70.

Salinas-Garcia, J.R., Velazquez-Garcia, J.D., Gallardo-Valdez, A., Diaz-Mederos, P., Caballero-Hernandez, F., Tapia-Vargas, L.M. and Rosales-Robles, E. (2002). Tillage effects on microbial biomass and nutrient distribution in soils under rain-fed corn production in central-western Mexico. Soil and Tillage Research. 66: 143-152.

Sharma, P., Abrol, V. and Sharma, R.K. (2011). Impact of tillage and mulch management on economics, energy requirement and crop performance in maize-wheat rotation in rainfed sub-humid Inceptisols, India. European Journal of Agronomy 34(1): 46-51.

Shiferaw, B., Prasanna, B.M., Hellin, J. and Bänziger, M. (2011). Crops that feed the world 6. Past successes and future challenges to the role played by maize in global food security. Food Security. 3: 307-327.

Singh, G., Marwaha, T.S. and Kumar, D. (2009). Effect of resource-conserving techniques on soil microbiological parameters under long-term maize (*Zea mays*)-wheat (*Triticum aestivum*) crop rotation. Indian Journal of Agricultural Sciences. 79(2): 94-100.

Singh, V.K., Dwivedi, B.S., Tiwari, K.N., Rani, M., Singh, S.K. and Timsina, J. (2014). Optimizing nutrient management strategies for rice-wheat system in the Indo-Gangetic Plains of India and adjacent region for higher productivity, nutrient use efficiency and profits. Field Crops Research 164: 30-34.

Six, J., Ogle, S.M., Breidt, F.J., Conant, R.T., Mosier, A.R. and Paustian, K. (2004). The potential to mitigate global warming with no-tillage management is only realized when practised in the long term. Global Change Biology 10: 155-160.

Srivastava, A.C. (2003). Energy savings through reduced tillage and trash mulching in sugarcane production. Applied Engineering in Agriculture. 19(1): 40-45.

Stagnari, F., Ramazzotti, S. and Pisante, M. (2009). Conservation agriculture: A different approach for crop production through sustainable soil and water management: A review. Springer Science+Business Media pp. 55-83.

Standley, J., Hunter, H.M., Thomas, G.A., Blight, G.W. and Webb, A.A. (1990). Tillage and crop residue management effect of Vertisols properties and grain sorghum growth over seven years in the semi-arid subtropics. Soil and Tillage Research. 18: 367-388.

Stott, D.E. andrews, S.S., Liebig, M.A., Wienhold, B.J. and Karlen, D.L. (2009). Evaluation of b-Glucosidase activity as a soil quality indicator for the soil management assessment framework. Soil Science Society American Journal. 74: 107-119.

Thierfelder, C., Baudron, F., Setimela, P., Nyagumbo, I., Mupangwa, W., Mhlanga, B., Lee, N. and Gerard, B. (2018). Complementary practices supporting conservation agriculture in southern Africa. A review. Agronomy for Sustainable Development 38: 1-22.

Thomas, G.A., Dalal, R.C. and Standley, J. (2007). No-till effects on organic matter, pH and cation exchange capacity and nutrient distribution in a Luvisol in the semi-arid subtropics. Soil and Tillage Research. 94: 295-304.

Tolk, J.A., Howell, T.A. and Evett, S.R. (1999). Effect of mulch, irrigation and soil type on water use and yield of maize. Soil and Tillage Research. 50: 137-147.

Utomo, M., Banuwa, I.S., Buchari, H. and Anggraini, Y. (2013). Long-term tillage and nitrogen fertilization effects on soil properties and crop yields. Journal Tanah Tropika. 18(2): 131-139.

Wilkins, D.E., Siemens, M.C. and Albrecht, S.L. (2002). Changes in soil physical characteristics during transition from intensive tillage to direct seeding. Transactions of the American Society of Agricultural Engineers. 45(4): 877-880.

Wright, A.L. and Hons, F.M. (2004). Soil aggregation and carbon and nitrogen storage under soybean cropping sequences. Soil Science Society America Journal. 68: 507-513.

Yang, X.M. and Wander, M.M. (1999). Tillage effects on soil organic carbon distribution and storage in a silt loam soil in Illinois. Soil and Tillage Research. 52: 1-9.

M.R, Y., C.M. Parihar, S.L., Jat, A.K., Singh, Kumar, R.. Yadav, R. K., Kuri, B.R., Parihar, M.D., Verma, A.P. and Jat, M.L. (2017). Long term effect of legume intensified crop rotations and tillage practices on productivity and profitability of maize vis-a-vis soil fertility in North-Western Indo-Gangetic Plains of India. Legume Research-An International Journal. 40: 282-290