REVIEW ARTICLE Agricultural Reviews, Volume 46 Issue 2: 272-279 (April 2025)

Role of Trisodium Citrate and Nanominerals in Mastitis Management in Dairy Animals: A Review

Chinmayee Sahu[1], Ayyasamy Manimaran[1], Arumugam Kumaresan[2], Duraisamy Rajendran[3], Muniandy Sivaram[4]

10.18805/ag.R-2612

ABSTRACT

Mastitis is the most common disease of dairy cattle across the globe including India and cause huge economic losses to dairy farmers. Mastitis is also the most common reason for antibiotic usage in dairy animals, which is an important reason for antimicrobial resistance development in human and veterinary important pathogen. Optimization of nutrition and improvement of immunity of lactating cows are important strategies to reduce the susceptibility to mastitis and thereby reduction of antibiotic use. Copper, Zinc and Manganese are important trace minerals for maintaining udder health and immunity but, their contents and bioavailability is affected by several factors. Citrate, the precursor for milk synthesis is an important determinant of udder health but, the role of citrate in udder health is not clear as milk citrate level is affected by various factors. This review discusses about the role of trisodium citrate and various forms of trace minerals in mastitis management in dairy animals.

Key words: Bioavailability, Dairy animals, Mastitis, Nano minerals, Trisodium citrate.

Mastitis is the most common and costly disease of dairy cattle across the globe including India. Among the animal diseases affecting the profitability to Indian dairy farmers and industry, mastitis is considered to be one of the most expensive diseases next to foot and mouth disease (Bardhan, 2013) through reduced milk production, milk quality, treatment and culling cost. Varshney and Naresh (2004) reported that the incidence of subclinical mastitis (SCM) is more common in India (10-50%) than clinical mastitis (CM: 1-10%) in cows and an annual loss to Indian dairy industry due to mastitis is about 2.37 thousand crore rupees of which, SCM accounted for approximately 70% of the loss. Other reports also estimated more incidence and economic loss due to SCM (58-72%; 4150-4365 crores) than CM (28-42%; 1700-3000 crores per annum) in India (Dua, 2001; Bansal and Gupta, 2009). Rathod *et al.* (2017) reported loss of INR 21,677- 88,340 per lactation due to SCM. Singh *et al*. (2014) reported more losses due to mastitis in high yielding crossbred (INR 1,314) than indigenous cows (INR 868) and buffaloes (INR 1, 272) per lactation. The importance of SCM is due to its more prevalence, it remains as the source of infection for herd mates, its longer duration of infections, further proceeds into CM, its difficulty of detection and it reduces the milk production and its quality (Tuteja *et al*., 1993; Seegers *et al*., 2003). Besides, the mastitis is the top most reason for antibiotic usage, which is an important public health concern due to antimicrobial resistance development in pathogen (Pol and Reugg, 2007). In order to reduce the antibiotic use, the optimization of housing, nutrition and improvement of immunity of lactating cows are important strategy to reduce the susceptibility to mastitis (Werven, 2018).

[1]Southern Regional Station (SRS), ICAR-National Dairy Research Institute (NDRI), Bengaluru-560 030, Karnataka, India.
[2]Theriogenology Laboratory, Southern Regional Station, ICAR-National Dairy Research Institute, Bengaluru-560 030, Karnataka, India.
[3]Division of Animal Nutrition, ICAR-National Institute of Animal Nutrition and Physiology, Bengaluru-560 030, Karnataka, India.
[4]Agricultural Statistics, Southern Regional Station, ICAR-National Dairy Research Institute, Bengaluru-560 030, Karnataka, India.

Corresponding Author: Ayyasamy Manimaran, Southern Regional Station (SRS), ICAR-National Dairy Research Institute (NDRI), Bengaluru-560 030, Karnataka, India.
Email: maranpharma@gmail.com

How to cite this article: Sahu, C., Manimaran, A., Kumaresan, A., Rajendran, D. and Sivaram, M. (2025). Role of Trisodium Citrate and Nanominerals in Mastitis Management in Dairy Animals: A Review. Agricultural Reviews. 46(2): 272-279. doi: 10.18805/ag.R-2612.

Submitted: 25-11-2022 **Accepted:** 27-03-2023 **Online:** 19-06-2023

Several studies reported the importance of Copper (Cu), Zinc (Zn) and Manganese (Mn) for maintenance of udder health and immunity in dairy animals (Sordillo *et al*., 1997; Bruno, 2010). In general, when mineral status is declined in animal body, first system to be compromised is immune system, followed by growth and fertility, before appearance of overt clinical symptoms (Wikse *et al.*, 1992). Several studies reported that supplementation of trace minerals during peripartum period resulted in improved functionality of neutrophil with lesser frequency and duration of mastitis (Mutoni *et al*., 2012; Dang *et al*., 2013). But the micronutrients contents vary in feedstuff and water; their absorption and utilization are also affected by their source

and patho-physiological status of animals (*e.g*., disease, rumen environment, mineral antagonism, *etc*.,) (Srinivas, 2012) and thus the current global recommendation of microminerals may not be sufficient (Socha *et al*., 2006). In this situation, increased level- or alternative source- of supplementation are options to increase the bioavailability; of which the former concept may not be a suitable strategy as several minerals have lesser levels of safety margin. For example, Cu toxicities reported in Jersey cattle at dietary levels of 37 ppm, three times the NRC (2001) requirement (Olson *et al*., 1999). On the other hand, nano formulations of trace minerals can effectively fulfil their requirement at reduced doses due to their more bioavailability (Rajendran *et al*., 2013).

Beside pathogenic factors, interference with availability of precursors for milk synthesis (*e.g*., citrate) affects udder health, particularly during early lactation period. Citrate is the main constituent of the buffer system in healthy udder for maintenance of milk pH at 6.5 and regulation of the homeostasis between Ca^{2+} and H^{+} ions. Citrate is also essential for sequestration of soluble Ca^{2+} in milk and thereby maintenance of milk fluidity (Shennan and Peaker, 2000). On the other hand, mastitic milk contained significantly lesser amount of citrate (Dhillon and Singh, 2013). Hence, deficiency of citrate in the udder would lead to the clumping of Ca^{2+}, which manifests as “flakes” in the mastitic milk and these flakes may cause injury and subsequent inflammatory reactions to udder parenchyma including changes in milk-blood barrier permeability. Based on this concept, several researchers evaluated the effects of trisodium citrate (TSC) supplementation on mastitic animals and found encouraging results. Dhillon *et al*. (1995) administered TSC and found decrease in bacterial count along with raised milk citric acid level and restoration of normal milk pH in dairy animals. They also found that administration of TSC replenished citrate deficiency and restored milk constituents on acute or subacute mastitic cows (Singh *et al*., 1997). TSC alone or with minerals also been recommended for prevention of mastitis in dairy animals by National Dairy Development Board (NDDB) (NDDB Annual Report, 2018). Under this program, they reported that feeding of 10 g of TSC for 10 days in feed or water for California Mastitis Test (CMT) positive animals resulted in reduction of pooled milk CMT positivity from 55% to 20% after seven rounds of TSC supplementation and testing. The reduction of CMT positivity of pooled milk sample was from 60% to 25% and 49% to 11% in cattle and buffalo, respectively (Kumar, 2016). On the other hand, several researchers reported that milk citrate level varies with stage of lactation, milk composition, season and diet and mammary epithelium is impermeable to citrate in both directions (Linzell *et al*., 1976; Holt and Muir, 1979; Braunschweig and Puhan, 1999). Therefore, it warrants further studies to understand the role of citrate in lactating dairy animals. In this review, we discussed about the role of TSC and various forms of trace minerals in mastitis management in dairy animals.

Effect of trisodium citrate on mastitis affected dairy animals

Citrate is an important constituent of milk buffer system and its fluidity. Citrate, an important intermediate molecule in tricarboxylic acid (TCA) cycle, is negatively correlated with ketone bodies level in milk and thus it can be an indicator of energy status in cows (Baticz *et al*., 2002). Milk citrate level varies with lactation stage, level of milk constituents (*e.g*., fatty acid levels), season and diet (Braunschweig and Puhan, 1999). Baticz *et al*. (2002) reported drop in citrate level during the metabolically critical period of first four weeks of lactation. In contrast, some researchers reported that variation in milk citrate with lactation stage is related to de novo synthesis of fatty acids and it is independent of diet and milk yield (Garnsworthy *et al*., 2006). Altogether, it indicated that further studies are required to understand the factors influencing the milk citrate level in lactating dairy animals.

Among the three immune responses of udder during mastitis, such as i) recruitment of PMN cells, ii) altered permeability of blood-milk barrier and iii) impaired synthetic and secretary activity of alveolar epithelial cells, one or more mechanism affects the milk composition. Normally, milk constituents like citrate, milk proteins, lactose, phosphate and calcium transportation across the mammary epithelium occurs *via* “exocytosis” process, while transportation of ions (Na^{+}, K^{+}, Cl^{-}), water and small molecules like glucose, amino acids occur *via* “trans membrane” pathways. Paracellular pathway allows direct transfer of interstitial fluid and serum components into milk via leaky tight junctions and it is activated during different physiological (*e.g*., colostrogenesis time, during late lactation or involution period, higher dose of oxytocin-mediated contraction of myoepithelial cells, *etc*.) and pathological conditions like mastitis. All these pathways are generally affected by the functional state of the mammary gland and regulated by hormones and growth factors (Shennan and Peaker, 2000; McManaman and Neville, 2003). Mammary epithelium is impermeable to citrate in both directions and thus level of milk citrate concentration is an indicator of mammary gland activity rather than general metabolism (Linzell *et al*., 1976). On the other hand, mastitic milk contained significantly lesser amount of citrate (Dhillon and Singh, 2013). Milk citrate is synthesized from glucose and acetate in mammary gland. About 130-160 mg of citrate per 100 ml of milk of cows and goats normally occur with function of maintaining lactogenesis. The increase of citrate concentration by 46 times in cows and 10 times in goats during 1-2 days prepartum to 2-3 days postpartum period, indicated the onset of secretion of milk (Peaker and Linzell, 1975). On the other hand, mastitis affected animals had lesser milk citrate concentration of about 33 mg/100 ml (Oshima and

Fuse, 1981; Dhillon *et al*., 1995). Oshima and Fuse (1981) reported decreased citrate values in SCM and the extent of decrease was proportional to the degree of mastitis. The citrate content in milk is reduced by its transfer into blood via passive diffusion pathway through paracellular route (Faulkner and Peaker, 1982). Dhillon *et al*. (1995). evaluated the effect of TSC (12 g/day for 7-10 days) on acute and sub-acute mastitis affected buffaloes and found significant reduction of bacterial colonies, pH and replenishment of milk citric acid. They also evaluated the effect of TSC (15 g for 6-8 days) on acute or sub-acute mastitis affected crossbred cows by CMT method and found improved milk citrate level, milk fat, milk protein and lactose content with simultaneous reduction of milk pH (Singh *et al*., 1997). Sarfaraz *et al*. (2009) reported that supplementation of TSC (30 mg/kg b.w. for 8 days) alone resulted 69% cure rate than TSC and levamisole combined treatment in SCM affected buffaloes. But, in term of bacteriological count, combined treatment group showed better responses (Yousaf *et al*., 2010). Some studies indicated that TSC was even superior to antibiotics in regards to the restoration of milk pH and other milk constituents in mastitis affected cows with increase of milk yield, fat, and SNF (Prakash *et al*., 2010). Sindhu *et al*. (2018) reported that milk pH and chloride levels were significantly reduced while milk citrate, lactose and calcium were improved after combined administration of TSC (30 mg/kg, five days) along with antibiotic (three to five days). Above clinical studies diagnosed the SCM and evaluated its treatment outcome often using CMT and pH strip methods, which are lesser sensitive methods. They also used various duration of TSC treatment and did not mention about role of patho-physiological factors on treatment outcome. Therefore, further studies are required to understand the therapeutic efficacy of TSC in SCM cows.

Reddy *et al*. (2017) observed that supplementation of coated TSC (20 g for 10 days, N: 48) in CMT positive cows, increased milk yield and fat content. Kumar (2016) reported increase of about one litre of milk per day and become CMT negative after TSC supplementation. Santoshi *et al*. (2018) evaluated combined supplementation of TSC (25 g) and Vitamin E (1000 IU/animal/day) in transition Sahiwal cows and found significantly higher body condition score (BCS) and dry matter intake (DMI) in treated cows. They also found more milk yield, milk fat and lactose with reduced milk SCC in supplemented cows. Stumpf *et al*. (2013) found no influence of sodium citrate (100 g/cow/day) and sodium bicarbonate ($NaHCO_3$) supplementation on the metabolic parameters, milk yield and its composition, milk SCC, milk quality, body weight and BCS of the cows. Mbonwanayo *et al*. (2016) reported no significant effects of TSC (30 mg/kg b.w. for 7 days) on daily milk yield, fat, and protein. But they found marginal decrease of milk SCC in TSC supplemented cows. Shaikh *et al*. (2019) observed that administration of TSC powder (30 mg/kg, BID for five days) was found to be more efficacious for prevention of SCM in cows maintained under conventional housing system than in loose housing system. Recently, we reported that supplementations of TSC alone cause no significant changes in milk yield and its composition, but reduced milk SCC and CMT score with more bacteriological cure rate in SCM affected cows (Sahu *et al.,* 2022).

Sodium citrate contains the sodium salt of citrate and it dissociates into sodium and citrate after its absorption. The citrate ions are further metabolized into bicarbonate ions, which act as blood buffering agent. Several studies investigated the effect of bicarbonate in dairy animals and found different results. For examples, Mckinnon *et al*. (1990) found no improvement in feed intake or milk production, fat percentage and rumen volatile fatty acid (VFA) production after bicarbonate supplementation. In contrast, Rogers *et al*. (1985) reported increase of milk yield, DMI and rumen VFA production in $NaHCO_3$ supplemented cows. Vicini *et al*. (1988) investigated the effect of bicarbonate supplementation along with *ad lib* or restricted ration and found increased DMI of cows fed ad libitum while restricted diet had lower milk yield, milk fat, protein and SNF. Feeding higher level of $NaHCO_3$ in diet not only increased DMI, milk yield, milk fat but also increased conception rate in early lactating Nili Ravi buffaloes during summer (Sarwar *et al*., 2007).

Effect of trace minerals on udder health of lactating dairy animals

Trace minerals such as Cu, Zn and Mn play an important role in maintaining the udder health and immunity of lactating dairy animals. Supplementation of Vitamin E, Se, Cu and Zn around peripartum period resulted in more rapid influx of neutrophil with its increased functionality in mammary gland and lesser duration of mastitis in Indian dairy cattle (Mutoni *et al*., 2012). Cortisol levels are also maintained at basal level in micronutrient supplemented animals (Dang *et al*., 2013). Supplementation of Vitamin E, Cu, Zn and combination of all these micronutrients significantly increased the phagocytic activity of neutrophils and lymphocytes proliferation during the peripartum period (Dang *et al*., 2013). Other studies, with Zn methionine supplementation showed reduced SCC (Kellogg, 1990). Whitaker *et al*. (1997) compared the effects of Zn (25 ppm) from organic and inorganic sources and found that source of Zn had no effect on intra-mammary infections (IMIs) and SCC. Cu supplementation during peripartum period (-60 to + 30 days) decreased the severity of *E. coli* induced mastitis and reduced the prevalence rate of IMIs due to major mastitis pathogens (Scaletti *et al*., 2003). Although, studies on the effect of Mn alone on immunological function are limited, increasing Mn status has been shown to enhance the killing ability of macrophages (Tomlinson *et al*., 2008). Since Cu, Zn and Mn are synergistically stimulating the superoxide dismutase activity and immune cell production, combined supplementation of these minerals is expected to improve the immune functions in dairy animals. For example, replacement of 30-33% inorganic

Cu, Mn and Zn with organic form resulted in 34-45% reduction in milk SCC in dairy cows (Popovic, 2004).

Several studies conducted in Indian dairy animals revealed differential effects of micronutrients. Kumar *et al*. (2009) reported that Cu supplementation (5 g/ week for 45 days as Cu sulphate) in SCM affected buffaloes before calving had no impact on recovery from SCM after calving (based on CMT), while Zn supplementation during peripartum period had some beneficial impact. However, none of the animals in both the groups developed CM. Malhotra (2011) supplemented Cu (20 ppm) during peri-partum (-60 to +60 days) period and found no significant effect on immunity, metabolic and endocrine profile of Karan Fries cows. Chandra *et al*. (2013) evaluated the effect of individual and combined supplementation of Vit. E (1,000 IU/cow/day) and Zn (60 ppm/ cow/day) during peri-partum (-60 to + 90 days) period and found better BCS, higher blood glucose and enhanced milk yield with lower non esterified fatty acid and lipid peroxidation level in combined group than the individually supplemented Sahiwal cows. Sharma *et al*. (2014) reported that nutritional supplementation ameliorated the SCM-induced metabolic, biochemical and milk compositional changes in cows. Similar to oral supplementations, injectable preparations were found to be differential effects. Warken *et al*. (2018) reported that subcutaneous mineral supplementation improved the immune response and minimized the oxidative stress in lactating dairy cows. In contrast, Ganda *et al*. (2016) reported that injectable trace minerals administration in cows with elevated SCC had no effect on milk yield, milk composition and milk SCC. Variations in beneficial effects of antioxidant minerals such as Cu, Zn and Mn supplementation in mastitic dairy cows could be due to supplementation levels, formulations, sources and subsequent differential bioavailability.

Bioavailability of trace minerals in dairy animals

Although well-balanced dairy diet contained high levels of trace minerals, it has no nutritional use, unless they are absorbed and utilized for numerous metabolic functions including immune response, growth, production and reproduction of dairy animals. Trace minerals deficiency is rarely identified as most of the time the deficiency causes subclinical infections rather than clinical manifestation at their earlier stage. Immunity and enzyme functions are first to be compromised during trace mineral status deficiency (Wikse *et al*., 1992) and thus, it is very critical to maintain the adequate level for optimal performance and health of dairy animals. Blood concentrations of micronutrients are more reliable indicator of mineral status of the cow than assessing dietary mineral status, as blood concentrations are significantly correlated with nutritional status of trace elements (Herdt and Hoff, 2011). They reported normal range for serum Cu as 0.6 to 1.1 μg/mL, Zn as 0.6 to 1.9 μg/mL and Mn as 0.9 to 6.0 ng/mL. However, several studies observed that plasma trace minerals concentrations and neutrophil functionality remain unchanged in additional supplemented group (Nemec *et al*., 2012; Dietz *et al*., 2017). Kincaid (1999) reported that liver concentration is more suitable than blood level of Cu, Zn and Mn in dairy animals.

The traditionally used inorganic mineral formulation in dairy animals has differential bioavailability due to their different solubility, molecular mass, electrical charge, pH properties and chemical reactive states that affect the absorption in the dairy cow. Besides, patho-physiological states of cows also influence the absorption. Srinivas (2012) reported variable content and bioavailability of cationic and anionic minerals (Table 1). NRC (2001) estimation also revealed absorption of 1 to 5% for Cu, 0.5 to 1% for Mn and 5 to 15% for Zn in mature cattle. With such low levels of actual absorption and variability of this mineral content in feedstuffs and rations, it is very difficult to feed the optimum requirement to dairy animals. The actual absorption is always low in oral feeding irrespective of sources due to rumen environment and other dietary antagonists. Species and breed also has some effect on trace mineral absorption (Blezinger, 2008). Kumar *et al*. (2017) also reported lesser absorption values of Mn, Cu, and Zn from maize germ oil cake in Indian dairy animals.

Absorption of Cu is much lower in ruminants than in non-ruminants, due to modifications that occur in the rumen environment. For instances, high dietary molybdenum with sulfur and iron results in thiomolybdates and copper sulfide formation in the rumen respectively, which greatly reduce Cu absorption and its metabolism. Dietary factors that affect bioavailability of Zn in ruminants are not well defined, though phytate does not seems to affect Zn absorption in ruminants because of microbial phytase enzyme mediated degradation of phytate in the rumen. Mn is very poorly absorbed in ruminants, and it could be due to high dietary calcium and phosphorus (Spears, 2003). Therefore, current dairy NRC, 2001 requirements for minerals such as Cu, Mn and Zn are may not be sufficient to meet the needs of the cow (Socha *et al*., 2006). They also studied the relationship between Mn intake and apparent Mn balance in dry and lactating dairy cows and found that feeding Mn above NRC requirement is necessary for achieving positive balance. Dose-dependent increase of Mn absorption suggested that a non-saturable paracellular absorption is

Table 1: Contents and bioavailability of trace minerals.

Minerals	Standard	Content	Bioavailability	Requirement (mg/kg DMI)
Cu	$CuSO_4$	25%	1-28%	4-15
Mn	$MnSO_4$	32%	4-12%	20-25
Zn	$ZnSO_4$	35%	10-50%	10-16

lactating cows. In the later strategy, feeding of chelated organic mineral has been practiced to increase the bioavailability of trace minerals (*e.g*., Zn-methionine, Cu-lysine, and Mn-methionine). Osorio *et al*. (2016) reported that supplementation of Zn, Mn, and Cu as amino acid complexes produced the positive response in milk yield and milk protein level with increased neutrophils and lymphocytes phagocytosis, and antioxidant capacity in dairy cows. Hackbart *et al*. (2010) reported that supplementation of organic trace minerals (Zn, Mn, Cu, and Cobalt) increased milk yield in mid-lactation, but did not affect postpartum follicular dynamics, embryo quality, or liver trace mineral concentrations. In general, it is commonly assumed that organic source is believed to improve the relative bioavailability than their inorganic counterparts (NRC, 2001). Meta-analysis study also showed that organic trace mineral supplementation could improve production and reproduction performances in lactating dairy cows (Rabiee *et al*., 2010). Apart from sources and formulations, supplementation of nanominerals to increase their bioavailability is a recently growing strategy as nanominerals have lesser antagonistic effects with other compounds at intestines with improved digestive efficiency, immunity and performance in livestock and poultry (Gopi *et al*., 2017).

Effect of nanominerals (Cu, Zn and Mn) on udder health and performance of ruminant animals

Rajendran *et al*. (2013) evaluated the effects of nano ZnO supplementation in SCM affected lactating HF crossbred cows and found increased milk yield and decreased milk SCC than other conventional formulations. The serum concentration of Zn was also found to be significantly higher in nano Zn supplemented cows than other group of cows. The results indicated that the more bioavailability and thus speedy recovery from SCM among nano Zn compared to other forms. Supplementation of nano ZnO, *in vitro*, showed an improvement in the growth of ruminal microorganisms, ruminal microbial protein synthesis and energy utilization efficiency in the early phase of incubation (Zhisheng, 2011). Cai *et al.,* (2021) reported that nano ZnO supplementation improved Zn bioavailability without affecting milk yield, health status, and mammary gland permeability in dairy cows. Nano Zn also increased blood Zn levels and immunity in Zn-deprived goats (Song *et al*., 2021). Similarly, nano Cu increased blood Cu content, antioxidant capacity, daily gain and wool yield in Cu-deprived sheep and goats (Shen *et al*., 2021; Min *et al*., 2022; Zhao *et al*., 2023). Singh *et al*. (2018) also reported that supplementation of nano ZnO significantly improved the plasma Zn levels in pre-ruminant lambs without causing toxicity. They suggested that nano ZnO formulation could be a better source of supplementation with reduced cost in future. In contrast, Zaboli *et al*. (2013) also reported either feed intake or average daily gain was not affected in kids supplemented with nano ZnO.

Abdollahi *et al*. (2020) found that nano ZnO increased the post-weaning feed intake, digestibility and blood Zn concentration compared to normal ZnO in calves. Chang *et al*. (2020) observed ZnO and Zn-methionine mediated reduction of diarrhoea incidence in dairy calves. They recommended ZnO supplementation during first 3 days of life and Zn-Met supplementation for the subsequent period. Combined supplementation of nano Cu and nano Zn improved young dairy calves' health by improving immunity and antioxidant status (Pandey *et al.,* 2022). They also found nano Cu supplementation at reduced levels, reduced oxidative stress through increased antioxidants levels and upregulation of antioxidant encoding genes and immune variables in Sahiwal heifers (Kushwaha *et al.,* 2022). Riazi *et al.* (2019) reported that the higher level of nano ZnO supplementation had no advantages on *in vitro* rumen fermentation. Several studies reported that oral supplementation of ZnO or nano ZnO had no significant effect on Zn blood level in sheep (Smith and Embling, 1993; Najafzadeh *et al*., 2013). Alijani *et al.* (2020) indicated that nano ZnO had advantages over inorganic ZnO in terms of increasing the Zn absorbability, rumen and blood antioxidant activity, blood IgG levels and decreasing blood urea nitrogen. Shafi *et al.* (2020) reported better efficacy of nano Zn on reduction of milk SCC than inorganic Zn in lactating Does.

Conclusion

The available literatures revealed that the supplementation of trisodium citrate had differential effects and further studies with more specific indicators of treatment outcome like milk SCC, bacteriological cure rate, *etc*., are required to understand the effects of trisodium citrate on udder health. Supplementation of trace minerals as organic form showed more beneficial effects than inorganic forms. Available studies revealed better efficacy of nanominerals than inorganic minerals on their bioavailability and improving udder health in dairy animals.

Conflict of interest

All authors declare that they have no conflict of interest.

References

Abdollahi, M., Rezaei, J., Fazaeli, H. (2020). Performance, rumen fermentation, blood minerals, leukocyte and antioxidant capacity of young Holstein calves receiving high-surface ZnO instead of common ZnO. Archives of Animal Nutrition. 74(3): 189-205.

Alijani, K., Rezaei, J., Rouzbehan, Y. (2020). Effect of nano-ZnO, compared to ZnO and Zn-methionine, on performance, nutrient status, rumen fermentation, blood enzymes, ferric reducing antioxidant power and immunoglobulin G in sheep. Animal Feed Science and Technology. 267: 114-532.

Bansal, B.K. and Gupta, D.K. (2009). Economic analysis of Bovine mastitis in India and Punjab- A review. Indian Journal of Dairy Science. 62: 337-345.

Bardhan, D. (2013). Economic losses due to clinical mastitis in organized dairy farms. Indian Journal of Dairy Science. 66(2): 168-172.

Baticz, O., To¨mo¨sko¨zi, S., Vida, L., Gaa´l, T. (2002). Relationship between concentration of citrate and ketone bodies in cow's milk. Acta Veterinaria Hungarica. 50: 253-261.

Blezinger, S. (2008). Trace mineral status in dairy cows: effects on health and performance. https://pdfs.semanticscholar.org/7488/214a21a6f83c929562fb5644bf8ff3bc8fbf.pdf. Accessed on July 2020.

Braunschweig, M. and Puhan, Z. (1999). Correlation between κ-casein variants and citrate content in milk quantified by capillary electrophoresis. International Dairy Journal. 9: 709-713.

Bruno, D.R. (2010). Mastitis, mammary gland immunity and nutrition. In Mid-South Ruminant Nutrition Conference, (Vol. 19).

Cai, J., Miao, C., Chen, Y., Xie, Y., Liu, J., Wang, D. (2021). Nano-sized zinc addition enhanced mammary zinc translocation without altering health status of dairy cows. Animal Nutrition. 7(4): 1024-1030.

Chandra, G., Aggarwal, A., Singh, A.K., Kumar, M., Upadhyay, R.C. (2013). Effect of vitamin E and zinc supplementation on energy metabolites, lipid peroxidation and milk production in peripartum sahiwal cows. Asian-Australasian Journal of Animal Sciences. 26(11): 1569.

Chang, M.N., Wei, J.Y., Hao, L.Y., Ma, F.T., Li, H.Y., Zhao, S.G., Sun, P. (2020). Effects of different types of zinc supplement on the growth, incidence of diarrhea, immune function and rectal microbiota of newborn dairy calves. Journal of Dairy Science. 103(7): 6100-6113.

Dang, A.K., Jamwal, M., Kaur, M., Prasad, S.P., Pal, S., De, K., Pathan, M.M., Swain, D.K., Mohapatra, S.K., Kapila, S., Kapila, R., Kaur, H., Mohanty, A.K., Prakash, B.S. (2013). Effect of micronutrient supplementation around calving on the plasma cortisol levels of Murrah buffaloes and Sahiwal and Karan Fries cows. Tropical Anima Health and Production. 45(4): 1047-1050.

Dhillon, K.S. and Singh, J. (2013). A new horizon in the pathobiology, aetiology and novel treatment of mastitis in buffalo. Buffalo Bulletin. 32(1): 26-34.

Dhillon, K.S., Singh, T.J., Sodhi, S.S., Sandhu, H.S., Dwivedi, P.N., Singh, J., Gilu, B.S. (1995). Milk bacteriology: Pre-and post-trisodium citrate mastitis treatment in buffaloes. Indian Journal of Animal Science. 65(1): 9-11.

Dietz, A.M., Weiss, W.P., Faulkner, M.J., Hogan, J.S. (2017). Short communication: Effects of supplementing diets of Holsteins with copper, zinc and manganese on blood neutrophil function. Journal of Dairy Science. 100(3): 2201-2206.

Dua, K. (2001). Incidence, etiology and estimated economic losses due to mastitis in Punjab and in India- An update. Indian Dairyman. 53(10): 41-48.

Faulkner, A. and Peaker, M. (1982). Reviews of the progress of Dairy Science: Secretion of citrate into milk. Journal of Dairy Research. 49: 159-169.

Ganda, E.K., Bisinotto, R.S., Vasquez, A.K., Teixeira, A.G.V., Machado, V.S., Foditsch, C., Bicalho, M., Lima, F.S., Stephens, L., Gomes, M.S., Dias, J.M., Bicalho, R.C. (2016). Effects of injectable trace mineral supplementation in lactating dairy cows with elevated somatic cell counts. Journal of Dairy Science. 99(9): 7319-7329.

Garnsworthy, P.C., Masson, L.L., Lock, A.L., Mottram, T.T. (2006). Variation of Milk Citrate with Stage of Lactation and De Novo Fatty Acid Synthesis in Dairy Cows. Journal of Dairy Science. 89: 1604-1612.

Goff, J.P. (2018). Invited review: Mineral absorption mechanisms, mineral interactions that affect acid–base and antioxidant status and diet considerations to improve mineral status. Journal of Dairy Science. 101: 2763-2813.

Gopi, M., Beulah, P.V., Dhinesh Kumar, R., Shanmathy, M., Prabakar, G. (2017). Role of nanoparticles in animal and poultry nutrition: Modes of action and applications in formulating feed additives and food processing. International Journal of Pharmacology. 13: 724-731.

Hackbart, K.S., Ferreira, R.M., Dietsche, A.A., Socha, M.T., Shaver, R.D., Wiltbank, M.C., Fricke, P.M. (2010). Effect of dietary organic zinc, manganese, copper and cobalt supplementation on milk production, follicular growth, embryo quality and tissue mineral concentrations in dairy cows. Journal of Animal Science. 88: 3856-3870.

Herdt, T.H. and Hoff, B. (2011). The use of blood analysis to evaluate trace mineral status in ruminant livestock. Veterinary Clinics: Food Animal Practice. 27: 255-283.

Holt, C. and Muir, D.D. (1979). Inorganic constituents of milk: I. Correlation of soluble calcium with citrate in bovine milk. Journal of Dairy Research. 46: 433-439.

Kellogg, D.W. (1990). Zinc methionine affects performance of lactating cows. Feedstuffs. 62(35): 15-28.

Kincaid, R.L. (1999). Assessment of trace mineral status of ruminants: A review. Proceedings of the American Society of Animal Science. 77(1): 1-10.

Kumar, A., Singh, R. and Kumar, R. (2009). Impact of supplementation therapy on subclinical mastitis. Indian Journal of Animal Science. 79(12): 1220-1221.

Kumar, A.V.H. (2016). A field approach to mastitis control in India. Keynote address in the "Collective Experiences" session at the 6th IDF International Mastitis Conference, Nantes, France.

Kumar, R.D., Kewalramani, N., Mani, V., Gupta, S., Parihar, D., Kujur, A.S.T. (2017). Evaluation of bioavailability of selected minerals from Maize Germ Oilcake in crossbred male calves. Indian Journal of Animal Research. 52(5): 730-734. doi: 10.18805/ijar.B-3266.

Kushwaha, R., Kumar, V., Kumar, M., Vaswani, S., Kumar, A., Choudhury, S. (2022). Nano copper supplementation increases superoxide dismutase and catalase gene expression profiles and concentration of antioxidants and immune variables in sahiwal heifers. Biological Trace Element Research. 10: 1-12.

Linzell, J.L., Mepham, T.B., Peaker, M. (1976). The secretion of citrate into milk. Journal of Physiology. 260: 739-750.

Malhotra, P. (2011). Influence of Copper supplementation on immune status, metabolic and endocrine profile of Karan Fries cows during pre- and post-partum period. MVSc thesis submitted to ICAR-National Dairy Research Institute, Karnal.

Mbonwanayo, D., Akal, C., Cedden, F., Bohloli, S. (2016). Trisodium citrate use in dairy cows and effects on daily milk yield and quality. Journal of Agricultural, Food and Environmental Sciences. 68: 34-37.

McKinnon, J.J., Christensen, D.A., Laarveld, B. (1990). The influence of bicarbonate buffers on milk production and acid-base balance in lactating dairy cows. Canadian Journal of Animal Science. 70(3): 875-886.

McManaman, J.L. and Neville, M.C. (2003). Mammary physiology and milk secretion. Advanced Drug Delivery Reviews. 55(5): 629-641.

Min, X., Yang, Q., Zhou, P. (2022). Effects of Nano-copper Oxide on antioxidant function of Copper-Deficient Kazakh Sheep. Biological Trace Element Research. 200(8): 3630-3637.

Mutoni, G., Prasad, S., De, K., Pal, S., Mukherjee, J., Kapila, S., Dang, A.K. (2012). Effect of supplementation of vitamin E, copper and zinc around peripartum on udder health, milk yield and composition of Sahiwal cows. Livestock Research for Rural Development. 24(12).

Najafzadeh, H., Ghoreishi, S.M., Mohammadian, B., Rahimi, E., Afzalzadeh, M.R., Kazemivarnamkhasti, M., Ganjealidarani, H. (2013). Serum biochemical and histopathological changes in liver and kidney in lambs after zinc oxide nanoparticles administration. Veterinary World. 6: 534-537.

NDDB. Annual Report, (2018). National Dairy Development Board, Govt. of India.

Nemec, L.M., Richards, J.D., Atwell, C.A., Diaz, D.E., Zanton, G.I., Gressley, T.F. (2012). Immune responses in lactating Holstein cows supplemented with Cu, Mn and Zn as sulfates or methionine hydroxyl analogue chelates. Journal of Dairy Science. 95: 4568-4577.

NRC, (2001). Nutrient Requirements of Dairy Cattle, 7th rev. edition. National Academy of Sciences, Washington, DC.

Olson, W.G., Murphy, M., Auza, N.J., Linn, J.G., Shaw, D. (1999). Review of Copper Toxicity in Dairy Cattle. Proceedings of Minnesota Dairy Health Conference, Bloomington, MN. 167-173.

Oshima, M. and Fuse, H. (1981). Citric acid concentration in subclinical mastitic milk. Journal of Dairy Research. 48: 387-392.

Osorio, J.S., Trevisi, E., Li, C., Drackley, J.K., Socha, M.T., Loor, J.J. (2016). Supplementing Zn, Mn and Cu from amino acid complexes and Co from cobalt glucoheptonate during the peripartal period benefits postpartal cow performance and blood neutrophil function. Journal of Dairy Science. 99(3): 1868-1883.

Pandey, P., Kumar, M., Kumar, V., Kushwaha, R., Vaswani, S., Kumar, A., Singh, Y., Shukla, P. K. (2022). The dietary supplementation of copper and zinc nanoparticles improves health condition of young dairy calves by reducing the incidence of diarrhoea and boosting immune function and antioxidant activity. Biological Trace Element Research. 10: 1-13.

Peaker, M. and Linzell, J.L. (1975). Citrate in milk: A harbinger of lactogenesis. Nature. 253(5491): 464-464.

Pol, M. and Ruegg, P.L. (2007). Treatment practices and quantification of antimicrobial drug usage in conventional and organic dairy farms in Wisconsin. Journal of Dairy Science. 90: 249-261.

Popovic, Z. (2004). Performance and udder health status of dairy cows influenced by organically bound zinc and chromium. Ph.D. Thesis, University of Belgrade.

Prakash, V., Yadav, M.P.S., Singh, S.P. (2010). Effect of different treatments of mastitis in cow and biochemical changes in their milk. Asian Journal of Animal Sciences. 4(2): 202-204.

Rabiee, A.R., Lean, I.J., Stevenson, M.A., Socha, M.T. (2010). Effects of feeding organic trace minerals on milk production and reproductive performance in lactating dairy cows: A meta-analysis. Journal of Dairy Science. 93: 4239-4251.

Rajendran, D., Kumar, G., Ramakrishnan, S., Shibi, T.K. (2013). Enhancing the milk production and immunity in Holstein Friesian cross bred cows by supplementing novel zinc oxide. Research Journal of Biotechnology. 8(5): 11-17.

Rathod, P., Shivamurty, V., Desai, A.R. (2017). Economic losses due to subclinical mastitis in dairy animals: A study in Bidar District of Karnataka. Indian Journal of Veterinary Sciences and Biotechnology. 13(1): 37-41.

Reddy, N.K., Gangaiah, M., Rao, D.B. (2017). Coated tri sodium citrate minimizes the pathogens and maximizes the milk and milk fat in sub clinical mastitis. Journal of Bio Innovation. 6(3): 386-390.

Riazi, H., Rezaei, J., Rouzbehan, Y. (2019). Effects of supplementary nano-ZnO on in vitro ruminal fermentation, methane release, antioxidants and microbial biomass. Turkish Journal of Veterinary and Animal Sciences. 43: 737-746.

Rogers, J.A., Muller, L.D., Davis, C.L., Chalupa, W., Kronfeld, D.S., Karcher, L.F., Cummings, K.R. (1985). Response of dairy cows to sodium bicarbonate and limestone in early lactation. Journal of Dairy Science. 68(3): 646-660.

Sahu, C., Manimaran, A., Kumaresan, A., Rajendran, D., Ezhuthupurakkal, P.B., Sharma, D., Dey, D.K. Sivaram, M. (2022). Supplementation of trisodium citrate improves recovery rate of subclinical mastitis in crossbred cows. Pharma Innovation Journal. SP 11(11): 1023-1030.

Santoshi, P., Oberoi, P.S., Alhussien, M.M., Dang, A.K. (2018). Combined effect of trisodium citrate and vitamin E supplementation during the transition period on body weight and other production parameters in Sahiwal cows. Indian Journal of Dairy Science. 71(1): 78-83.

Sarfaraz, I., Yousaf, A., Muhammad, G., Abbas, R.Z., Zafar, M.A., Ahmad, T. (2009). Evaluation of tri-sodium citrate alone and in combination with levamisolehcl in the treatment of sub-clinical mastitis in dairy Buffaloes. Pakistan Journal of Zoology Supplement Series. 9: 839-843.

Sarwar, M., Shahzad, M. A., Nisa, M. U. (2007). Influence of varying level of sodium bicarbonate on milk yield and its composition in early lactating Nili Ravi buffaloes. Asian-Australasian Journal of Animal Sciences. 20(12): 1858-1864.

Scaletti, R.W., Trammell, D.S., Smith, B.A., Harmon, R.J. (2003). Role of dietary copper in enhancing resistance to escherichia coli mastitis. Journal of Dairy Science. 86: 1240-1249.

Seegers, H., Fourichon, C., Beaudeau, F. (2003). Production effects related to mastitis and mastitis economics in dairy cattle herds. Veterinary Research. 34(5): 475-491.

Shafi, B.U.D., Kumar, R., Jadhav, S.E., Kar, J. (2020). Effect of zinc nanoparticles on milk yield, milk composition and somatic cell count in early-lactating barbari does. Biological Trace Element Research. 196(1): 96-102.

Shaikh, S.R., Digraskar, S.U., Siddiqui, M.F.M.F., Syed, A.M., Ajabe, J.S. (2019). Prophylactic potential of tri-sodium citrate on subclinical mastitis in cow in different housing system. International Journal of Livestock Research. 9(9): 198-206.

Sharma, D., Singh, S.K., Srivastava, M., Sachan, P., Srivastava, A., Singh, A.P. (2014). Evaluation of metabolic status and milk compositions of indigenous cattle with subclinical mastitis and its amelioration by nutritional supplementations. Indian Journal of Animal Science. 84(8): 846-850.

Shen, X., Song, C., Wu., T. (2021). Effects of Nano-copper on Antioxidant Function in Copper-Deprived Guizhou Black Goats. Biological Trace Element Research. 199(6): 2201-2207.

Shennan, D.B. and Peaker, M. (2000). Transport of milk constituents by the mammary gland. Physiological Reviews. 80(3): 925-51.

Sindhu, O.K., Baby, P.G., Narayanapillai, U., Ajithkumar, S. (2018). Effect of Tri Sodium Citrate in the Treatment of Clinical Mastitis in Cattle. International Journal of Innovative Research in Science, Engineering and Technology. 7(8): 9040-9043.

Singh, D., Kumar, S., Singh, B., Bardhan D. (2014). Economic losses due to important diseases of bovines in central India. Veterinary World. 7(8): 579-585.

Singh, K.K., Maity, S.B., Maity, A. (2018). Effect of nano zinc oxide on zinc bioavailability and blood biochemical changes in pre-ruminant lambs. Indian Journal of Animal Sciences. 88(7): 805-807.

Singh, T.J., Singh, S., Dhillon, K.S. (1997). Effect of trisodium citrate mastitis treatment on some milk constituents in cows. Indian Journal of Animal Sciences. 67(12): 1055-1056.

Smith, B.L. and Embling, P.P. (1993). Sequential changes in the development of the pancreatic lesion of zinc toxicosis in sheep. Veterinary Pathology. 30(3): 242-247.

Socha, T.M., Tomlinson, D.J., Ward, T.L. (2006). Meeting the mineral needs of dairy cows. Advances in Dairy Technology: Western Canadian Dairy Seminar.18: 213-234.

Song, C., Gan, S., He, J., Shen, X. (2021). Effects of Nano-zinc on immune function in qianbei-pockmarked goats. Biological Trace Element Research. 199(2): 578-584.

Sordillo, L.M., Shafer-Weaver, K., Derosa, D. (1997). Immunobiology of the mammary gland. Journal of Dairy Science. 80: 1851-1865.

Spears, J.W. (2003). Trace Mineral Bioavailability in Ruminants. The Journal of Nutrition. 133(5): 1506S-1509S.

Srinivas, B. (2012). Homeostasis regulation and bioavailability of cation and anion minerals in ruminants in the reproductive of functional physiological chemistry. Indian Journal of Dairy Science. 65(1): 1-7.

Stumpf, M.T., Fischer, V., Kolling, G.J., Zanela, M.B., Ribeiro, M.E.R., Abreu, A.S.D. (2013). Metabolic attributes, yield and stability of milk in Jersey cows fed diets containing sodium citrate and sodium bicarbonate. Pesquisa Agropecuária Brasileira. 48(5): 564-567.

Tomlinson, D.J., Socha, M.T., DeFrain, J.M. (2008). Role of trace minerals in the immune system. In *Proc*. Penn State Dairy Cattle Nutrition Workshop. Grantville, PA. pp. 39-52.

Tuteja, F.C., Kapur, M.P., Sharma, A., Vinayak, A.K. (1993). Studies on bovine sub clinical mastitis: Prevalence and microflora. Indian Veterinary Journal. 70: 787-791.

Varshney, J.P. and Naresh, R. (2004). Evaluation of homeopathic complex in the clinical management of udder diseases of riverine buffaloes. Homeopathy. 93(01): 17-20.

Vicini, J.L., Cohick, W.S., Clark, J.H., McCutcheon, S.N., Bauman, D.E. (1988). Effects of feed intake and sodium bicarbonate on milk production and concentrations of hormones and metabolites in plasma of cows. Journal of Dairy Science. 71(5): 1232-1238.

Warken, A.C., Lopes, L.S., Bottari, N.B., Glombowsky, P., Galli, G.M., Morsch, V.M., Schetinger, M.R.C., Da Silva, A.S. (2018). Mineral supplementation stimulates the immune system and antioxidant responses of dairy cows and reduces somatic cell counts in milk. Anais da Academia Brasileira de Ciências. 90(2): 1649-1658.

Werven, T.V. (2018). New developments in mastitis research. Proceedings of WCDS Advances in Dairy Technology. 30: 299-306.

Whitaker, D.A., Eayres, H.F., Aitchison, K., Kelly, J.M. (1997). No effect of a dietary zinc proteinate on clinical mastitis, infection rate, recovery rate and somatic cell count in dairy cows. Veterinary Journal. 153: 197-204.

Wikse, S.E., Herd, D., Field, R., Holland, P. (1992). Diagnosis of copper deficiency in cattle. Journal of the American Veterinary Medical Association (USA). 200: 1625-1629.

Yousaf, A., Sarfaraz, I., Zafar, M.A., Abbas, R.Z., Hussain, A., Manzoor, A. (2010). Effect of treatment with tri-sodium citrate alone and in combination with levamisole hcl on total milk bacterial count in dairy buffalo suffering from sub-clinical mastitis. Revista Veterinaria. 21(1): 187-189.

Zaboli, K., Aliarabi, H., Bahari, A.A., Abbasalipourkabir, R. (2013). Role of dietary nano zinc oxide on growth performance and blood levels of mineral: A study on in Iranian Angora (Markhoz) goat kids. Journal of Pharmaceutical and Health Sciences. 2(1): 19-26.

Zhao, K., Shen, X., Zhou, P.,Wu, J. (2023). Effects of Nano-Cu2O on the Productivity in the Cu-Stripped Chinese Merino Sheep. Biological Trace Element Research. 201(3): 1181-1187.

Zhisheng, C.J. (2011). Effect of nano-zinc oxide supplementation on rumen fermentation *in vitro*. Chinese Journal of Animal Nutrition. 23(8): 1415-1421.