REVIEW ARTICLE Agricultural Reviews, Volume 46 Issue 1: 109-115 (February 2025)

Crop Wild Relatives (CWRs) a Genetic Pool for Crop Improvement: A Review

Amandeep Kaur[1], Manjeet Kaur Sangha[1]

10.18805/ag.R-2613

ABSTRACT

Deleterious effects of climatic changes and diverse environmental conditions, possess a major threat to agriculture in terms of growth and yield. So, to meet these challenges, there is a need to develop new crop varieties containing traits resistant against abiotic and biotic stresses and enhanced input use efficiency. Compared with domesticated cultivars, crop wild relatives maintains much higher level of genetic diversity. The high nutrient and bioactive content of many wild edible plants make them valued contributors for balanced diet and play significant role in human nutrition. In this review article, we discussed the importance of crop wild relative in crop improvement programs by giving examples of wild relatives that have been used to enhance abiotic and biotic stress tolerance and yield in various crop species. The advancement in DNA sequence technology enables plant breeders to produce high quality food. Further, combination of re-sequencing and de novo sequencing are efficiently used to explore genomic variation in crop wild relatives and moreover the genome analysis study results in detection of beneficial alleles in wild relatives which has been lost in cultivated species. Thus, conservation of genetic pool of crop wild relatives is essential to maintain sustainability of agriculture and food production.

Key words: Biotic and abiotic stress, Crop wild relatives, Nutritional improvement.

Abbreviations: CWRs: Crop Wild Relatives; QTL: quantitative trait locus; ILs: Introgression libraries; TYLCY: Tomato yellow leaf curl virus; YVMD: Yellow vein mosaic disease; CGMMV: Cucumber green mottle mosaic virus.

Now, a day's challenges faced by a global food security are complex and compounding. The changing climatic conditions such as increasing temperature along biotic and abiotic stresses like salinity and drought condition and new emerging strains of pest and diseases possess a major threat to the agriculture in terms of growth and yield. Moreover, the process of domestication has also results in reduction of diversity in modern crops at both the local and genomic level (Kaur *et al*., 2018). For example more than half of the genetic variation has been lost in cultivated rice (Xu *et al*., 2012) and significant reduction in maize (Wright *et al*., 2005) and cultivated soybean (Zhou *et al*., 2015) has observed in comparison to its progenitors counterparts. So, to meet these challenges, there is a need to develop new crop varieties containing traits resistant against abiotic and biotic stresses such as drought and heat tolerance, increased pest and disease resistance and enhanced input use efficiency.

Crop wild relatives (CWRs) are the primary reserve of genetic variation and these have been widely underutilized in crop improvement. In the 20th century, crop wild relatives were used in sugar cane crop improvement and further their effectiveness was known in breeding programs of major crops like rice and wheat in the 1940s and 1950s (Plucknett *et al*., 2014). The genetic base of modern cultivars is narrow and this poses major hindrance for crop improvement efforts. Germplasm characterization studies have also shown that cultivated plants are less tolerant to abiotic and biotic stresses and nutritional quality when compared to wild relatives (Holland *et al*., 2009). The wild variants are stress tolerant as they have developed stress response mechanisms which may be attributed to higher content of antioxidants and polyphenols. For example the spelt varieties of wheat were depicted to be endowed with high antioxidant activity. As a result the spelt varieties required no pesticide use and lower nitrogen fertilization (Gawlik-Dziki *et al*., 2012). Some of the wild accessions also depicted higher input use efficiency and higher productivity.

Wild food can also contribute to nutrient diversity and richness. Wild edible plants are important components of global food basket and constitute a significant part of human diet. The high nutrient and bioactive content of many wild edible plants make them valued contributors for balanced diet and play significant role in human nutrition, especially as sources of phyto-nutraceuticals. Their consumption, together with staple food crops can provide a balanced and

[1]Department of Biochemistry, Punjab Agricultural University, Ludhiana-141 004, India.

Corresponding Author: Amandeep Kaur, Department of Biochemistry, Punjab Agricultural University, Ludhiana-141 004, India. Email: amandeepkaurbrar413@gmail.com

How to cite this article: Kaur, A. and Sangha, M.K. (2025). Crop Wild Relatives (CWRs) a Genetic Pool for Crop Improvement: A Review. Agricultural Reviews. 46(1): 109-115. doi: 10.18805/ag.R-2613.

Submitted: 29-11-2022 **Accepted:** 22-02-2023 **Online:** 16-03-2023

healthy diet to malnourished people. Substantial economic and nutritional gains have been achieved by consuming edible wild okra (Gutiérrez *et al*., 2017). Loskutov *et al*., (2017) reported that biochemical like fatty acids, amino acids and sugars exhibited wider range in wild species in comparison to the cultivated ones. The wild relatives of tomato depicted higher total soluble solids and ascorbic acid content than cultivated tomato (Kumar *et al*., 2019). High yielding varieties with proven tolerance to biotic and abiotic stresses, superior nutritional profiles and ability to adapt to the changing environment are needed for continued agricultural sustainability. So, Crop wild relatives (CWR) can be exploited to enhance genetic diversity in cultivated crops which offer a diverse array of traits with the potential to decrease the amount of yield loss. The wild relatives serve as signiûcant gene pool for improving the existing crop plants for adaptation, tolerance, yield and nutritional quality (Chauhan *et al*., 2021) and find application in nutraceutical, pharmaceutical, food and cosmetic industries (Singh and Abhilash, 2018). In this review article, we discussed some representative examples in which wild relatives were used against stress resistance and improving yield and nutritional quality in their cultivated descendants.

CWRs in crop improvement

Improvement in yield and quality trait factors

Over the past decades, great progress has been made to produce nutritionally improved food crops. The genomic diversity within wild species provides useful traits for improving the nutritional value of specific crops (Nawaz *et al*. 2018). Biofortified staple crops such as fruits and vegetables are generated which harbor essential micronutrients that become a very important part of human nutrition. This article explores the improvement in nutritional status of crop and its positive impact on human health. In the following paragraph, several examples of nutritionally enhanced crops will be discussed which have been developed by using biotechnological approaches.

Increase in nutritional quality is either achieved *via* altering the biochemical structure of plants or by modifying the genotypes of staple crops for example; nutritional improvement in case of potato is achieved through interspecific hybridization involving wild and cultivated species. Hameed *et al*. (2018) studied that cultivated potato *i.e*. *solanum tuberosum* contain fewer amount of protein (0.85-4.2%) lacking in essential amino acids but introgression of protein enhanced wild gene *Amaranthus hypochondriacus (AmA1)* in cultivated potato raised the protein content upto 60%. Similarly, Starch amylose content also has been increased in potato tubers by introducing recessive allele *IAm* (Increased amylose) from wild potato *solanum sandemanii* into *Solanum tuberosum* which result in generation of potato tuber with enhanced amylose content upto 28-59%. Actually IAm trait was introgressed to *S. tuberosum* cultivars by marker assisted technique to increase the amylose content. The increase in amylose content might be due to high granule associated phosphorylase 1 (Pho1) and decreased starch synthase (SS) activity (Krunic *et al*. 2018). Gur and Zamir (2004) also depicted that introgression of three independent yield-promoting genomic regions from wild species *Solanum pennellii* into *Solanum lycopersicum* lead to generation of hybrids with 50% increase in yield over the leading variety under both wet and dry field conditions that received 10% of the irrigation water. The probability of increasing seed yield by using wild relatives was seen to be unlikely. But Frey (1976) observed that crossbreeding between *Avena sterilis* and *Avena sativa* resulted in increase in seed yield by 25–30%. After this, various examples of crop wild relatives contributing to increase in yield has been reported. Recently in wild tomato, quantitative trait locus (QTL) has been identified which contribute to increase in yield.

Tomato shows genetic variability for beta carotene content that had been utilized to breed varieties with high carotene content (Chauhan *et al*., 2021). The highest levels of vitamin C, lycopene, solids contents and phenolics were found in wild relative *i.e*. *Lycopersicon pimpinellifolium* (Colak *et al*. 2020). The crossing of *Lycopersicon pimpinellifolium* with *Lycopersicon chmielewskii* resulted in 2.4% increase in solid content of the fruit reported by Esquinas-Elcázar (1981) which was equivalent to worth of estimated 250 million US$ in California. Hefferon *et al*. (2015) also studied that tomatoes have been transformed with the grape encoding gene *i.e.* stilbene synthase in presence of fruit specific promoter. They found that these transgenic tomato plants are able to accumulate trans-resveratrol-glucopyranoside and trans-resveratrol and reached up to the level of 53 µg/g upon ripening. Resveratrol is bioactive compound that can helps to improve the nutritional value of tomatoes. Crop wild relatives of tomato such as *S*. *chilense*, *S*. *peruvianum*, *S*. *habrochaites* and *S*. *pimpinellifolium* were also found to be resistant to pest and diseases and abiotic stress tolerance (Ebert and Schafleitner, 2015). Certain accessions of *S*. *Habrochaites* and *S*. *Chilense* were resistant to leaf curl begomoviruses (Hajjar and Hodgkin, 2007).

Wild germplasm has limited use in quality characteristics, because of their genetic complexity. However, wild grasses have a higher level of protein as compared to its related domesticated cultivars. Nevo (2007) also reported that durum wheat varieties were improved in protein content through crossing with wild relative, *Triticum turgidum var dicoccoides* which had high protein content. The accession FA15-3 accumulates 40% protein in presence of nitrogen. Further, FA15-3 lines of wheat have been used to transfer the *Gpc-B1* gene (for high protein) into hard red spring wheat which resulted up to 3% more protein than the parallel lines. Alleles from *Oryza rufipogon* increased grain weight in a Korean wild rice cultivar (Xie *et al*. 2006). Kaur *et al*., (2020) also studied that promoter element of wild *Oryza rufipogon* accession (IRGC89224) has lowest expression of *Phospholipase D alpha1* (OsPLDα1) enzyme which is responsible for rice bran oil rancidity. Therefore, *Oryza rufipogon* accession (IRGC89224) have been used as

potential donor in back crossing programmes to transfer the desired trait into elite cultivars of rice to improve shelf life of rice bran. High protein content of seed has also been incorporated from wild *Cajanus* species, such as *C. sericeus*, *C. scarabaeoides* and *C. albicans* into cultivated pigeonpea (Singh and Abhilash 2018). Similarly, Barabeau and Hilu (1993) reported that crop wild progenitor of fingermillet *E.corocana subsp*. *Africana* has higher value of protein which ranges from 7.5 to 11.7%. The Fe and Ca content of wild relatives were also significantly greater than its cultivated cultivars. Moreover, these wild species also found to have high lysine and other essential amino acids content. Thus, results showed that the nutritional value of fingermillet may be significantly improved by selective crossbreeding of its wild and cultivated cultivars. However, the nutritional value of grain is higher than cereals and making them essential component of balance diet.

Among grains, soybean [*Glycine max* (L.) *Merril*] is the world's most important seed legume and also considered as an oilseed crop. Although its seeds are rich source of protein (Qin *et al*., 2014), but it has lower amount of sulphur containing amino acid such as methionine, which confines their nutritional quality. However the latest studies suggested that the seed specific expression of enzyme of methionine biosynthesis pathway *i.e. Arabidopsis thaliana cystathionine ã-synthase* (*AtCGS*) give rise to higher methionine content in the seeds of three soybean cultivars, Bert and Misuzudaizu and Zigongdongdou (Cohen *et al*., 2016).

Abiotic and biotic resistance

Crop Wild relatives (CWRs) found in an extensive environment conditions act as a good source of genotypes that are adapted to new environments in agricultural production region. Adaption to new environment means adaption to biotic or abiotic stresses that can be discovered by whole genome analysis. The analysis of genomic variation in progenitors in different environmental conditions may bring better understanding to point that how plants are adapted to various climatic disparities under natural selection (Henry and Nevo, 2014). So, this idea may be defined as a way to utilize the CWRs to increase stress tolerance and thereby improve productivity in crops such as the major cereal grains, wheat, rice, maize, grain legumes and fruits.

Abiotic stress tolerance

Salinity and Drought are the two most important environmental factors that limit production of any crop (Feyisa *et al*., 2022). Effect of both stresses have been extremely observed in crops like tomatoes, soybeans, cereals and as well as in their wild relatives (Qi *et al*., 2014). Here, we discuss the utilization of CWRs for salt and drought tolerance in wheat, soybeans and barley respectively.

Wheat is the third most produced cereal after maize and rice and it is significantly affected by salinity (Mujeeb-Kazi and De Leon 2002). Continuously, efforts have been made to produce a salt tolerant wheat variety by transferring an allele TmHKT1;5-A from its wild relative *Triticum monococcum* by using crossbreeding method. Munns *et al*. (2012) suggested that TmHKT1;5-A gene reduces the sodium concentration in the plant leaves that prevents loss of yield under salinity stress and moreover results also showed that new salt tolerant variety, produce 25% greater yield as compared to its Tamaroi parent. Ebert and Schafleitner (2015) also reported that two Na+/K+ transporter gene from wild *S*. *Cheesmaniae* act as a salt stress tolerant genes.

Soybean [*Glycine max* (L.) *Merril*] can also affected by salinity stress and results upto 40% decrease in yield (Chang *et al*., 1994). A salt tolerant gene *GmCHX1* was recently recognised in the wild relative of soybean called Glycine soja (Qi *et al*., 2014). Moreover, the finding of *GmCHX1* and another salt tolerant gene, *GmSALT3* (Guan *et al*. 2014), proposed that salt tolerant alleles might have been lost in soybean during domestication. Both the salt tolerant genes (*GmCHX1* and *GmSALT3*) help in regulating ion homeostasis and production of commercial soybean varieties with improved salt tolerance. The CWR of sunflower, *Helianthus paradoxus* is used for tolerance to salinity in sunflower and *Helianthus argophyllus* for tolerance to drought stress (Miller and Seiler 2003).

Drought is another important constraint that affects growth and development of crop production. Although several drought tolerant genes have been identified in wild relatives of crops such as barley (Suprunova *et al*., 2007), rice (Quan *et al*., 2010) and tomatoes (Fischer *et al*., 2013) but the applications of these gene for improving drought tolerance have not proven as successful as expected. The development of backcross introgression libraries (ILs) by using single gene of wild parent provides a useful substitute method for the transfer of drought tolerant genes. Honsdorf *et al*. (2014) a reported that wild barley *Hordeum spontaneum* introgressed lines were used to induce drought tolerance. The introgression lines were also prepared by crossing between the German spring barley cultivar Scarlett (*Hv*) and the Israeli wild barley to improve yield, pathogen resistance and malting quality traits.

Biotic stress tolerance

Plant yield are also reduced due attack from various biotic stresses such as pathogens like bacteria, viruses, fungi, insect pests and nematodes. Even after use of limited insecticides and pesticides, these pathogens continuously causing harm to the crops because they evolve very quickly and become tolerant. To overcome this problem, plant breeders have been exploiting exotic genetic resources such as crop wild relatives by using various strategies of biotechnology to develop broad spectrum biotic stress resistant varieties (Hajjar and Hodgkin 2007).

For example bacterial leaf blight, has been one of the most widespread rice disease caused by *Xanthomonas oryzae pv*. *oryzae* (Xoo). However, introduction of new bacterial blight-resistant genes *Xa21* from wild rice variety *Oryza longistaminata A*. *Chev*. *et Roehrich* into *Oryza sativa* has proved as a significant example of using wild relatives

in cultivated rice for conferring leaf blight-resistance in rice (Zhou *et al*. 2011). Ballini *et al*. (2007) also reported that bacterial leaf blight resistance has been obtained from introgression of resistant gene *Pi33* from wild rice, *Oryza rufipogon* into rice blast resistance variety IR64. Similarly, in United State, corn blight disease in maize was resolved by the introduction of blight resistance genes from the wild Mexican maize *Tripsacum dactyloides* L. Into commercial corn lines (Zhang *et al*., 2016). Among diploid wheat *Triticum monococcum*, progenitor of common hexaploid wheat has been used to mark the traits related to resistance genes against leaf rust and powdery mildew. Using marker assisted selection, one adult plant resistance gene (APR) for leaf rust and stripe rust has been transferred from *T*. *monococcum* to bread wheat WL711 and one gene for leaf rust has been transferred to PBW343 background (Singh *et al*., 2007). In case of cultivated oats, the wild progenitor *Avena barbata* used as a gene reservoir for powdery mildew pathogen resistance (Swarbreck *et al*., 2011).

Oilseeds are also major food crops, known for their protein and oil rich properties but major drawback in oilseed production are abiotic and biotic stresses. Wild black mustard *i.e*. *Brassica nigra* represents wild gene pool to improve cultivated Brassicas against various pathogens *e.g*. interspecific allopolyploids between *B*. *rapa*, *B*. *Fruticulosa* and *B*. *nigra* aid the use of CWRs in Brassicas as bridge species (Chen *et al*., 2011*). L*. *grandiflorum* a CWR of linseed has potential for linseed bud fly and Alternaria blight (Allaby *et al*., 2005). Among CWRs of pulses, *Vignamungo var*. *sylvestris*, *V.tribolata* and *V*. *radiatavar*. *sublobata* have provided resistance to yellow mosaic virus. *V*. *vexillata* has resistance to cowpea pod sucking bug and bruchids and it is crossable with *V*. *radaita* and *V*. *unguiculata* (Kaur *et al*., 2018).

Among different vegetable crops, potato is genetically more uniform which make it more vulnerable to abiotic and biotic stresses. Late blight is one of most important disease of potato which is caused by the *Phytophthora infestans (Mont.) de Bary*. Late blight resistance genes were transferred from wild species namely *S*. *stoloniferum* and *S*. *Demissum* into cultivated potato in different parts of the world (Bethke *et al*., 2017). Many novel genes for late blight resistance, virus resistance and other useful traits are available in diploid wild potato species like *Solanum tuberosum, S. pinnatisectum*, *S*. *acaule, S*. *brachistotrichum, S*. *cardiphyllum, S*. *jamesii, S*. *stoloniferum, S*. *Polyadenium etc.* Similarly, in tomato, the improvement in biotic resistance has benefitted significantly when various traits from wild relatives were transferred into cultivated species (*Solanum lycopersicum*). These traits include resistance to bacteria, fungi, viruses, insect pests and nematodes. The wild relatives of tomato that have been used for introgression are *Solanum habrochaites*, *S. chilense*, *S. peruvianum* and *S. pimpinellifolium*. These varieties act as richest source of genetic diversity. For example, five Ty genes that exhibit wide range of tolerance to tomato yellow leaf curl virus (TYLCY) were successfully introgressed into cultivated varieties (Menda *et al*., 2014). Nowicki *et al*. (2012) reported that genetic resistance to tomato late blight has been identified in *S*. *habrochaites*, *S*. *pennellii* and *S*. *Pimpinellifolium* and three main late blight resistance genes namely *Ph-1, Ph-2, Ph-3* have been reported in four *S*. *pimpinellifolium* accessions and among these three genes, *Ph-3* is considered most effective source of late blight resistance.

In okra, yellow vein mosaic disease (YVMD) is the most serious disease among biotic stresses (Kumar and Reddy, 2015) whereas leaf hopper, shoot and fruit borer are the major insect pests. Resistance genes for YVMD, leaf hopper, shoot and fruit borer are observed in okra wild species *A*. *manihot*, *A*. *Moschatus* and *A*. *tuberculatus* respectively (Singh *et al*., 2006). Gangopadhyay *et al*. (2017) also reported *A*. *manihot*, *A*. *caillei* and *A*. *moschatus* possess resistant against YVMD, while *A*. *moschatus*, *A*. *caillei, A*. *tuberculatus*, *A*. *manihot* to leaf hopper and shoot and fruit borer. In addition to this, a single dominant resistant gene against powdery mildew (*Peronospora destructor*) was introgressed from wild onion species namely *A*. *roylei* into cultivated onion (Kim *et al*., 2016).

Among cucurbits, wild *Cucumisfigarei* exhibited absolute resistance to cucumber green mottle mosaic virus (CGMMV), Fusarium wilt and higher level of resistance to downy mildew (Kaur *et al*., 2018). The wild species *C*. *anguria* and *C*. *ficifolius* showed no symptoms and reported resistance to CGMMV virus (Ruiz *et al*., 2021). However, wild species *Cucumis hardiwickii* has been reported with strong resistance to powdery and downy mildew diseases (Pitchaimuthu *et al*., 2012).

Limitation to use CWRs within breeding programmes

There are numerous obstacles that prevent use of CWRs as a source of potential alleles that can be incorporated into elite cultivated germplasm.

1. The compatibility, relatedness and crossability of crop wild relatives to their cultivated counterparts inhibit the direct introgression of CWR trait into cultivated genotype. For example, in cotton highly disease resistant sources were identified (such as *G*. *Somalense* J.B. Hutch; *Gossypium longicalyx* J.B. Hutch and B.J.S. Lee.) but due to their genetic incompatibility and ploidy, breeders were unable to use these resources (Tirnaz *et al*., 2022).
2. Trait identification and selection may be significantly influenced by environment as there are radically different selection regimes in a wild region compared to a domesticated region and a trait that is useful in a domesticated state may not be useful in the wild and vice versa. For example, Parker *et al*. (2020) proposed tha the decreased-pod dehiscence (PD) among domesticated common bean is due to the different fitness landscape imposed by domestication, where stronger selection pressure was used against PD in the arid conditions of North Mexico compared to tropical lowlands, where environmental humidity masks susceptibility to PD and reduces selection pressure against it (Parker *et al*., 2020).
3. Even if beneficial wild derived traits are introgressed into elite material, linkage drag can often have a negative

effect on yield or yield-related traits. For example, introduction of disease resistance genes improve some resistance but are detrimental to other agronomic traits (Trinaz *et al*., 2022).

4. Furthermore, when CWR genetic material is introduced into cultivated germplasm, sterility issues can arise, most commonly at the F1 or BC1 generation (Wang *et al*. 2020; Bohra *et al*., 2022).

There are also several challenges to using CWR in breeding that have been alleviated by the availability of more genomic resources and advances in laboratory techniques such as genome editing methods (CRISPR/Cas9), marker-assisted selection techniques, or RenSeq technologies (Bohra *et al*., 2022). But lack of knowledge about gene-trait relationships in wild species, uncertainty about how allelic combinations will be expressed in different cultivated crop backgrounds and challenges with transferring genes of interest into crops is major drawback.

CONCLUSION

Crop Wild relatives (CWRs) as a source of abiotic and biotic resistance are imperative for the genetic improvement of crops in multiple ways. To cope with problems arising from environmental changes and pathogens and population growth, it is essential to use wild genetic diversity to inflate agriculture production. Developed biotechnological applications such as genomics-assisted breeding, are used for the transfer of favorable traits from wild relatives into cultivated ones, despite incompatibility barriers. Nevertheless, now the advancement in DNA sequence technology particularly the combination of resequencing and de novo sequencing is efficiently used to explore genetic variation in crop wild relatives. The conservation of the genetic pool of wild relatives of the crop will be important for providing the key agronomic traits to breeding programs. Moreover, the utilization of the biodiversity present in the wild gene reservoir will allow cultigens to adapt rapidly to varying climatic conditions and enhance agricultural production to ensure global food and nutrition security.

Conflict of interest

The author declare that there to be no conflict of interest.

REFERENCES

Allaby, R.G., Peterson, G.W., Merriwether, D.A. and Fu, Y.B. (2005). Evidence of the domestication history of flax (*Linum usitatissimum* L.) from genetic diversity of the sad2 locus. Theoretical and Applied Genetics. 11*:* 58-65.

Ballini, E., Berruyer, R., Morel, J.B., Lebrun, M.H., Nottéghem, J.L. and Tharreau, D. (2007). Modern elite rice varieties of the 'Green Revolution' have retained a large introgression from wild rice around the Pi33 rice blast resistance locus. New Phytologist. 175: 340-350.

Barbeau, W.E., and Hilu, K.W. (1993). Protein, calcium, iron and amino acid content of selected wild and domesticated cultivars of finger millet. Plant Foods for Human Nutrition. 43: 97-104.

Bethke, P.C., Halterman, D.A. and Jansky, S. (2017). Are we getting better at using wild potato species in light of new tools?. Crop Science. 57: 1241-1258.

Bohra, A., Kilian, B., Sivasankar, S., Caccamo, M., Mba, C., McCouch, S.R. and Varshney, R.K. (2022). Reap the crop wild relatives for breeding future crops. Trends in Biotechnology. 40: 412-431.

Chang, R.Z., Chen, Y.W., Shao, G.H. and Wan, C.W. (1994). Effect of salt stress on agronomic characters and chemical quality of seeds in soybean. Soybean Science. 13: 101-105.

Chauhan, A., Sharma, D. and Banyal, S.K. (2021). Potential, challenges and strategies involved in gene introgression from wild relatives of vegetable crops: A review. Agricultural Reviews. 42(4): 390-397. doi: 10.18805/ag.R-2036.

Chen, S., Nelson, M.N., Chèvre, A.M., Jenczewski, E., Li, Z., Mason, A.S. and Cowling, W.A. (2011). Trigenomic bridges for Brassica improvement. Critical Reviews in Plant Sciences. 30: 524-547.

Cohen, H., Shir, O.M., Yu, Y., Hou, W., Sun, S., Han, T. and Amir, R. (2016). Genetic background and environmental conditions drive metabolic variation in wild type and transgenic soybean (*Glycine max*) seeds. Plant, Cell and Environment. 39: 1805-1817.

Çolak, N.G., Eken, N.T., Ülger, M., Frary, A. and Doğanlar, S. (2020). Mapping of quantitative trait loci for antioxidant molecules in tomato fruit: Carotenoids, vitamins C and E, glutathione and phenolic acids. Plant Science. 292: 110393.

Ebert, A.W. and Schafleitner, R. (2015). Utilization of wild Relatives in the Breeding of Tomato and Other Major Vegetables. Crop Wild Relatives and Climate Change (John Wiley and Sons, Inc Hoboken, NJ, USA). pp. 141-172.

Esquinas-Elcazar, J.T. (1981). Genetic resources of tomatoes and wild relatives. International Board for Plant Genetic Resources (IBPGR), Rome.

Feyisa, A.S. (2022). Potential impact of using genetically modified crops in Ethiopia: A Review. Agricultural Reviews. 43(4): 419-427. doi: 10.18805/ag.R-213.

Fischer, I., Steige, K.A., Stephan, W. and Mboup, M. (2013). Sequence evolution and expression regulation of stress-responsive genes in natural populations of wild tomato. PLoS One. 8: e78182.

Frey, K.J. (1976). Plant breeding in the seventies: Useful genes from wild plant species. Egypt Journal of Genetics and Cytology. 5: 460-482.

Gangopadhyay, K.K., Singh, A., Bag, M.K., Ranjan, P., Prasad, T.V., Roy, A. and Dutta, M. (2017). Diversity analysis and evaluation of wild *Abelmoschus* species for agro-morphological traits and major biotic stresses under the north western agro-climatic condition of India. Genetic Resources and Crop Evolution. 64: 775-790.

Gawlik-Dziki, U., Sìwieca, M. and Dziki, D. (2012). Comparison of phenolic acids profile and antioxidant potential of six varieties of spelt (*Triticum spelta* L.). Journal of Agricultural and Food Chemistry. 60: 4603-4612.

Guan, R., Qu, Y., Guo, Y., Yu, L., Liu, Y., Jiang, J. and Qiu, L. (2014). Salinity tolerance in soybean is modulated by natural variation in G m SALT 3. The Plant Journal. 80: 937-950.

Gur, A., Zamir, D. and Dangl, J. (2004). Unused natural variation can lift yield barriers in plant breeding. PLoS Biology. 2: e245.

Gutiérrez, J.A., Van, T.R., Hoekstra, R. and Hintum, T.J.V. (2017). Crop wild relatives range shifts and conservation in Europe under climate change. Diversity Distribution. 23: 739-50.

Hajjar, R. and Hodgkin, T. (2007). The use of wild relatives in crop improvement: A survey of developments over the last 20 years. Euphytica. 156: 1-13.

Hameed, A., Zaidi, S.S.E.A., Shakir, S. andMansoor, S. (2018). Applications of new breeding technologies for potato improvement. Frontiers in Plant Science. 9: 925.

Hefferon, K.L. (2015). Nutritionally enhanced food crops; Progress and perspectives. International Journal of Molecular Sciences. 16: 3895-3914.

Henry, R.J. andNevo, E. (2014). Exploring natural selection to guide breeding for agriculture. Plant Biotechnology Journal. 12: 655-662.

Holland, D., Hatib, K. and Bar-Ya'akov, I. (2009). Pomegranate: Botany, Horticulture, Breeding. Horticultural Reviews. 35: 127-191.

Honsdorf, N., March, T.J., Berger, B., Tester, M. andPillen, K. (2014). High-throughput phenotyping to detect drought tolerance QTL in wild barley introgression lines. PloSone. 9: e97047.

Kaur, A., Neelam, K., Kitazumi, A., Kaur, K., Sharma, P., Mangat, G.S. and Singh, K. (2020). Novel cis-acting regulatory elements in wild *Oryza* species impart improved rice bran quality by lowering the expression of Phospholipase D alpha1 enzyme (OsPLDα1). Molecular Biology Reports.

Kaur, V., Shubha, K., Kumar, P. and Manju (2018). Role of Crop Wild relatives in crop improvement under changing climatic conditions. Crop Improvement for Sustainability. pp.13-35.

Kim, S., Kim, C.W., Choi, M.S. and Kim, S. (2016). Development of a simple PCR marker tagging the *Allium roylei* fragment harboring resistance to downy mildew (Peronospora destructor) in onion (*Allium cepa* L.). Euphytica. 208: 561-569.

Krunic, S.L., Skryhan, K., Mikkelsen, L., Ruzanski, C., Shaik, S.S., Kirk, H.G. and Blennow, A. (2018). Non GMO potato lines with an altered starch biosynthesis pathway confer increased amylose and resistant starch properties. Starch Stärke. 70: 1600310.

Kumar, P.P., Acharya, B., Dutta, S., Saha, S., Choudhury, J. and Hazra, P. (2019). Comparative expression of different characters in cultivated tomato and its wild relatives. Journal of Pharmacognosy and Phytochemistry. 8: 3410-3419.

Kumar, S. and Reddy, M.T. (2015). Morphological characterization and agronomic evaluation of yellow vein mosaic virus resistant single cross hybrids for yield and quality traits in okra [*Abelmoschus esculentus* (L.) *Moench*]. Open Access Library Journal. 2: 1.

Loskutov, I.G., Shelenga, T.V., Konarev, A.V., Shavarda, A.L., Blinova, E.V., and Dzubenko, N.I. (2017). The metabolomic approach to the comparative analysis of wild and cultivated species of oats (*Avena* L.). Russian Journal of Genetics: Applied Research. 7: 501-508.

Menda, N., Strickler, S.R., Edwards, J.D., Bombarely, A., Dunham, D.M., Martin, G.B. and Mueller, L.A. (2014). Analysis of wild-species introgressions in tomato inbreds uncovers ancestral origins. BMC Plant Biology. 14: 1-16.

Miller, J.F. and Seiler, G.J. (2003). Registration of five oilseed maintainer (HA 429-HA 433) sunflower germplasm lines. Crop Science. 43: 2313.

Mujeeb-Kazi, A. and De Leon, J.L.D. (2002). Conventional and Alien Genetic Diversity for Salt Tolerant Wheats: Focus on Current Status and New Germplasm Development. In: Prospects for saline agriculture. Springer, Dordrecht. (pp. 69-82).

Munns R., James R.A., Xu B., Athman A., Conn S.J., Jordans C. *et al*. (2012). Wheat grain yield on saline soils is improved by an ancestral Na+ transporter gene. Nature Biotechnology. 30: 360-364.

Nawaz, M.A., Yang, S.H. and Chung, G. (2018). Wild soybeans: An opportunistic resource for soybean improvement. Rediscovery of Landraces as a Resource for the Future (September).

Nevo, E. (2007). Evolution of wild wheat and barley and crop improvement: studies at the Institute of Evolution. Israel Journal of Plant Sciences. 55: 251-262.

Nowicki, M., Foolad, M.R., Nowakowska, M. and Kozik, E.U. (2012). Potato and tomato late blight caused by *Phytophthora* infestans: An overview of pathology and resistance breeding. Plant Disease. 96: 4-17.

Parker, T.A., BernyMier y Teran, J.C., Palkovic, A., Jernstedt, J. and Gepts, P. (2020). Pod indehiscence is a domestication and aridity resilience trait in common bean. New Phytologist. 225: pp.558-570.

Pitchaimuthu, M., Souravi, K., Ganeshan, G., Kumar, G.S. and Pushpalatha, R. (2012). Identification of sources of resistance to powdery and downy mildew diseases in cucumber [*Cucumis sativus* (L.)]. Pest Management in Horticultural Ecosystems. 18: 105-107.

Plucknett, D.L. and Smith, N.J. (2014). Gene banks and the world's food. Princeton University Press.

Qi, X., Li, M.W., Xie, M., Liu, X., Ni, M., Shao, G. and Lam, H.M. (2014). Identification of a novel salt tolerance gene in wild soybean by whole-genome sequencing. Nature Communications. 5: 1-11.

Qin, P., Song, W., Yang, X., Sun, S., Zhou, X., Yang, R. and Ren, G. (2014). Regional distribution of protein and oil compositions of soybean cultivars in China. Crop Science. 54: 1139-1146.

Quan, R., Hu, S., Zhang, Z., Zhang, H., Zhang, Z. and Huang, R. (2010). Overexpression of an ERF transcription factor TSRF1 improves rice drought tolerance. Plant Biotechnology Journal. 8: 476-488.

Ruiz, L., López, C., Picó, B. and Janssen, D. (2021). Resistance to cucumber green mottle mosaic virus in Cucumismelo. Plants. 10: 1077.

Singh, A. and Abhilash, P.C. (2018). Wild Relatives of Cultivated Plants in India. A Reservoir of Alternative Genetic Resources and More, AK Singh, Springer (2017) 309 pp., Price 156.99£, ISBN: 9789811051166.

Singh, B., Rai, M., Kalloo, G., Satpathy, S. and Pandey, K. (2006). Wild taxa of okra (*Abelmoschus* species): Reservoir of genes for resistance to biotic stresses. Acta Horticulturae. 752: 323-328.

Singh, K., Chhuneja, P., Ghai, M., Kaur, S., Goel, R.K., Bains, N.S. and Dhaliwal, H.S. (2007). Molecular Mapping of Leaf and Stripe Rust Resistance Genes In *T. Monococcum* and Their Transfer to Hexaploid Wheat. In Wheat Production in Stressed Environments. Springer, Dordrecht. (pp. 779-786).

Suprunova, T., Krugman, T., Distelfeld, A., Fahima, T., Nevo, E. and Korol, A. (2007). Identification of a novel gene (Hsdr4) involved in water-stress tolerance in wild barley. Plant Molecular Biology. 64: 17-34.

Swarbreck, S.M., Lindquist, E.A., Ackerly, D.D. and Andersen, G.L. (2011). Analysis of leaf and root transcriptomes of soil-grown *Avena barbata* plants. Plant and Cell Physiology. 52: 317-332.

Tirnaz, S., Zandberg, J., Thomas, W.J., Edwards, D. and Batley, J. (2022). Application of crop wild relatives in modern breeding: An overview of resources, experimental and computational methodologies. Frontiers in Plant Science. p. 4597.

Wang, M., Yang, J., Wan, J., Tao, D., Zhou, J., Yu, D. and Xu, P. (2020). A hybrid sterile locus leads to the linkage drag of interspecific hybrid progenies. Plant Diversity. 42: pp. 370-375.

Wright, S.I., Bi, I.V., Schroeder, S.G., Yamasaki, M., Doebley, J.F., McMullen, M.D. and Gaut, B.S. (2005). The effects of artificial selection of the maize genome. Science. 308: 1310-1314.

Xie, X., Song, M.H., Jin, F., Ahn, S.N., Suh, J.P., Hwang, H.G. and McCouch, S.R. (2006). Fine mapping of a grain weight quantitative trait locus on rice chromosome 8 using near-isogenic lines derived from a cross between *Oryza sativa* and *Oryza rufipogon*. Theoretical and Applied Genetics. 113: 885-894.

Xu, X., Liu, X., Ge, S., Jensen, J.D., Hu, F., Li, X. and Wang, W. (2012). Resequencing 50 accessions of cultivated and wild rice yields markers for identifying agronomically important genes. Nature Biotechnology. 30: 105-111.

Zhang, H., Li, C., Davis, E.L., Wang, J., Griffin, J.D., Kofsky, J. and Song, B.H. (2016). Genome-wide association study of resistance to soybean cyst nematode (Heteroderaglycines) HG Type 2.5. 7 in wild soybean (*Glycine soja*). Frontiers in Plant Science. 7: 1214.

Zhou, Y.L., Uzokwe, V.N., Zhang, C.H., Cheng, L.R., Wang, L., Chen, K. and Li, Z.K. (2011). Improvement of bacterial blight resistance of hybrid rice in China using the Xa23 gene derived from wild rice (*Oryza rufipogon*). Crop Protection. 30: 637-644.

Zhou, Z., Jiang, Y., Wang, Z., Gou, Z., Lyu, J., Li, W. and Tian, Z. (2015). Resequencing 302 wild and cultivated accessions identifies genes related to domestication and improvement in soybean. Nature Biotechnology. 33: 408-414.