REVIEW ARTICLE Agricultural Reviews, Volume 47 Issue 2: 227-237 (April 2026)

# Effective Microorganisms (EM): A Promising Approach to Sustainable Dairy Production: A Review

R.M.H. Shaill Rathnayke[1,2], W.A.D. Nayananjalie[2], A.M.J.B. Adikari[2], U.L.P. Mangalika[3]

10.18805/ag.RF-393

## ABSTRACT

The dairy sector is vital to global food production and the economy. In dairy farming, the cost incurred for animal nutrition accounts for approximately two-thirds of total production costs and contributes significantly to the ecological footprint. Therefore, reducing production costs while maintaining optimal animal performance is essential. Thus, alternative, sustainable feeding sources should be researched. Effective microorganisms (EM) are considered a mixture of beneficial bacterial inoculants and have the potential to be used as an alternative feeding supplement to enhance the digestibility and subsequently enhance the production. Effective microorganism technology was developed in Japan in 1980 and since then, limited studies have focused on animal nutrition. For this study, a narrative review process was employed. Google Scholar's advanced search function was employed to filter the papers, with the keywords "Effective Microorganisms" OR "EM" OR "Direct Feed Microorganisms" OR "Probiotics" AND "Dairy Cow" OR "Cow" OR "Dairy Production" AND "Cow Health" OR "Gut Health" OR "Nutrition". Recently published, peer-reviewed articles written only in English, journal articles and leading web articles and reports were selected for the synthesis of the results. Based on synthesized knowledge, EM supplementation has significantly improved nutrient digestibility, gut health, milk production and overall performance in cows while reducing manure nutrient excretion.

**Key words:** Animal welfare, Microbial synergy, Nutrient bioavailability, Rumen health, Sustainable agriculture.

In the modern world, there is a growing trend of exploring alternative feed ingredients for the dairy sector to enhance production efficiency (Vlaicu and Untea, 2024). The term "effective microorganisms (EM)" is often used to introduce consortia of different genera of beneficial microorganisms, which could help the digestion process and nutrient absorption, as well as have probiotic effects (Abraheem *et al*., 2024; Natt and Katyal, 2022; Sugandhi *et al*., 2018). Effective microorganisms contain the strains of lactic acid bacteria (LAB), yeast, actinomycetes, photosynthetic bacteria and substrates such as molasses (Ezeagu *et al*., 2023). In addition, EM could be contained with beneficial enzymes and some crude extracts. Malaysian company (EMRO, 2024) has defined "EM as a feed additive that contains only naturally occurring three major genera of microorganisms, which are yeast, LAB and photosynthetic bacteria". The fermentation process of these consortia of bacteria secretes additional beneficial substrates of bioavailable vitamins, chelated minerals and organic acids as well as antioxidants (Sawant *et al*., 2025).

Effective microorganisms have been applied in the livestock industry for purposes such as reducing odour and manure management (Tu-Kitaw *et al*., 2013). They have also been used as feed supplements to improve animal performance, including milk and egg production (Yitbarek, 2023; Kumar *et al*., 2017). Effective microorganisms can be supplied through feed, water and in some cases, via fermented products such as Bokashi.

Various interventions aimed at improving animal performance have been discussed in the literature, however, relatively few studies have focused specifically on dairy cows.

[1]Postgraduate Programme, Faculty of Agriculture, Rajarata University of Sri Lanka, Puliyankulama, Anuradhapura 50000, Sri Lanka.
[2]Department of Animal and Food Science, Faculty of Agriculture, Rajarata University of Sri Lanka, Puliyankulama, Anuradhapura 50000, Sri Lanka.
[3]Department of Animal Production and Health, Getambe, Peradeniya 20400, Sri Lanka.

**Corresponding Author:** W.A.D. Nayananjalie, Department of Animal and Food Science, Faculty of Agriculture, Rajarata University of Sri Lanka, Puliyankulama, Anuradhapura 50000, Sri Lanka.
Email: deepthin@agri.rjt.ac.lk
ORCIDs: https://orcid.org/0009-0004-2470-0412, https://orcid.org/0000-0002-5225-0074, https://orcid.org/0000-0002-7877-4597, https://orcid.org/0000-0002-4902-1908

**How to cite this article:** Rathnayke, R.M.H.S., Nayananjalie, W.A.D., Adikari, A.M.J.B. and Mangalika, U.L.P. (2026). Effective Microorganisms (EM): A Promising Approach to Sustainable Dairy Production: A Review. *Agricultural Reviews*. **47(2):** 227-237. doi: 10. 18805/ag.RF-393.

**Submitted:** 03-06-2025 **Accepted:** 24-12-2025 **Online:** 19-01-2026

Existing research has primarily focused on odour reduction, then it has expanded to one of the mitigation strategies of livestock greenhouse gas (GHG) emission (Riaz *et al*., 2024). Moreover, with its beneficial probiotic activity, EM enhance the gut health and subsequently improves the animal performance as well (Fig 1).

However, limited studies have focused on the effects of EM on dairy cattle performance, focusing on the essential areas of nutrient digestibility, manure management and gut function (Nalla *et al*., 2022). At present, the dairy sector

faces different challenges, including high feed costs, seasonal forage scarcity, poor-quality feed and climate variability (Damunupola *et al.,* 2022; Vidanarachchi *et al*., 2019). Hence, it is worth reviewing the EM technology as a sustainable approach to improve nutrient utilisation, milk production and potentially reduce environmental impact. Therefore, this review summarises the current knowledge, study gaps and suggestions for future research to optimise the use of EM in dairy production.

## Effective microorganisms on milk yield and quality in dairy farming

Milk production is the main purpose of dairy cattle farming and improvement in production over time is important for profitable dairy farming (OECD, 2025). Today, the best nutritional regime and feeding management that influences milk production is of great importance (Qin *et al*., 2025). Hence, the investigation for other sustainable alternatives for feeding is imperative (Ngesi *et al*., 2025; Chanda *et al*., 2024). Therefore, this section centres around the effect of EM on milk production, with a particular focus on nutrient digestion, absorption and their overall influence on dairy performance.

The rumen is the largest of the anterior fore-stomach compartments of ruminant herbivores and together with the reticulum, it creates a spacious fermentation vessel (Pokhrel and Jiang, 2024; Liu *et al*., 2023; Gupta *et al*., 2016). In the rumen, there are a massive population of bacteria, fungi and protozoa, which is held at ~39°C under a continuous culture system (Mayulu and Christiyanto, 2025; Zahra *et al*., 2022). As already mentioned, dietary substrates passing to the rumen are fermented by the rumen microbes and the products of fermentation, which are volatile fatty acids (VFA), are the major source of energy in cows (Dhakal *et al*., 2024; Fang *et al*., 2021) and microbial protein and gases such as methane ($CH_4$) and carbon dioxide ($CO_2$) (Iddi-Mrutu *et al*., 2023; Parchami *et al*., 2024). Addition of EM to the system improves the rumen ecosystem and microbial diversity (Gao *et al*., 2022; Lv *et al*., 2020). With the process of fermentation, it is believed that EM can improve fermentation and subsequently milk synthesis (Jiang *et al*., 2025; Marinho *et al*., 2025; Tadesse *et al*., 2024).

It was found that microbial solution supplementation enhanced the cellulolytic bacterial population and their activity in the rumen (Nair *et al*., 2025). It causes more extensive degradation of complex fibrous feed ingredients and results in greater VFA production, especially in the form of acetate and propionate (Jiang *et al*., 2025). Faster release of acetate and propionate and faster production by efficient fibre digestion subsequently increase milk fat and volume (Yang *et al*., 2024; Shi *et al*., 2023; Mitchell *et al*., 2023). Alternatively, stimulating production or supplementation of butyrate enhances the rumen epithelial development and function by facilitating the required energy for better rumen wall function and performance (Fukumori *et al*., 2022; Steele *et al*., 2016). Based on the above discussion, EM as the probiotic significantly helps to improve milk production and its quality by enhancing the availability of VFA for better cow performance (Cappellozza *et al*., 2023; Nalla *et al*., 2022).

Supplementation of EM in small ruminants has been shown to improve ruminal degradability and improve the nitrogen utilisation and microbial protein synthesis (Direkvandi *et al*., 2020). The EM affects the rumen microbial population and optimises the rumen microbial protein synthesis, which is achieved by optimising the conditions in the rumen. Produced microbial proteins are a great source of amino acids, which drives protein synthesis for milk production (Tadesse *et al*., 2024; Zeeshan *et al*., 2023). Further, studies have shown that supplementation of microbial-based feeding enhances the body weight and milk yield in cows by efficient utilisation of available energy and microbial protein, especially increasing the synthesis of casein, which is the dominant protein found in milk (Tadesse *et al*., 2024; Carresi *et al*., 2024).





**Fig 1:** The role of effective microorganisms in enhancing dairy production.

Rumen fat digestion occurs only in minute quantities because dietary fat can be detrimental to the microbes. To alleviate this problem, microbes process fats by lipolysis and biohydrogenation (Bionaz *et al*., 2020; Vasta *et al*., 2019). Lipolysis in the rumen secretes glycerol, which may be fermented by the microbes into propionates and other VFAs that may enhance milk yield and performance in cows (Yang *et al*., 2025; Kupczyñski *et al*., 2020; Torres *et al*., 2021). Such changes in energy production processes in the rumen are accomplished through intervention from efficient microorganisms that modify and change the microbial consortia within the rumen, with the end goal being to maximise milk yield (Mohanty *et al*., 2023).

As shown in Fig 2, supplementation with microbial blends including EM supports quality milk production (Polyorach *et al*., 2023) by producing beneficial enzymes, minerals and vitamins, such as B vitamins (thiamine, riboflavin, pyridoxine, *etc*.), which are crucial for fibre, protein and fat digestion and absorption (Shukla *et al*., 2024). By enhancing nutrient absorption, EM plays a crucial role in improving overall milk yield and composition.

Other than that, recent studies have shown that high-yielding and longer lifetime dairy cows reduce the ruminal bacterial diversity, due to a greater adaptation to lactation-specific diets. Thus, supplementation with direct-fed microbials such as *Lentilactobacillus buchneri, Bifidobacterium longum* and *Pediococcus pentosaceus,* improved digestibility of both crude protein and neutral detergent fibre while improving milk protein percentage through optimization of ruminal fermentation and microbial protein synthesis (Yang *et al*., 2025; Mu *et al*., 2019; Tong *et al*., 2018). This may be literary evidence that the supplementation of EM could support and improve the longer lifetime of dairy rumen microbial diversity. Moreover, supplementation with EM results in an increase in milk yield together with improved quality parameters of milk, thereby supporting longevity of production through improved interactions between the microbiome and host (Yang *et al*., 2025; Wu *et al*., 2024).

## Effective microorganisms on nutrient utilisation and digestive efficiency in dairy cows

This section explores the effect of EM on nutrient digestibility, enzyme activities and the digestion of fibre, fat and protein, as well as its role in nutrient absorption. Many studies in dairy cows and other ruminants showed that beneficial microorganisms (*e.g*., yeast, *Bacillus* spp.) interact with native rumen microbes and promote the proliferation of cellulolytic and amylolytic bacteria (such as *Fibrobacter* and *Ruminococcus*) while suppressing pathogenic and competitive microbes (Halfen *et al*., 2021; Lopreiato *et al*., 2020).

Usually, a cow's diet consists of high-fibrous feed ingredients, which are particularly difficult to digest without fermentation (Jardstedt *et al*., 2018). Supplementation with microbial additives (EM) has been shown in some studies to increase the fibrinolytic bacterial population (*e.g*., *Ruminococcus, Fibrobacter*) in the rumen, which may enhance the digestion process by accelerating the fermentation process (Wang *et al*., 2024; Wei *et al*., 2022; Mamuad *et al*., 2019). Ingested fibrous materials are fermented and broken down by the rumen native microbes (*Ruminococcus spp, Fibrobacter spp*, *etc*.) in a blend with EM by producing enzymes that break cellulose and hemicellulose and subsequently produce simple sugars such as glucose, which are the base substrates for VFA production (Firrincieli *et al*., 2024; Weimer, 2022). In anaerobic conditions in the rumen, sugars (glucose) produced from fibrous feeds are converted into pyruvate through glycolysis and then converted into different VFS (*e.g*., acetate, propionate, butyrate), which supply energy to the cows (Zhu *et al*., 2025; Hassan and Karsl, 2022). Effective microorganisms influence the process of fiber digestion by altering the rumen environment (Tadesse *et al*., 2024; Sheikh *et al*., 2018).





**Fig 2:** Biological mechanisms of effective microorganisms in improving milk yield and quality of dairy cow.

Focusing on N utilization and protein digestion, cows ingest N and protein sources from the legume forages, by-products, concentrates like soybean meal, cotton seed meal and supplemented non-protein nitrogen (NPN) compounds (Castro-Montoya *et al*., 2021; Wilson *et al*., 2020). These proteins basically can be divided into two groups; rumen degradable protein (RDP), which is crucial for producing microbial protein and rumen undegradable protein (RUP), which is bypassed through the rumen and absorbed in the small intestine (Putri *et al*., 2021; Ahvenjärvi and Huhtanen, 2018). Both are crucial in fulfilling cows' protein requirements.

By improving the rumen ecosystem and microbial population, microbial supplementation enhances the N and protein digestion process (Jiang *et al*., 2025; Cappellozza *et al*., 2023; Hassanat and Benchaar, 2021). When RDP comes from the feed materials, they are broken down into peptides, amino acids and ammonia ($NH_3$) by the microbes through the process called proteolysis (Putri *et al*., 2021; Cherif *et al*., 2018). Then the process of deamination separates the amine group (-$NH_3$) by the enzymes and subsequently produces $NH_3$ (Gervais *et al*., 2017). By utilizing these $NH_3$ microbes produce microbial proteins, which are digested and absorbed in the small intestine. Moreover, by a similar mechanism, NPNs are utilized to produce microbial protein in the rumen, which converts urea into $NH_3$ (Lima *et al*., 2023; Zurak *et al*., 2023). For the efficient production of microbial protein, EM contributes by stabilizing the rumen pH, enhancing the fermenting process by altering the whole rumen environment.

While focusing on fat digestion, it is unlike fiber and protein digestion in the rumen, excessive fats could be toxic to the rumen microbes (Lee *et al*., 2019). Dietary fats are digested by lipolysis and biohydrogenation processes with minimal intervention of rumen microbes (Huang *et al*., 2022). Through the lipolysis process, the rumen microbes break down the complex fats, such as triglycerides and glycolipids, into fatty acids and glycerol by the enzyme lipase (Hao *et al*., 2021). Then the glycerol converts into VFA through microbial fermentation, which produces energy (Cantalapiedra-Hijar *et al*., 2016). Apart from that, the microbes convert unsaturated fatty acids in feed material such as oleic acid and linoleic acids into saturated fatty acids by adding additional hydrogen ($H^+$) atoms (Wei *et al*., 2018). This conversion reduces the harmful effect which can be caused by rumen microbes (Karangiya *et al*., 2016). The small intestine is where the saturated fatty acids go next and then they get mixed with the bile and pancreatic enzymes of the pancreas, which digest them into free fatty acids and monoglycerides that can be absorbed into the blood (Kand *et al*., 2018). Further, EM supplementation increases lipolysis and biohydrogenation by optimizing conditions for fat digestion and absorption (Owens and Basalan, 2016).

As stated in Fig 3, EM supplementation increases the nutrients' digestibility and absorption through optimum utilization of feed resources while enhancing FCE.





**Fig 3:** Understanding the digestive benefits of effective microorganisms in dairy cow nutrition.

Furthermore, recent findings by Liu *et al*. (2025) have highlighted that feed sorting behaviour may alter the rumen microbial ecosystem and certain metabolisms to a degree that negatively impacts the efficiency of fermentation and nutrient utilisation, underscoring the need to stabilize specific microbial populations to enhance digestive efficiency, even with balanced rations. Moreover, when comparing Sanhe with Holstein cows, differences in rumen microbiome composition were observed, indicating that relative to breed and parity, some microbiomes matured, resulting in increased binding of ATP transporters in lactating cows. This led to the enrichment of transport and bioconversion of certain monosaccharides and amino acids, which indirectly influenced digestion and nutrient metabolism (Liu *et al*., 2025). Besides, the study conducted by Alaa *et al*. (2025), which focuses on integrated metagenomic and single-cell RNA sequencing techniques deconstructing microbial consortia, exposed certain bacterial genomes linked to core cellulose, xylan and pectin polysaccharides with prominent fibre digestion efficiency, while identifying some epithelial cell subtypes that specifically absorb and metabolise VFA. Thus, improvements in microbial ecosystem functionality are able to maximise nutrient recovery within the digestive tract, reducing feeding costs and concomitant nutrient losses to the environment (Alaa *et al*., 2025; Liu *et al*., 2025).

## Effective microorganisms on blood plasma metabolites in dairy cows

Examination of blood plasma concentration is the key method for understanding how animals metabolize ingested feeds and concentrates, as well as their nutrient absorption efficiency. This section summarizes the impact of EM on important plasma parameters in dairy cows.

Blood plasma concentration can be assessed using haematological techniques, which are valuable for accurately predicting nutrition requirements, monitoring animal health and identifying metabolic disorders and other diseases. Table 1 shows the effect of EM on blood plasma concentrations, focusing on plasma nutrient levels, compositions, hormones and metabolic indicators.

As shown in Table 1, EM positively influences the important blood plasma parameters such as SUN, NEFA, Insulin-like hormones and immunoglobulins. These parameters are key indicators of efficient nutrient utilization and enhanced immunity, both of which may contribute to improved milk production when EM is supplemented.

## Effective microorganisms on manure excretion and methane emission patterns in dairy cows

Effective microorganisms have shown insightful results in manure nutrient excretion and management (Mohamed *et al*., 2018). This section focuses on the specific role of EM in manure composition and its effect on the environment.

As previously discussed, EM shows a promising intervention for digestion and nutrient absorption. Oyebade *et al*., (2023) clearly showed that the supplementation of EM increases the N utilization and protein absorption and reduces the manure N retention, enteric $CH_4$ formation and $NH_3$ emission by modifying the rumen environment. In addition, when $NH_3$ volatilization is decreased, urinary N losses are decreased (Kenney, 2013). Thus, EM application might be useful to the environment in terms of reducing malodor and the eutrophication of water bodies resulting from excessive nutrient inflow.

The EM aids the breakdown of fibrous constituents and enhances nutrient utilization toward faecal solids reduction, owing to augmented microbial activity (Maylem *et al*., 2024; Williams *et al*., 2023; Seo *et al*., 2010). In addition, EM increases a number of enzymatic processes, which results in the lowering of Phosphorus (P) and organic matter excretion in the manure (Ort *et al*., 2018).

**Table 1:** Effects of effective microorganisms on blood plasma concentration in dairy cows.

| Observed effects | Details | Reference |
|---|---|---|
| Improved immune function | Increased neutrophil and monocyte activity, enhancing immune response | Brodzki *et al*., 2024 |
| Increased serum enzyme (GGT) and glucose metabolism | Improved liver function and glucose utilization in lactating cows | Liepa and Viduža, 2018 |
| No significant effect on plasma glucose or hydroxybutyrate | No changes in glucose or ketone metabolism, but better protein content | Kassa and Diffe, 2022 |
| Lower serum urea nitrogen (SUN) and better nitrogen efficiency | Reduced waste nitrogen excretion, improving protein utilization | Oyebade *et al*., 2023 |
| Higher plasma NEFA levels and energy mobilization | Increased fat metabolism, beneficial for early lactation cows | Seo *et al*., 2010 |
| Enhanced plasma hormones (IGF-1) and better feed efficiency | Improved milk yield and feed efficiency with Bacillus subtilis DFMs | Maylem *et al*., 2024 |
| Higher serum IgG and IgA levels (immune boost) | Strengthened immune defence and reduced infection risk | Ort *et al*., 2018 |
| Modulation of plasma metabolites (ammonia-N, VFAs, *etc*.) | Improved ruminal fermentation and microbial protein synthesis | Williams *et al*., 2023 |





**Fig 4:** Impact of effective microorganism supplementation on manure excretion and composition in dairy cows.

Decreased digestion and nutrient absorption efficiency would increase the volume and solid content of the manure, both of which EM could alleviate.

Further, it was reported that using EM reduces volatile compounds in manure such as $NH_3$ and hydrogen sulfide ($H_2S$), which causes malodour and sometimes even mortality of cows due to toxicity would be minimized. Further, it is concluded that effective microorganisms are beneficial in reducing the manure VFA content, which may cause instability and odour production (Bastami *et al*., 2016). Those works claim that EM controls the emission of a broad range of volatile compounds and their effects.

Moreover, contemporary mitigation strategies have shown that probiotics and direct-fed microbials concurrently address enteric $CH_4$ production and manure-based greenhouse gas emissions and meta-analyses have shown that supplementation of probiotics decreases daily $CH_4$ emission by about 11%, while improving nitrogen utilisation efficiency that consequently reduces nitrogen excretion in manure (Xie *et al*., 2025). Methanotroph-based probiotics resulted in significant $CH_4$ reductions in beef steers, with some results exceeding 40 to 50% during *in vitro* trials, thus showing an opportunity for specialised microbial consortia to consume $CH_4$ produced during rumen fermentation directly (Tseten *et al*., 2025). Further, comprehensive works showed that nitrous oxide and $CH_4$ emissions from dairy manure are greatly influenced by the storage period, along with management practices. Probiotics supplementation enhances nutrient retention in animals, thereby reducing the greenhouse gas emission potential from manure storage systems (Cherif *et al*., 2018). Application of EM in manure management, therefore, results in a dual environmental benefit through a reduction of enteric $CH_4$ by 10 to 15% along with a further decrease in nitrogen excretion up to 15-20%, as well as reducing $NH_3$ volatilisation from manure by 25 to 30%, thus mitigating the atmospheric greenhouse gas contributions and nutrient pollution, further contributing toward sustainable dairy production systems (Xie *et al*., 2025; Tseten *et al*., 2025; Zahra *et al*., 2024).

Fig 4 shows, EM minimizes the environmental effects by reducing nutrient excretion and the formation of GHG and odour-forming gases by efficient nutrient utilization.

## CONCLUSION

The application of effective microorganisms in dairy farming presents a promising and sustainable approach to enhancing milk production, optimizing dairy nutrition and managing manure excretion. By improving rumen fermentation, balancing microbial ecology enhancing nutrient digestibility, EM contributes to increased milk yield, improved quality and greater feed conversion efficiency. Besides, EM enhances better N and protein utilization, reduces $CH_4$ emissions and minimizes nutrient excretion in manure, thus minimizing the ecological footprint and supporting sustainable dairy farming.

As the industry trends toward eco-friendly and cost-effective feeding concepts, EM supplementation emerges as a potential alternative to conventional feed additives. However, further research is needed to fully understand its modes of action, long-term impacts and interaction with other dietary components.

## ACKNOWLEDGEMENT

The authors gratefully acknowledge the financial support provided by the National Science Foundation (NSF) of Sri Lanka through Grant No: NSF/RD/06/RG/2024/AG/01.

### Conflict of interest

All authors have no conflict of interest in this study.

## REFERENCES

Abraheem, B.A., Hussein, A.A. and Shachai, N.F. (2024). Effect of effective microorganisms on vegetative growth of wheat. *Agricultural Science Digest.* **44(5):** 859-863. doi: 10.18805/ag.DF-609.

Ahvenjärvi, S. and Huhtanen, P. (2018). Effects of intraruminal urea-nitrogen infusions on feed intake, nitrogen utilization and milk yield in dairy cows. *Journal of Dairy Science.* **101(10):** 9004-9015. doi: 10.3168/jds.2018-14617.

Alaa, M.A., Zhang, H., Duan, H., Zhang, J. and Mao, S. (2025). Metagenomic analysis reveals rumen microbiome enrichment and functional genes adjustment in carbohydrate metabolism induced by different sorting behavior in mid-lactation dairy cows. *Animal Microbiome*. **7(1):** 82. doi: 10.1186/s42523-025-00439-3.

Bastami, M.S.B., Jones, D.L. and Chadwick, D.R. (2016). Reduction of methane emission during slurry storage by the addition of effective microorganisms and excessive carbon source from brewing sugar. *Journal of Environmental Quality.* **45(6):** 2016-2022. doi: 10.2134/jeq2015.11.0568.

Bionaz, M., Vargas-Bello-Pérez, E. and Busato, S. (2020). Advances in fatty acids nutrition in dairy cows: From gut to cells and effects on performance. *Journal of Animal Science and Biotechnology.* **11(1):** 512-520. doi: 10.1186/s40104-020-00512-8.

Brodzki, P., Gorzkoś, H., Marczuk, J., Lisiecka, U., Junkuszew, A., Krakowski, L. and Głodkowska, K. (2024). The influence of probiotic administration on the phagocytic and oxidative burst activity of neutrophils and monocytes in the peripheral blood of dairy cows during different lactation periods. *Journal of Veterinary Research*. **68(3):** 401. doi: 10.2478/jvetres-2024-0043.

Cantalapiedra-Hijar, G., Fouillet, H., Huneau, J.F., Fanchone, A., Doreau, M., Noziere, P. and Ortigues, M.I. (2016). Relationship between efficiency of nitrogen utilisation and isotopic nitrogen fractionation in dairy cows: Contribution of digestion metabolism. *Animals*. **10(2):** 221-229. doi: 10.1017/S1751731115002025.

Cappellozza, B.I., Sousa, D.O., Alveblad, C., Queiroz, O., Joergensen, J.N. and Bengt-Ove, R. (2023). Effects of supplementing a Bacillus-based direct-fed microbial on performance, nutrient digestibility, rumen fermentation characteristics and metabolic responses of lactating dairy cows. *Journal of Dairy Science*. **5(2):** 4-39. doi: 10.3168/jdsc.2023-0439.

Carresi, C., Pauletto, M., Fiore, E., Musolino, V. and Britti, D. (2024). Green veterinary pharmacology and toxicology: A "One health" approach milestone. *Frontiers in Veterinary Science*. **11:** 1476877. doi: 10.3389/fvets.2024.1476877.

Castro-Montoya, J., Kand, D., Selje-Assmann, N. and Dickhoefer, U. (2021). The effects of rumen nitrogen balance on intake, nutrient digestibility, chewing activity and milk yield and composition in dairy cows vary with dietary protein sources. *Journal of Dairy Science.* **104(4):** 4236-4250. doi: 10.3168/jds.2020-19129.

Chanda, T., Khan, M.K.I., Chanda, G.C. and Debnath, G.K. (2024). Correlation between milk urea nitrogen (MUN) levels with metabolizable energy (ME) and crude protein (CP) provided with ratio of roughage concentrate by supplied feed of dairy cattle and its effect on milk yield and milk composition. *Agricultural Reviews*. **45(3):** 546-551. doi: 10.18805/ag.RF-280.

Cherif, C., Hassanat, F., Claveau, S., Girard, J., Gervais, R. and Benchaar, C. (2018). Faba bean (*Vicia faba*) inclusion in dairy cow diets: Effect on nutrient digestion, rumen fermentation, nitrogen utilization, methane production and milk performance. *Journal of Dairy Science.* **101(10):** 8916-8928. doi: 10.3168/jds.2018-14890.

Damunupola, D.A.P.R., Munasinghe, M.A.L.S.S., Rasika, D.M.D. and Weerathilake, W.A.D. . (2022). Sri Lanka's dairy sector: Where to move and what to do, prediction and a trend analysis. *Sri Lankan Journal of Agriculture and Ecosystems.* **4(1):** 81-95. doi: 10.4038/sljae.v4i1.74.

Dhakal, R., Alves, L., Sapkota, R., Khanal, P., Ellegaard, J.L., Winding, A. and Hansen, H.H. (2024). Temporal dynamics of volatile fatty acids profile, methane production and prokaryotic community in an *in vitro* rumen fermentation system fed with maize silage. *Frontiers in Microbiology*. **15:** 1599. doi: 10.3389/fmicb.2024.1271599.

Direkvandi, E., Mohammadabadi, T. and Salem, A.Z.M. (2020). Effect of microbial feed additives on growth performance, microbial protein synthesis and rumen microbial population in growing lambs. *Translational Animal Science*. **4(4):** 203. doi: 10.1093/tas/txaa203.

EMRO-Applied Animal Production. (2024). [Accessed: 2024.12.24]. Retrieved from https://www.em-la.com/en/aplicacoes-e-uso/producao-animal/.

Ezeagu, G.G., Omotosho, A.O. and Suleiman, K.O. (2023). Effective microorganisms, their products and uses: A review. *Nile Journal of Engineering and Applied Science.* **1(1):** 42-53. doi: 10.5455/NJEAS.147954.

Fang, L., Li, Y., Xue, F., Mao, S., Xiong, B., Ma, Z. and Jiang, L. (2021). Effects of bamboo leaf extract on the production performance, rumen fermentation parameters and rumen bacterial communities of heat-stressed dairy cows. *Animal Bioscience.* **34(11):** 20-27. doi: 10.5713/ab.20.0527.

Firrincieli, A., Minuti, A., Cappelletti, M., Ferilli, M., Ajmone-Marsan, P., Bani, P., Petruccioli, M. and Harfouche, A.L. (2024). Structural and functional analysis of the active cow rumen's microbial community provides a catalogue of genes and microbes participating in the deconstruction of cardoon biomass. *Biotechnology for Biofuels and Bioproducts*. **17(1):** 95-110. doi: 10.1186/s13068-024-02495-4.

Fukumori, R., Doi, K., Mochizuki, T., Oikawa, S., Gondaira, S., Iwasaki, T. and Izumi, K. (2022). Sodium butyrate administration modulates the ruminal villus height, inflammation related gene expression and plasma hormones concentration in dry cows fed a high fiber diet. *Animal Science Journal*. **93(1):** 37-91. doi: 10.1111/asj.13791.

Gao, Y., Wei, W., Tian, F., Li, J., Wang, Y., Qi, J. and Xue, S. (2022). Corn straw total mix dietary supplementation of *Bacillus subtilis* enhanced growth performance of lambs by favorably modulating rumen bacterial microbiome. *Fermentation.* **9(1):** 32. doi: 10.3390/fermentation9010032.

Gervais, W.M., Xygalatas, D., McKay, R.T., Van Elk, M., Buchtel, E.E., Aveyard, M. and Bulbulia, J. (2017). Global evidence of extreme intuitive moral prejudice against atheists. *Nature Human Behaviour*. **1(8):** 0151.

Gupta, M., Khan, N., Rastogi, A. and Varun, T.K. (2016). Nutritional drivers of rumen development: A review. *Agricultural Reviews*. **37(2):** 148-153. doi: 10.18805/ar.v37i2.10740.

Halfen, J., Carpinelli, N., Del Pino, F.A.B., Chapman, J.D., Sharman, E.D. Anderson, J.L. and Osorio, J.S. (2021). Effects of yeast culture supplementation on lactation performance and rumen fermentation profile and microbial abundance in mid-lactation holstein dairy cows. *Journal of Dairy Science.* **104(11):** 11580-11592. doi: 10.3168/jds.2020-19996.

Hao, Y., Guo, C., Gong, Y., Sun, X., Wang, W., Wang, Y., Yang, H., Cao, Z. and Li, S. (2021). Rumen fermentation, digestive enzyme activity and bacteria composition between pre-weaning and post-weaning dairy calves. *Animals.* **11(9):** 2527. doi: 10.3390/ani11092527.

Hassan, M.A.S. and Karsl, M.A. (2022). The effects of some feed additives in nutrition of ruminant animals. *Zenodo-European Organisation for Nuclear Research.* **5(2):** 45-97. doi: 10.5281/zenodo.7024597.

Hassanat, F. and Benchaar, C. (2021). Corn silage-based diet supplemented with increasing amounts of linseed oil: Effects on methane production, rumen fermentation, nutrient digestibility, nitrogen utilization and milk production of dairy cows. *Journal of Dairy Science.* **104(5):** 5375-5390.

Huang, G., Guo, L., Chen, M., Wu, X., Tang, W., Zheng, N., Zhao, S., Zhang, Y. and Wang, J. (2022). Biohydrogenation pathway of α-linolenic acid in rumen of dairy cow *in vitro. Animals.* **12(4):** 502. doi: 10.3390/ani12040502.

Iddi-Mrutu, R., Mustapha Umar, K., Abdulhamid, A., Agaba, M. and Abdussamad, A.M. (2023). Microbial engineering to mitigate methane emissions in ruminant livestock-A review. *Proceedings of the National Academy of Sciences.* doi: 10.48550/arXiv.2307.14372.

Jardstedt, M., Hessle, A., Nørgaard, P., Frendberg, L. and Nadeau, E. (2018). Intake and feed utilization in two breeds of pregnant beef cows fed forages with high-fiber concentrations. *Journal of Animal Science.* **96(8):** 3398-3411. doi: 10.1093/jas/sky199.

Jiang, A., Liu, Z., Yang, Z., Zhang, S., Wu, J., Zhou, C. and Tan, Z. (2025). Direct-fed microbials optimize ruminal fermentation, microbial ecosystem and milk quality to enhance the lactation performance of Sanhe dairy cows. *Animal Microbiome*. **7(1):** 437-442. doi: 10.1186/s42523-025-00437-5.

Kand, D., Raharjo, I.B., Castro-Montoya, J. and Dickhoefer, U. (2018). The effects of rumen nitrogen balance on *in vitro* rumen fermentation and microbial protein synthesis vary with dietary carbohydrate and nitrogen sources. *Animal Feed Science and Technology.* **241:** 184-197.

Karangiya, V.K., Savsani, H.H. and Ribadiya, N.K. (2016). Use of densified complete feed blocks as ruminant feed for sustainable livestock production: A review. *Agricultural Reviews.* **37(2):** 141-147. doi: 10.18805/ar.v37i2.10739.

Kassa, T. and Diffe, Z. (2022). The role of direct-fed microbes to ruminants: A review. *Global Journal of Animal Scientific Research.* **10:** 1-13.

Kenney, N. (2013). Impact of direct-fed microbials on nutrient utilization in beef cattle. Thesis and Dissertation, University of Kentucky, USA.

Kumar, P., Jain, K.K., MunilKumar, S. and Sudhagar, S.A. (2017). Alternate feeding strategies for optimum nutrient utilization and reducing feed cost for semi-intensive practices in aquaculture system-A review. *Agricultural Reviews.* **38(2):** 145-151. doi: 10.18805/ag.v38i02.7946.

Kupczyñski, R., Szumny, A., Wujcikowska, K. and Pachura, N. (2020). Metabolism, ketosis treatment and milk production after using glycerol in dairy cows: A review. *Animals.* **10(8):** 1379. doi: 10.3390/ani10081379.

Lee, C., Feyereisen, G.W., Hristov, A.N., Dell, C.J., Kaye, J. and Beegle, D. (2019). Effects of dietary protein concentration on ammonia volatilisation, nitrate leaching and plant nitrogen uptake from dairy manure applied to lysimeters. *Journal of Environmental Quality.* **43(1):** 398-408.

Liepa, L. and Viduža, M. (2018). The effect of peroral administration of *Lactobacillus* fermentum culture on dairy cows' health indices. *Macedonian Veterinary Review.* **41(2):** 2018-0017. doi: 10.2478/macvetrev-2018-0017.

Lima, J., Ingabire, W., Roehe, R. and Dewhurst, R.J. (2023). Estimating microbial protein synthesis in the rumen-can 'Omics' methods provide new insights into a long-standing question. *Veterinary Sciences*. **10(12):** 679-679. doi: 10.3390/vetsci10120679.

Liu, Y., Min, Q., Tang, J., Yang, L., Meng, X., Peng, T. and Jiang, M. (2023). Transcriptome profiling in rumen, reticulum, omasum and abomasum tissues during the developmental transition of pre-ruminant to the ruminant in yaks. *Frontiers in Veterinary Science.* **10(12):** 706. doi: 10.3389/fvets.2023.1204706.

Liu, Z., Jiang, A., Ma, D., Liu, D., Han, X., Zhao, M., Zhou, C. and Tan, Z. (2025). The impact of rumen microbial composition on apparent digestibility, rumen fermentation and metabolism in sanhe cows and holstein cows of different parities under identical dietary conditions. *Frontiers in Veterinary Science*. **11:** 1463209-1463215. doi: 10.3389/fvets.2024.1463209.

Lopreiato, V., Mezzetti, M., Cattaneo, L., Ferronato, G., Minuti, A. and Trevisi, E. (2020). Role of nutraceuticals during the transition period of dairy cows: A review. *Journal of Animal Science and Biotechnology.* **11(1):** 96. doi: 10.1186/s40104-020-00501-x.

Lv, F., Wang, X., Pang, X. and Liu, G. (2020). Effects of supplementary feeding on the rumen morphology and bacterial diversity in lambs. *Zoological Science.* **8:** e9353. doi: 10.7717/peerj.9353.

Mamuad, L.L., Kim, S.H., Biswas, A.A., Yu, Z., Cho, K.K., Kim, S.B., Lee, K. and Lee, S.S. (2019). Rumen fermentation and microbial community composition influenced by live enterococcus faecium supplementation. *Applied Microbiology and Biotechnology Express.* **9(1):** 123. doi: 10.1186/s13568-019-0848-8.

Marinho, M.N., Perdomo, M.C., Simões, B.S., Husnain, A., Arshad, U., Figueiredo, C.C., Peixoto, P.M., Monteiro, H.F. Anderson, B. and Santos, J.E.P. (2025). Impact of dietary supplementation of rumen native microbes on ruminal microbiome, ruminal fermentation and total-tract digestibility in holstein cows. *Journal of Dairy Science*. **108(11):** doi: 10.3168/jds.2025-26900.

Maylem, E.R., Vargiu, M., Melton, C., Stiger, C., Thompson, J.S., O'Neill, J. and Spicer, L.J. (2024). Effect of a Bacillus subtilis-based direct-fed microbial, on milk yield, milk components, feed intake and plasma hormones and metabolites in lactating Holstein cows. *Animal Feed Science and Technology.* **318:** 116-136. doi: 10.1016/j.anifeedsci.2024.116136.

Mayulu, H. and Christiyanto, M. (2025). Literature review rumen microbiota mysteries: Nutrition, function and its impact on livestock productivity. *American Journal of Animal and Veterinary Sciences*. **20(2):** doi: 10.3844/ajavsp.2025.180.191.

Mitchell, K.E., Kienzle, S.L., Lee, C., Socha, M.T., Kleinschmit, D.H. and Firkins, J.L. (2023). Supplementing branched-chain volatile fatty acids in dual-flow cultures varying in dietary forage and corn oil concentrations. II: Biohydrogenation and incorporation into bacterial lipids. *Journal of Dairy Science*. **106(11):** 7548-7565. doi: 10.3168/jds.2022-23192.

Mohamed, M., Amarasinghe, A.A.S.S.M. and Nayananjalie, W.A.D. (2018). Use of effective microorganisms in dry zone dairy farms in Sri Lanka. *International Journal of Livestock Research.* **8(4):** 50-57.

Mohanty, P.P., Nagalakshmi, D., Tarjan, K. and Sriharsha, K.V. (2023). Effect of dietary supplementation of chromium and yeast on *in vitro* dry matter degradability and rumen fermentation pattern. *Indian Journal of Animal Research.* **57(2):** 201-206. doi: 10.18805/IJAR.B-4758.

Mu, Y., Lin, X., Wang, Z., Hou, Q., Wang, Y. and Hu, Z. (2019). High production dairy cattle exhibit different rumen and fecal bacterial community and rumen metabolite profile than low production cattle. *Microbiology Open*. **8(4):** 00673. doi: 10.1002/mbo3.673

Nair, S.S., Williams, R.O., Neachtain, A.S., Tognelli, R., Chandra, S., Alvarez-Hess, P.S., Cheng, L., Giri, K. and Jacobs, J.L. (2025). The effect of ruminal fluid adaptation to a direct fed microbial: *In vitro* methane production and fermentation characteristics. *Fermentation.* **11(8):** 473-478. doi: 10.3390/fermentation11080473.

Nalla, K., Manda, N.K., Dhillon, H.S., Kanade, S.R., Rokana, N., Hess, M. and Puniya, A.K. (2022). Impact of probiotics on dairy production efficiency. *Frontiers in Microbiology.* **13:** 805- 963. doi: 10.3389/fmicb.2022.805963.

Natt, S.K. and Katyal, P. (2022). Current trends in non-dairy probiotics and their acceptance among consumers: A review. *Agricultural Reviews*. **43(4):** 450-456. doi: 10.18805/ag.R-2172.

Ngesi, A., Laswai, G., Msalya, G., Lyatuu, E., Komwihangilo, D. and Mrode, R. (2025). An assessment on the feeding practices and other factors influencing dairy production under smallholder systems of Kilimanjaro region in Tanzania. *International Journal of Animal Science and Technology.* **9(2):** 87-99. doi: 10.11648/j.ijast.20250902.17.

OECD (2025). OECD-FAO Agricultural outlook 2025-2034. Organisation for Economic Co-operation and Development Publishing. Paris. doi: 10.1787/601276cd-en.

Ort, S.B., Aragona, K.M., Chapman, C.E., Shangraw, E., Brito, A.F., Schauff, D.J. and Erickson, P.S. (2018). The impact of direct fed microbials and enzymes on the health and performance of dairy cows with emphasis on colostrum quality and serum immunoglobulin concentrations in calves. *Journal of Animal Physiology and Animal Nutrition.* **102(2):** 641-652. doi: 10.1111/jpn.12806.

Owens, F.N. and Basalan, M. (2016). Ruminal Fermentation. In: Rumenology. Springer International Publishing. pp (63-102).

Oyebade, A.O., Lee, S., Sultana, H., Arriola, K., Duvalsaint, E., De Guzman, C.N. and Vyas, D. (2023). Effects of direct-fed microbial supplementation on performance and immune parameters of lactating dairy cows. *Journal of Dairy Science*. **106(12):** 8611-8626. doi: 10.3168/jds.2022-22898.

Parchami, M., Rustas, B.O., Taherzadeh, M.J. and Mahboubi, A. (2024). Effect of agro-industrial by-products derived from volatile fatty acids on ruminant feed *in vitro* digestibility. *Animals*. **14(16):** 23-30. doi: 10.3390/ani14162330.

Pokhrel, B. and Jiang, H. (2024). Postnatal growth and development of the rumen: Integrating physiological and molecular insights. *Biology.* **13(4):** 269. doi: 10.3390/biology13040269.

Polyorach, S., Nampukdee, R., Wanapat, M., Kang, S., Cherdthong, A., Poungchompu, O., Gunun, P., Gunun, N., Foiklang, S., Thinowong, A., Jindatajak, Y., Lapmee, A. and Norrapoke, T. (2023). Microbial fermented liquid supplementation improves nutrient digestibility, feed intake and milk production in lactating dairy cows fed total mixed ration. *Animals.* **13(5):** 933. doi: 10.3390/ani13050933.

Putri, E.M., Zain, M., Warly, L. and Hermon, H. (2021). Effects of rumen-degradable-to-undegradable protein ratio in ruminant diet on *in vitro* digestibility, rumen fermentation and microbial protein synthesis. *Journal of Veterinary World.* **14(3):** 640-648. doi: 10.14202/vetworld.2021.640-648.

Qin, D., Gao, R. and Ji, D. (2025). Synergy between genetics and nutrition: A systematic review of factors affecting performance in dairy cows. *Frontiers in Animal Science*. **6(1):** doi: 10.3389/fanim.2025.1653052.

Riaz, F., Rauf, N., Bilal, M., Hamza, M.A., Irfan, M., Akhter, S., Ahmed, H., Talha, M., Hussnain, A., Haider, I. and Javaid, T. (2024). Use of Probiotics and Prebiotics in Livestock Productivity. In: Complementary and Alternative Medicine: Prebiotics and Probiotics. [Farooqi, S.H., Aqib, A.I., Zafar, M.A., Akhtar, T. and Ghafoor, N. (Eds)], Unique Scientific Publishers, Faisalabad, Pakistan. pp 31-38. doi: 10.47278/book.CAM/2024.328.

Sawant, S.S., Park, H.Y., Sim, E.Y., Kim, H.S. and Choi, H.S. (2025). Microbial fermentation in food: Impact on functional properties and nutritional enhancement, a review of recent developments. *Fermentation*. **11(1):** 15. doi: 10.3390/fermentation11010015.

Seo, J.K., Kim, S.W., Kim, M.H., Upadhaya, S.D., Kam, D.K. and Ha, J.K. (2010). Direct-fed microbials for ruminant animals. *Asian-Australasian Journal of Animal Sciences*. **23(12):** 1657-1667. doi: 10.5713/ajas.2010.r.08.

Sheikh, G.G., Ganai, A.M., Reshi, P.A., Bilal, S., Mir, S. and Masood, D. (2018). Improved paddy straw as ruminant feed: A review. *Agricultural Reviews*. **39(2):** 137-143. doi: 10.18805/ag.R-1667.

Shi, R., Dong, S., Mao, J., Wang, J., Cao, Z., Wang, Y., Li, S. and Zhao, G. (2023). Dietary neutral detergent fiber levels impacting dairy cows' feeding behavior, rumen fermentation and production performance during the period of peak-lactation. *Animals*. **13(18):** 2876. doi: 10.3390/ani13182876.

Shukla, N., Tripathi, A., Tiwari, A. and Sharma, K. (2024). Probiotics in dairy farming: A holistic approach to animal nutrition. *International Journal of Advanced Biochemistry Research*. **8(5):** 360-373. doi: 10.33545/26174693.2024.v8.i5sf.1206.

Steele, M.A., Penner, G.B. and Chaucheyras, D.F. (2016). Development and physiology of the rumen and the lower gut: Targets for improving gut health. *Journal of Dairy Science*. **99(6):** 4955-4966. doi: 10.3168/jds.2015-10351.

Sugandhi, G.P. (2018). Probiotics conventional benefits and cautions in intake-A review. *Agricultural Reviews*. **39(3):** 251-254. doi: 10.18805/ag.R-1749.

Tadesse, E.T., Minneeneh, G.A., Mummed, Y.Y., Dejene, M. and Kebreab, E. (2024). Effect of effective microorganisms treated wheat bran supplementation on intake, digestibility and milk yield and composition of lactating cows fed natural pasture hay as a basal diet. *Livestock Research for Rural Development*. **36(2):** 11.

Tong, J., Zhang, H., Yang, D., Zhang, Y., Xiong, B. and Jiang, L. (2018). Illumina sequencing analysis of the ruminal microbiota in high-yield and low-yield lactating dairy cows. *Plos One*. **13(11):** 0198225. doi: 10.1371/journal.pone.0198225

Torres, R.N.S., Bertoco, J.P.A., de Arruda, M.C.G., Rodrigues, J.L., Coelho, L.M., Paschoaloto, J.R., de Almeida Júnior, G.A., Ezequiel, J.M.B. and Almeida, M.T.C. (2021). Meta-analysis of the effect of glycerin inclusion in dairy cattle diet on milk fatty acid profile. *Translational Animal Science*. **5(1):** 12. doi: 10.1093/tas/txab012.

Tseten, T., Sanjorjo, R.A., Son, J.W., Baik, K.S., Berdos, J.I., Kim, S.H., Yoon, S.H., Kang, M.K., Kwon, M., Lee, S.S. and Kim, S.W. (2025). Reduction of enteric methane emission using methanotroph-based probiotics in hanwoo steers. *Animal Microbiome*. **7(19):** 385. doi: 10.1186/s42523-025-00385-0.

Tu-Kitaw, G., Daba, T., Kehaliw, A., Assefa, G., Dejene, M. and Fekadu, D. (2013). Evaluation of effective microbes (EM) as non-conventional feed supplement to lactating crossbred cows and barn malodor control under Ethiopian condition. *Journal of Renewable Agriculture*. **1(8):** 141-146. doi: 10.12966/jra.11.02.2013.

Vasta, V., Daghio, M., Cappucci, A., Buccioni, A., Andrea, S., Viti, C. and Mele, M. (2019). Invited review: Plant polyphenols and rumen microbiota responsible for fatty acid biohydrogenation, fiber digestion and methane emission: Experimental evidence and methodological approaches. *Journal of Dairy Science*. **102(5):** 3781-3804. doi: 10.3168/jds.2018-14985.

Vidanarachchi, J.K., Chathurika, H.M.M., Dias, H.M., Gedara, P.K., Silva, G.L.L.P., Perera, E.R. K. and Perera, A.N.F. (2019). Dairy industry in Sri Lanka: Current status and way forward for a sustainable industry. Sri Lanka Association of Animal Production, University of Peradeniya: Peradeniya, Sri Lanka. pp 1-38.

Vlaicu, P.A. and Untea, A.E. (2024). Feeding strategies and quality assessments of animal-derived products. *Agriculture*. **14(11):** 1949. doi: 10.3390/agriculture14111949.

Wang, K., Jiang, M., Chen, Y., Huang, Y., Cheng, Z., Datsomor, O., Jama, S.M., Zhu, L., Li, Y., Zhao, G. and Lin, M. (2024). Changes in the rumen development, rumen fermentation and rumen microbiota community in weaned calves during steviol glycosides treatment. *Frontiers in Microbiology*. **15:** 1395665. doi: 10.3389/fmicb.2024.1395665.

Wei, W., Wang, L., Bao, P., Shao, Y., Yue, H., Yang, D. and Wang, H. (2018). Metal-Free C (sp2)-H/N-H cross-dehydrogenative coupling of quinoxalinones with aliphatic amines under visible-light photoredox catalysis. *Organic Letters*. **20(22):** 7125-7130. doi: 1021/acs.orglett.8b03079.

Wei, W., Zhen, Y., Wang, Y., Shahzad, K. and Wang, M. (2022). Advances of rumen functional bacteria and the application of micro-encapsulation fermentation technology in ruminants: A review. *Fermentation*. **8(10):** 564. doi: 10.3390/fermentation8100564.

Weimer, P.J. (2022). Degradation of cellulose and hemicellulose by ruminal microorganisms. *Microorganisms*. **10(12):** 2345. doi: 10.3390/microorganisms10122345.

Williams, S.R.O., Jacobs, J.L., Chandra, S., Soust, M., Russo, V.M., Douglas, M.L. and Hess, P.S.A. (2023). The effect of direct-fed lactobacillus species on milk production and methane emissions of dairy cows. *Animals*. **13(6):** 1018. doi: 10.3390/ani13061018.

Wilson, R.L., Bionaz, M., MacAdam, J.W., Beauchemin, K.A., Naumann, H.D. and Ates, S. (2020). Milk production, nitrogen utilization and methane emissions of dairy cows grazing grass, forb and legume-based pastures. *Journal of Animal Science*. **98(7):** 220. doi: 10.1093/jas/skaa220.

Wu, R., Chang, S., Zhang, H., Yang, X., Gu, R., Wang, S., Liu, X., Liu, X., Bat Ochir, M. and Wu, J. (2024). Compound probiotics microcapsules improve milk yield and milk quality of dairy cows by regulating intestinal flora. *Food Bioengineering*. **3(1):** 110-125. doi: 10.1002/fbe2.12084.

Xie, X., Cao, Y., Li, Q., Li, Q., Yang, X., Wang, R., Zhang, X., Tan, Z., Lin, B. and Wang, M. (2025). Mitigating enteric methane emissions: An overview of methanogenesis, inhibitors and future prospects. *Animal Nutrition*. **21:** 84-96. doi: 10.1016/j.aninu.2025.02.001.

Yang, J., Li, Y., Sun, M., Guo, S., Lin, P., Wang, A. and Jin, Y. (2025). Understanding the differences in rumen bacteria and their impact on dairy cows' production performance: A review. *Animal Nutrition*. **22(1):** 259-279. doi: 10.1016/j.aninu.2025.04.006.

Yang, J., Li, Y., Sun, M., Zhang, Y., Guo, S., Zhou, D., Lin, P., Wang, A. and Jin, Y. (2025). Associations of rumen and rectum bacteria with the sustained productive performance of dairy cows. *Frontiers in Microbiology*. **16:** 1565034. doi: 10.3389/fmicb.2025.1565034.

Yang, J., Zhao, S. and Lin, B. (2024). Effect of commercial fibrolytic enzymes application to normal- and slightly lower energy diets on lactational performance, digestibility and plasma nutrients in high-producing dairy cows. *Frontiers in Veterinary Science.* **11(1):** 1302034. doi: 10.3389/fvets.2024.1302034.

Yitbarek, M.B. (2023). Effects of microorganism supplementation on egg quality and production. *Journal of World's Poultry Research.* **13(4):** 406-418. doi: 10.36380/jwpr.2023.43.

Zahra, R., Madjid, T. and Nadia, H. (2022). Indigestible foreign bodies in the rumen-reticulum of cattle slaughtered at batna slaughterhouse, Algeria: A postmortem study. *Agricultural Reviews.* **43(2):** 260-263. doi: 10.18805/ag.RF-222.

Zahra, W.A., Shiddieqy, M.I., Anisa, R., Yani, A. and Purwanto, B.P. (2024). The dynamics of nitrous oxide and methane emissions from various types of dairy manure at smallholder dairy farms as affected by storage periods. *Waste Management.* **183:** 10-20. doi: 10.1016/j.wasman.2024.04.039.

Zeeshan, M., Masood, S., Ashraf, S., Bokhar, S.G., Zainab, H., Ijaz, S. and Usman, M.M. (2023). Efficacy of mannan-oligosaccharide and live yeast feed additives on performance, rumen morphology, serum biochemical parameters and muscle morphometric characteristics in buffalo calves. *Journal of Population Therapeutics and Clinical Pharmacology*. **31(6):** 1401-1410. doi: 10.53555/jptcp.v31i6.6684.

Zhu, W., Guo, J., Li, X., Li, Y., Song, L., Li, Y., Feng, B., Bao, X., Li, J., Gao, Y. and Xu, H. (2025). Effects of gallic acid on *in vitro* ruminal fermentation, methane emission, microbial composition and metabolic functions. *Animals.* **15(13):** 1959. doi: 10.3390/ani15131959.

Zurak, D., Kljak, K. and Aladrović, J. (2023). Metabolism and utilisation of non-protein nitrogen compounds in ruminants: A review. *Journal of Central European Agriculture*. **24(1):** 1-14. doi: 10.5513/JCEA01/24.1.3645.