RESEARCH ARTICLE Agricultural Science Digest, Volume 45 Special Issue: 118-123 (November 2025)

# Analysis of Metabolomic Compounds and Antioxidant Potential of Solanaceae Family Plants using Gcms after Elicitor Induction Methyl Jasmonate and Salicylic Acid

Henny Lieke Rampe[1], Hanny Hesky Pontororing[1], Hanry Jefry Lengkong[1], Meytij Jeanne Rampe[2], Vistarani Arini Tiwow[3], Wilson Marthin Moniaga[4]

10.18805/ag.DF-687

## ABSTRACT

**Background:** Solanaceae plants have the potential to be a source of food, medicine, industrial materials and antioxidants. Antioxidants are compounds that can delay or prevent oxidation caused by free radicals significantly. Metabolomic analysis is a method used to study metabolite profiles. It is necessary to use elicitors to optimize the synthesis of metabolomic compounds and antioxidant potential: methyl jasmonate and salicylic acid as elicitors.

**Methods:** This research aims to analyze the Solanaceae plant's metabolomic compounds and antioxidant potential using GCMS after the elicitors methyl jasmonate and salicylic acid induction. Experimental method research with stages: 1) Observation, collection and identification of Solanaceae plants, 2) Preparation of experimental land, 3) Planting of plant collections, 4) Application of elicitor methyl jasmonate 60 ppm and salicylic acid 0.45 g/L, 5) Metabolomic and antioxidant analysis on fresh extracts of samples. Testing antioxidant activity using a spectrophotometer with DPPH solution. Metabolomic profile analysis using GCMS.

**Result:** The results of the non-targeted metabolomic analysis with GCMS and antioxidant activity with $IC_{50}$ in the *Solanum genus*, namely *Solanum betaceum, Solanum melongena*, *Solanum torvum*, *Solanum Lycopersicum* and *Solanum tuberosum,* respectively identified 18, 16, 14, 13 and 20 compounds. The $IC_{50}$ values were 51.32 mg/L (strong category), 56.64 mg/L (strong category), 65.86 mg/L (strong category), 113.60 mg/L (medium category), 102.71 mg/L (medium category), respectively. In the *Capsicum* genus, namely *Capsicum frutescens* and *Capsicum annuum*, 21 and 37 compounds were identified, respectively, with $IC_{50}$ values of 124.76 mg/L (medium category) and 63.23 mg/L (strong category). In *Physalis angulata*, 19 compounds were identified, $IC_{50}$ was 44.04 mg/L (very strong category), six compounds were identified in *Nicotiana tabacum* and $IC_{50}$ was 67.55 mg/L (strong category). Solanaceae plants have moderate to very strong antioxidant potential.

**Key words:** Antioxidants, Metabolomics, Solanaceae family.

## INTRODUCTION

The Solanaceae family is a group of shrubs or trees with single or compound leaves and trumpet-shaped flowers. Some important genera from the Solanaceae family are *Solanum*, *Capsicum*, *Nicotiana* and *Physalis*. Solanaceae plants have the potential to be a source of food, medicine, industrial materials and antioxidants (Artanti and Lisnasari, 2018; Krisnawati and Febrianti, 2019; Azzikri *et al.,* 2020). Antioxidants are compounds that can significantly delay or prevent oxidation caused by free radicals. Antioxidants result from metabolite activity produced by various compounds, which can be studied through metabolomic analysis (Kumari *et al.,* 2023).

Several studies on the antioxidant potential of Solanaceae plants have been carried out on *Capsicum annuum* (Ananta and Anjasmara, 2022), *Capsicum frutescens* (Misfadhila *et al.,* 2022), *Solanum tuberosum* (Hellmann *et al.,* 2021; Dewi *et al.,* 2024). *Solanum lycopersicum* (Pujiastuti *et al.,* 2019; Hasfikasari, 2024), *Physalis angulata* (Putra *et al.,* 2023; Sukmawati *et al.,* 2024), *Datura metel* (Mbida *et al.,* 2022). *Solanum torvum* (Helilusiatiningsih and Soenyoto, 2020), *Nicotiana tabacum* (Al-Lahham *et al.,* 2020; Zou *et al.,* 2021). Antioxidant activity was determined through decolorization of 2,2-diphenyl-1-picrylhydrazyl (DPPH). The use of DPPH in the antioxidant potential test has been carried out on extracts (Leaves and Fruit) of *Cucurbita maxima* (Wafa, 2024),

[1]Department of Biology, Faculty of Mathematics and Natural Sciences, Sam Ratulangi University, Indonesia.
[2]Department of Chemistry, Faculty of Mathematics and Natural Sciences, Manado State University, Indonesia.
[3]Department of Physics, Faculty of Mathematics and Natural Sciences, Manado State University, Indonesia.
[4]Department of Pests and Diseases, Faculty of Agriculture, Sam Ratulangi University, Indonesia.

**Corresponding Author:** Henny Lieke Rampe, Department of Biology, Faculty of Mathematics and Natural Sciences, Sam Ratulangi University, Indonesia. Email: hennyrampe@unsrat.ac.id
ORCID: https://orcid.org/my-orcid?orcid=0000-0001-5533-5009

**How to cite this article:** Rampe, H.L., Pontororing, H.H., Lengkong, H.J., Rampe, M.J., Tiwow, V.A. and Moniaga, W.M. (2025). Analysis of Metabolomic Compounds and Antioxidant Potential of Solanaceae Family Plants using Gcms after Elicitor Induction Methyl Jasmonate and Salicylic Acid. *Agricultural Science Digest*. **45 (Special Issue):** 118-123. doi: 10.18805/ag.DF-687.

**Submitted:** 15-10-2024 **Accepted:** 03-06-2025 **Online:** 27-06-2025

*Syzygium cumini* (Devi *et al.,* 2023), Turnip Roots (*Brassica rapa*) Beniwal *et al.,* (2022), Nardostachys Jatamansi (Pant *et al.,* 2024).

Metabolomic analysis is a method used to study metabolite profiles. It is necessary to use elicitors to optimize the synthesis of metabolomic compounds and antioxidant potential methyl jasmonate and salicylic acid as elicitors (Diaz *et al.,* 2016). Jasmonic acid can induce hormone synthesis and gene expression (Huang *et al.,* 2017; Ruan *et al.,* 2019). Jasmonic acid is a natural compound synthesized by plants as a resistance response to pathogen attack and external damage (Siddiqi and Husen, 2019; Gomi, 2020; Sanchez *et al.,* 2020). Salicylic acid is an endogenous hormone that plays a role in plant physiological processes, protein synthesis and abiotic stress (Khatun *et al.,* 2016).

Research on induction of the elicitor methyl jasmonate (MJ) (Rampe *et al.,* 2022), a combination of the elicitors salicylic acid and methyl jasmonic (Rampe *et al.,* 2023) has been carried out on sweet potatoes. The results showed increased quantitative morphoanatomical characters and flavonoid and tannin content. Tam-Ho *et al.* (2020) found that applying exogenous jasmonic acid and methyl jasmonate induced reactive oxygen species and resistance mechanisms with the accumulation of secondary metabolites. Testing of MJ *in vitro* cultures has induced antioxidant enzyme activity, expression of resistance-related genes and increased production of secondary metabolites. The potential for methyl jasmonate to induce secondary metabolite biosynthesis has been carried out by Sanchez *et al.* (2020) on the suspension culture of *Piper cumanense*, Nguyen *et al.* (2019) on the root culture of *Centella asiatica*. Solanaceae plants have antioxidant potential (Ralte *et al.,* 2021) because they contain metabolomic compounds, namely alkaloids, flavonoids, saponins, tannins and terpenoids. This research focuses on the induction of elicitors methyl jasmonate and salicylic acid on antioxidant potential and metabolomic compound content.

## MATERIALS AND METHODS

Research activities were carried out in experimental gardens located in Tomohon City, North Sulawesi Province and the advanced biology laboratory of the Biology Department at Sam Ratulangi University Indonesia. GCMS analysis at the North Sulawesi Regional Police Forensic Laboratory, in 2024.

### Research procedures

a. Observation, collection and identification of *Solannaceae* plants in Tomohon City.
b. Preparation of experimental land. The experimental plots were 10 × 10 m in size.
c. Planting a collection of solanaceae plants. Each plot is planted with 16 plants.
d. The elicitor methyl jasmonate 60 ppm and salicylic acid 0.45 g/L, 25 days after planting (DAP) and 50 DAP were applied by spraying on the whole plant. Control without methyl jasmonate and salicylic acid treatment.
e. Metabolomic and antioxidant analysis on fresh extracts of samples. The materials used were fruits of *Solanum betaceum*, *Solanum melongena*, *Solanum torvum*, *Solanum lycopersicum*, *Capsicum frutescens*, *Capsicum annuum*. Furthermore, leaves on *Physalis angulata* and *Nicotiana tabacum* and tubers on *Solanum tuberosum*. The material (fruit/leaf/tuber) of each plant sample was pulverized, squeezed with a filter cloth; then, the extract was filtered to obtain the filtrate.

### Antioxidant activity testing

Each test sample was made into a stock solution of 500 ppm by weighing the sample 0.005 g and then dissolving it in 10 mL methanol to obtain a sample of 500 ppm. Next, the mother liquor was diluted to 20, 40, 60, 80 and 100 ppm. 1.5 mg of fresh sample extract was dissolved in 1 mL of DMSO to obtain a solution with a concentration of 50/mL. The sample was incubated for 20 minutes. Extract 2 mL of the sample and add 2 mL of DPPH, the sample is incubated for 30 minutes in dark conditions. Antioxidant activity was determined through decolorization of DPPH at a wavelength of 514 nm using a UV-Vis spectrophotometer.

Through a linear equation that states the relationship between the concentration of the test extract (x) and free radical scavenging activity (y). The antioxidant activity of the sample extract was determined based on the percentage of inhibitory power relative to the control using the equation of Handayani *et al.* (2018), namely.

$$\% \text{ Inhibition} = \frac{\text{A DPPH - A Sample}}{\text{A DPPH}} \times 100$$

The level of antioxidant power using the DPPH method: $IC_{50}$ value (ppm)<50 (very strong), 50-100 (strong), 100-150 (medium), 150-200 (weak) and >200 (very weak) (Molyneux, 2004).

Metabolomic compound analysis using GCMS version of Agilent 8890 GC System. The sample used was a fresh extract solution of Solanaceae plants. The sample extract was dissolved in methanol (1:1) and 2 μL was injected into the GCMS.

## RESULTS AND DISCUSSION

The samples used were fresh extract solutions from Solanaceae plants. The entire area was then identified qualitatively through the method of comparing the spectrum data obtained with the spectrum data in the data bank. Next, look for the compound chain in PubChem and then copy the chain obtained to way2drug.com to find out the potential of the metabolite compound. The study is focused on identifying large percent areas that show the greatest composition of compounds contained in the sample.

The GCMS results of terong belanda/tamarillo (*Solanum betaceum*) (Fig 1), obtained 18 metabolite compounds, three compounds that dominate with the highest area (%) namely thiocyanic acid, retention time (rt)

8.952, area 8.64 %; 4H-pyran-4-one, rt 7,720 area 5.25%; 2H-imidazole-2-one, rt 4.558 area 6.39%. Based on DPPH test, $IC_{50}$ value = 51.32 mg/L strong antioxidant category, $IC_{50}$ value of control 82.16 mg/L. The GCMS results of terong ungu/purple eggplant (*Solanum melongena*) fruit (Fig 1) showed that 16 metabolite compounds were obtained, three compounds that dominated with the highest area (%) namely 4H-pyran-4-one, rt 7.745, area 10.62%; 2-Cyclopenten-1-one rt. 4,721 area 13.36%, propane rt. 5,744 area 11.28%. Based on the DPPH test, $IC_{50}$ value = 56.64 mg/L in the category of strong antioxidant properties, IC50 value of control 84.34 mg/L.

The GCMS results of takokak (*Solanum torvum*) leaves (Fig 1) showed that 14 metabolite compounds were obtained, three compounds that dominated namely phytol rt.18.461 area 37.13%; alpha-L-Galactopyranoside, rt. 13,783 area 14.03%; 2-Propen-1-ol, rt 5.563 area 6.36%, Based on DPPH test, $IC_{50}$ value = 65.86 mg/L strong antioxidant category, $IC_{50}$ value of control 94.64 mg/L. The GC-MS results of tomat / tomato (*Solanum lycopersicum*) (Fig 1) showed that 13 metabolite compounds were obtained, three compounds that dominated with the highest area (%) namely methyl glyoxal, rt 3,517 area 37.44%; (furan-2-yl) methanol, rt. 4,860 area 11.41%; 1,3-Cyclohexane-11, rt. 5,706 area 9.69%. Based on the DPPH test, $IC_{50}$ value = 113.60 mg/L in the medium antioxidant category, $IC_{50}$ value of control 136.12 mg/L. The GCMS results tubes of kentang / potato (*Solanum tuberosum*) (Fig 1) showed that 20 metabolite compounds were obtained, three compounds that dominated with the highest area (%) namely Hexadecanoic acid, rt.16,706 area 22.66%; 2-pyrrolidinone rt. 6,861 area 15.44%; methyl stearate, rt.18,552 area 1.74%. Based on the DPPH test, $IC_{50}$ value = 102.71 mg/L in the medium antioxidant category, $IC_{50}$ value of control 144.07 mg/L.

The GCMS results of cabai rawit/cayenne pepper (*Capsicum frutescens*) fruit (Fig 2), obtained 21 metabolite compounds, the three compounds that dominated were Capsaicin, rt. 22,379 area 63.15%; Dihydrocapsaicin, rt.22,521 area 19.92%; acetamide, rt.18,895 area 3.25%.





**Fig 1:** GCMS results of *S. betaceum* (A), *S. melongena*) (B), *S. torvum* (C), *S. lycopersicum* (D) and *S. tuberosum* (E).





**Fig 2:** GCMS results of *C. frutescens* (A) and *C. annuum* (B).





**Fig 3:** GCMS results of *P. angulata* (A) and *N. tabacum* (B).

Based on the DPPH test, IC50 value = 124.76 mg/L in the medium antioxidant category, $IC_{50}$ value of control 165.94 mg/L. The GCMS results of cabai merah/red chili (*Capsicum annuum*) fruit (Fig 2), obtained 37 metabolite compounds, three compounds that dominated namely O-Isopropylhydroxylamine rt 3,198 area 27.84%; 6-Oxa-bicyclo (3.1.0) hexan-3-one rt 4.639 area 13.08%; Acetamide, rt 18.903, area 6.27%. Based on the DPPH test, IC50 value = 63.23 mg/L in the category of strong antioxidant properties, $IC_{50}$ value of control 83.14 mg/L.

The GCMS results of ciplukan/morel berry (*Physalis angulata*) (Fig 3) obtained 19 metabolite compounds, three compounds that dominated namely 4H-Pyran-4-one rt 7,681 area 7.40%; Cyclopropane rt 4,632 area 5.14%; Topotecan rt 5,577 area 4.40%. Based on the DPPH test, the $IC_{50}$ value 44.04 mg/L in the category of very strong antioxidant, $IC_{50}$ value of control 79.16 mg/L. Furthermore, six metabolite compounds were obtained from tembakau /tobacco leaves (*Nicotiana tabacum*), three compounds dominated Pyridine, rt. 10,425 area 88.97%; Caryophyllene oxide, rt. 19.768 area 3.63% and Scopoletine, rt.17.383 area 1.33%. Based on the DPPH test, $IC_{50}$ value = 67.55 mg/L category of strong antioxidant, $IC_{50}$ value of control 98.17 mg/L.

Metabolomics is the systematic study of global metabolite profiles in biological samples. Metabolomics analysis is an approach used to determine metabolite profiles. Untargeted GCMS is considered an appropriate method for analyzing metabolites in liquid samples. Based on Fig 1, the five species of the *Solanum* genus show chromatograms with different numbers of identified compounds, namely *Solanum tuberosum*, 20 compounds; Solanum *betaceum*, 18 compound and *Solanum melongena*, 16 compounds. Compounds found in the five types of plants include propane, acetaldehyde, 4H-Pyran-4-one-TU, 2-propen-1-ol, propen and 1,2 ethanediamine. Based on (Fig 2), it can be seen that there are differences in the chromatograms of the *Capsicum* genus species. In *Capsicum annuum*, 37 compounds were obtained and 21 compounds were found. Compounds identified in the two types of plants include capsaicin, acetamide, ethylene oxide, butanoic acid and acetic acid. The diversity of chromatograms obtained in the Solanaceae tribe shows differences in specific phenotypic and genotypic characteristics and can also be influenced by external factors. Applying elicitors can increase growth, including synthesizing potential antioxidant compounds. The results of research by Mendoza *et al.* (2018) showed that the application of 300 μM salicylic acid and 3 μM methyl jasmonate increased the content of fenolic compounds and flavonoids and induced the phenylpropanoid metabolite pathway.

Antioxidant activity is known from the $IC_{50}$ value. The $IC_{50}$ value is inversely proportional to the antioxidant content. A material that has a small $IC_{50}$ value indicates that the

material has a large antioxidant content because $IC_{50}$ represents the sample concentration required to reduce 50% of the absorbance of a reducible DPPH solution (Rusmana *et al.,* 2017). Solanaceae family plants have different antioxidant potentials from medium to very high categories. Antioxidant properties are closely related to metabolite compounds. Some classic non-enzymatic antioxidants are vitamins such as vitamin C or E, but flavonoids and other phenolic compounds are also potent antioxidants.

## CONCLUSION

The results of non-targeted metabolomic analysis with GCMS and antioxidant activity with $IC_{50}$ of the *Solanum* genus in *Solanum betaceum*, *Solanum melongena*, *Solanum torvum*, *Solanum lycopersicum*, *Solanum tuberosum* identified 18, 16, 14, 13 and 20 compounds respectively. The $IC_{50}$ values were 51.32 mg/L (strong category), 56.64 mg/L (strong category), 65.86 mg/L (strong category), 113.60 mg/L (medium category), 102.71 mg/L (medium category), respectively. In the *Capsicum* genus, namely *Capsicum frutescens* and *Capsicum annuum*, 21 and 37 compounds were identified respectively, with $IC_{50}$ values of 124.76 mg/L (medium category) and 63.23 mg/L (strong category). In *Physalis angulata* 19 compounds were identified, $IC_{50}$ was 44.04 mg/L (very strong category), 6 compounds were identified in *Nicotiana tabacum*, $IC_{50}$ was 67.55 mg/L (strong category). Solanaceae plants have moderate to very strong antioxidant potential.

## ACKNOWLEDGEMENT

The present study was supported by the Directorate of Higher Education, Ministry of Education, Culture, Research and Technology of the Republic of Indonesia and Sam Ratulangi University for providing research funding through Fundamental Research in 2024.

### Conflict of interest

The authors declare no conflict of interest.

## REFERENCES

Al-Lahham, S., Sbieh, R., Jaradat, N., Almasri, M., Mosa, A., Hamayel, A. and Hammad, F. (2020). Antioxidant, antimicrobial and cytotoxic properties of four different extracts derived from the roots of *Nicotiana tabacum* (L.). *European Journal of Integrative Medicine*. **33(2):** 101039.

Ananta, I.G.B.T. and Anjasmara, D.G.A. (2022). Antioxidant and antibacterial pof red chillies extract (*Capsicum annum* var. Longum). *Medicamento Scientific Journal.* **8:** 48-55.

Artanti, A.N. and Lisnasari, R. (2018). Test of the antioxidant activity of *Solanum* family leaf ethanol extract using the DPPH free radical reduction method. *Journal of Pharmaceutical Science and Clinical Research.* **3:** 62-69.

Azzikri, Wardana, S.T. and Harmida. (2020). Ethnobotany of solanaceae medicinal plants in the Kerinci Tribe community in the Lempur Region, Kerinci Regency, Jambi Province. *Sribios: Sriwijaya Bioscientia.* **2:** 172-179.

Beniwal, R., Singh, S., Devi, P. (2022). Effect of extraction solvents on phytochemicals and antioxidant potential of turnip roots *Brassica rapa* (L.). *Agricultural Science Digest*. 1-5. doi: 10.18805/ag.D-5551.

Devi, R., Singh, S., Moond, M., Beniwal, R. and Kumar, S. (2023). Phytochemical screening and antioxidant potential of *Syzygium cumini* leaves. *Agricultural Science Digest*. 1-6. doi: 10.18805/ag.D-5628.

Dewi, L., Slamet, T. and Zianida, N. (2024). Antioxidant activity of ethanol extract of black potato [*Plectranthus rotondifolius* (Poir) Spreng] leaves using the DPPH (2-2-diphenyl-1-picrylhydrazyl) method. *Pharmaceutical Scientific Journal.* **12:** 1-7.

Diaz, M.R., Lobos, T., Cardemil., Nesi, A.N., Retamales, J., Jaakola, L., Alberdi, M. and Fonseca, A.R. (2016). Methyl jasmonate: An alternative for improving the quality and health properties of fresh fruits. *Molecules*. **21:** 567.

Gomi, K. (2020). Jasmonic acid: An essential plant hormone. *Int. J. Mol. Sci.* **21:** 1261.

Handayani, S., Najib, A. and Wati, N.P. (2018). Test the antioxidant activity of daruju *Acanthus ilicifolius* (L.) leaf extract using the 1,1-diphenyl-2-picrylhidrazil free radical scavenging method. *Indonesian Phytopharmaceutical Journal (JFFI)*. **5:** 299-308.

Hasfikasari, P., Faradiba and Amin, A. (2024). Review article: Antioxidant activity of tomato fruit extract (*Solanum lycopersicum* L.). *Makassar Natural Product Journal.* **2:** 43-50.

Helilusiatiningsih, N. and Soenyoto, E. (2020). Analysis of bioactive antioxidant compounds and nutrients in pokak eggplant (*Solanum torvum*) fruit as a functional food ingredient. *Buana Sains*. **20:** 7-19.

Hellmann, H., Goyer, A. and Navarre, D.A. (2021). Antioxidants in potatoes: A functional view of one of the world's major food crops. *Molecule.* **26(9):** 2446.

Huang, H., Liu, B., Liu, L. and Song, S. (2017). Jasmonate action in plant growth and development. *Journal of Experimental Botany.* **68:** 1349-1359.

Khatun, S., Roy, T.S., Haque, M.N. and Alamgir, B. (2016). Effect of plant growth regulators and their time of application on yield attributes and quality of soybean. *Int. Journal of Plant and Soil Science*. **11:** 1-9.

Krisnawati, Y. and Febrianti, Y. (2019). Identification of solanaceae family plants found in Tugumulyo District. *BIOSFER, J. Bio. and Biology Education.* **4(2):** 73-84.

Kumari, V., Kumar, D. and Bhardmaj, R. (2023). Metabolome: Analysis, nutrient and antioxidant potential of aerial and underground parts of *Ajuga parviflora* Benth. *Microchemical Journal*. **187:** 108451.

Mbida, H., Tsala, D.E., Aboubakar, S., Habtemariam, S., Edmond, J.J., Bakwo, E.F. and Minkande, J.Z. (2022). Research article: Antioxidant activity of aqueous extract of leaves and seeds of *Datura metel* (Solanaceae) in frog's heart failure model. Hindawi Evidence-Based Complementary and Alternative Medicine. Article ID: 5318117.

Mendoza, D., Cuaspud, O., Arias, J.P., Ruiz, O. and Arias, M. (2018). Effect of salicylic acid and methyl jasmonat in the production of phenolic compounds in plant cell suspension cultures of *Thevetia peruviana*. Biotechnol REp (Amst). **19:** e00273.

Misfadhila, S., Wulandari, N., Yetti, R.D., Sarina, G. and Haris, M. (2022). Test of the antioxidant activity of ethanol extract of red cayenne pepper [*Capsicum annuum* Var. Frutescens (L.) Kuntze] using the ferric reducing antioxidant power (FRAP) method. *Higea Pharmacy Journal.* **14:** 50-57.

Molyneux, P. (2004). The use of the stable free radical diphenylpicrylhydrazyl (DPPH) for estimating antioxidant activity. Songklanakarin. *J. Sci. Technol.* **26:** 211-219.

Nguyen, K.V., Pongkitwitoon, B., Pathomwichaiwat, T., Viboonjun, U. and Prathanturarug, S. (2019). Effects of methyl jasmonate on the growth and triterpenoid production of diploid and tetraploid *Centella asiatica* (L.) Urb. hairy root cultures. *Scientific Reports.* **9:** 18665.

Pant, H.C., Rautela, I., Pant, H.V., Kumar, A., Kumar, P., Fatima, K. and Gaurav, N. (2024). Comparison of antioxidant properties and flavonoid of natural and *in vitro* cultivated Nardostachys Jatamansi. *Agricultural Science Digest.* **44(3):** 406-413. doi: 10.18805/ag.D-5654.

Pujiastuti, A. and Kristen, M. (2019). Formulation and mechanical stability test of hand and body lotion tomato juice (*Licopersicon esculentum* Mill.) as an antioxidant. *Indonesian Pharmaceutical Journal.* **16:** 42-55.

Putra, G.N.T.M. and Astuti, N.M.W. (2023). Review: Study of phytochemical content, antioxidant activity and toxicity of ciplukan (*Physalis angulata* L.). *COMSERVA: (Journal of Research and Community Service)*. **3:** 2168-2179.

Ralte, L., Bhardwaj, U. and Sing, T. (2021). Traditionally used edible solanaceae plants of Mizoram, India have high antioxidant and antimicrobial potential for effective phytopharmaceutical and nutraceutical formulations. *Heliyon.* **7(9):** e07907.

Rampe, H.L., Siahaan, R. and Moniaga, W.M. (2022). Potential of exogenous methyl jasmonate elicitor as an agent to induce resistance of sweet potato (*Ipomoea batatas* L.) to herbivorous insect pests. Sam Ratulangi University Flagship Applied Research. pp: 62.

Rampe, H.L., Siahaan, R. and Pontororing, H.H. (2023). Analysis of genetic diversity and genetic relationships of sweet potato (*Ipomoea batatas* L.) based on morphological characters after induction of the elicitors methyl jasmonate and salicylic acid. Sam Ratulangi University Flagship Basic Research. pp: 48.

Ruan, J., Zhou, Y., Zhou, M., Yan, J., Khurshid, M., Weng, W., Cheng, J. and Zhang, K. (2019). Jasmonic acid signaling pathway in plants. *Int. J. Mol. Sci.* **20(10):** 2479.

Rusmana, D., Wahyudianingsih, R., Elisabeth, M., Balqis, Maesaroh and Widowati, W. (2017). Antioxidant activity of *Phyllantus niruri* extract, rutin and quercetin. *The Indo. Biomed. J.* **9:** 84-90.

Sanchez, L.K.R., Bernala, J.E.P., Torres, M.A.S., Casas, X.M., Suarez, L.E.C., Rodriguez, J.A.P. and Ladinoa, O.J.P. (2020). Effect of methyl jasmonate and salicylic acid on the production of metabolites in cell suspensions cultures of *Piper cumanense* (Piperaceae). *Biotechnology Reports*. **28:** e00559.

Siddiqi, K.S. and Husen, A. (2019). Plant response to jasmonates: Current developments and their role in changing environment. *Bulletin of the National Research Centre.* **43:** 153.

Sukmawati, Rahmawati and Aviah, A.A. (2024). Antioxidant activity of ciplukan (*Physalis angulata* L.) stem ethanol extract using DPPH. *Makassar Pharmaceutical Science Journal.* **1:** 373-381.

Tam-Ho, T., Murthy, H.N. and Park, S.Y. (2020). Methyl jasmonate induced oxidative stress and accumulation of secondary metabolites in plant cell and organ cultures. *Int. J. Mol. Sci.* **21:** 716.

Wafa, N. (2024). Total phenolic, flavonoid contents and antioxidant activity of ethanolic extracts (leaves and fruit) of *Cucurbita maxima* Duch. ex Lam. *Agricultural Science Digest*. **44(1):** 114-117. doi: 10.18805/ag.DF-463.

Zou, X., Amrit, B.K., Abdur, R., Saeed, M., Al-Awthan, Y.S., Al-Duais, M.A., Bahattab, O., Khan, M.H. and Suleria, H.A.R. (2021). Screening of polyphenols in tobacco (*Nicotiana tabacum*) and determination of their antioxidant activity in different tobacco varieties. *ACS Omega.* **6(39).** https://doi.org/10.1021/ACSOMEGA.1C03275