REVIEW ARTICLE Indian Journal of Agricultural Research, Volume 59 Issue 2: 179-186 (February 2025)

Salinity Tolerance Mechanisms in Rice: A Review

O.P. Reshna[1], R. Beena[1]

10.18805/IJARe.A-5760

ABSTRACT

Rice (*Oryza sativa* L.) is one of the most important cereal crop globally and it is the staple food of more than half of world population. Current average yield is 10 to 15% lower than its potential yield. There are many reasons for this yield gap such as biotic and abiotic stresses, management strategies as well as nutrient deficiencies. Salinity is one of the most serious factors limiting the productivity of rice, with adverse effects on germination, plant vigor and crop yield. This salinity may be natural or induced by agricultural activities such as irrigation or the use of certain types of fertilizer.Salinity induced ionic imbalance and osmotic stress affect water and nutrient uptake, stomatal closure, gas exchange, photosynthesis, transpiration, carbon assimilation and hence decrease in rice yield. Studying the response of rice at physiological, genetic and molecular level is the mandate to develop salt tolerant rice varieties. This review described the impact of salt stress in rice, various types of salt tolerance mechanisms in rice, including the ion homeostasis, production of compatible organic solutes, antioxidative genes, salt responsive regulatory elements, role of ion transporters and channel proteins. Further, the future perspective of developing salt-tolerant varieties using landraces *via*., marker assisted breeding, genome editing tool, utilization of beneficial microorganisms.

Key words: Molecular mechanism, Physiological mechanism, Rice, Salinity, Transporters.

Rice is one of the world's most important cereal crop which is grown over 156 mha area globally and is staple food crop of more than 50 per cent population worldwide (FAO, 2008). To maintain food security, rice production must be increased quantitatively and qualitatively to meet the requirements of the growing population in the twenty-first century (Rejeth *et al*., 2020). Although rice is adapted to a wide range of ecosystems it is susceptible to climatic changes leading to low rice productivity (Beena, 2005). Changing climate leads to periods of drought, frequent floods, sea water inundations, *etc*. which reduce the yield potential of current rice varieties. Among abiotic stresses, salinity is the second most devastating constraint in rice production after drought, affecting approximately 1 billion ha of land globally. Soil salinity affects about 800 million hectares of arable lands worldwide (Munns and Tester, 2008). Excessive use of irrigation water with improper drainage, poor quality irrigation water containing an excess level of salts, and flooding from seawater are the causes of salinity. Direct accumulation of salts disturbs metabolic processes and all major morpho-physiological and yield-related traits including tiller number, panicle length, spikelet number per panicle (Khatun *et al*., 1995), grain filling (Rao *et al*., 2013), plant biomass (Zeng *et al*., 2007) and photosynthesis (Baker, 2008), leading to significantly decreased yield (Flowers and Yeo, 1981).

Salinity stress triggers a wide variety of plant responses, ranging from morphological, physiological, biochemical and molecular changes to plant system. Salt stress is first perceived by the root system and impairs plant growth both in the short term, by inducing osmotic stress caused by reduced water availability, and in the long term, by salt-induced ion toxicity due to nutrient imbalance in the cytosol. Therefore, the two main threats imposed by salinity are induced by osmotic stress and ionic toxicity associated with excessive Cl^- and Na^+ uptake, leading to Ca^{2+} and K^+ deficiency and to other nutrient imbalances (Marschner, 1995). In addition, salt stress is also manifested as oxidative stress mediated by ROS. All these responses to salinity contribute to the deleterious effects on plants. However, tolerance can come with trade-offs; for example, it has long been known that stress-tolerant plants have reduced growth rates and productivity (Beena *et al*., 2018c; Nithya *et al*., 2021). Hence, understanding the basis of tolerance is also important to understand the trade-offs between tolerance and growth/ productivity for potent crop improvement.

[1]Department of Plant Physiology, College of Agriculture, Vellayani Kerala Agricultural University, Vellayani-695 522, Kerala, India.

Corresponding Author: R. Beena, Department of Plant Physiology, College of Agriculture, Vellayani Kerala Agricultural University, Vellayani-695 522, Kerala, India.
Email: beenaajithkumar@gmail.com

How to cite this article: Reshna, O.P. and Beena, R. (2025). Salinity Tolerance Mechanisms in Rice: A Review. Indian Journal of Agricultural Research. 59(2): 179-186. doi: 10.18805/IJARe.A-5760.

Submitted: 02-03-2021 **Accepted:** 05-07-2021 **Online:** 09-08-2021

Impacts of salinity stress on rice growth and development

Salt stress severely influences the morphology of rice plants which comprise of stunted plant growth, chlorosis, leaf burning, leaf rolling, poor root growth, wilting and senescence (Zhang *et al*., 2012). Salinity bring down the photosynthetic rate and associated parameters such as transpiration,

stomatal conductance and intercellular CO_2, concentration (Munns, 2002). Salinity affects the pollination, fertilization and causes spikelet sterility in rice (Pearson and Bernstein,1959); delays heading and affect yield components (Grattan *et al*., 2002).

The biochemical impacts typically comprise of oxidative stress and ionic stress. Salt stress increases ROS production which can elevate oxygen-induced cellular damage. With the rise in salinity level, indices of oxidative stress *viz*., superoxide radical and H_2O_2, content increased in leaves of all the cultivars and this increase was more notable in salt-sensitive varieties and non-significant in resistant cultivars. There was 10 fold increase in Na^+ concentration in salt sensitive "IR29" compared to salt tolerant rice variety "Pokkali" (Golldack *et al*., 2003). Elevated concentration of cytoplasmic Na^+ and reduction of K^+ result in changes of membrane potential, osmotic pressure, turgor pressure, calcium signaling, reactive oxygen species signaling, transcriptional regulation, alteration of gene expression, modification of protein expression pattern and spectra of *siRNA*s, signaling molecules, phytohormones and metabolites. NaCl at a concentration as low as 50 mM is lethal at the seedling stage (Yeo *et al*., 1990). High sodium concentrations over 100 mM often have inhibiting effect on protein synthesis at least in salt-sensitive glycophytes like rice (Flowers and Yeo, 1981).

Na^+ is the principal poisonous ion in salinized soil and low Na^+/K^+ ratio in the cytoplasm is essential to maintain a number of enzymatic reactions. Plants under salt stress normally absorb Na^+ and simultaneously inhibit K^+ absorption. High salt domains can shatter the ion homeostasis of plant cells, knock down the ionic balance and affect the distributions of ion at whole plant levels under salinity, increased osmotic pressure of the soil solution which restricts the water absorption and entry of water into the seeds, interruption of nutrient uptake, induces stomatal closure (Almansouri *et al*., 2001).

Salinity alters a wide array of metabolic processes in growing plants and induces changes in contents and activities of many enzymes. Sugar metabolism is critically affected when the plant grown under salinity condition. Under salinity, the starch content in roots declined more in salt sensitive *cvs*. Ratna and Jaya than in salt tolerant *cvs*. CSR-1 and CSR-3. The content of reducing and non-reducing sugars and the activity of sucrose phosphate synthase was increased more in the sensitive than in the tolerant cultivars. Acid invertase activity decreased in shoots of the salt tolerant cultivars, whereas increased in salt sensitive cultivars. Starch phosphorylase activity decreased in all cultivars (Dubey and Singh, 1999). To better understand the impacts of salinity on plant growth, it is necessary to study the changes in metabolites involved in respiratory metabolism. A considerable reduction in saccharide abundance, including glucose, fructose, xylose, and sucrose has been reported in rice exposed to salinity (Sanchez *et al*., 2008). These saccharides are required as sources of glucose for glycolysis and may also contribute to osmotic adjustment in plants.

Mechanisms for salinity tolerance in rice

It is essential to know the basic molecular mechanisms of salt tolerance for the crop improvement programs to develop high yielding salt tolerant varieties. Salt tolerance is regarded as a complex quantitative trait which is controlled by multiple genes. Many adaptive responses at molecular, cellular and physiological level results in tolerance or resistance of rice plant to salt stress. There are mainly three mechanisms observed in rice.

Ion exclusion

Selective uptake of salt by root cells is the initial strategy followed by rice plant to tackle the salinity stress. Rahman *et al*. (2001) reported that root cap cell expansion is an adaptive response to salinity in rice plants and this root cap cells may function in Na^+ exclusion under salinity stress. Once water is taken up by the root hair, it travels from cell to cell through intracellular regions across the cortex to handover the salt in xylem through apoplastic pathway. Suberin lamellae and casparian strip limit and force both water and salts to cross the plasma membrane into the cytoplasm, where they can continue on the symplast pathway, leading to decrease in Na^+ accumulation in the shoot (Cai *et al*., 2011). With increase in NaCl concentration in rhizosphere, there will be a fall off of transpiration rate which lowers water potential in roots, and the transport of ABA from root to shoot and give rise to stomatal closure and prevent dehydration (Zhang *et al*., 2006).

Osmotic stress tolerance

Osmotic tolerance is the plant's ability to tolerate the physiological drought aspect of salinity stress and to maintain leaf expansion and stomatal conductance by maintaining the tissue turgor (Rajendran *et al*., 2009). The early symptom of salinity stress on plants is osmotic stress which arise due to the accumulation of salts in the rhizosphere that affects negatively the plant growth (Munns and Tester, 2008). Compatible solutes accumulate in the cytosol, contribute to the decrease of cytoplasmic water potential and act as osmoprotectants. Rice plant under osmotic stress accumulates more carbohydrates or trehalose than proline (Nounjan and Theerakulpisut, 2012). Trehalose bring down Na^+ accumulation, growth inhibition and salt (an osmotically regulated gene) expression at low to medium concentration and at higher concentration hinder NaCl-induced loss of chlorophyll in blades, enhance rice growth and preserve root integrity (Garcia *et al*., 2012). Karthikeyan *et al.* (2011) observed high accumulation of proline in salt tolerant transgenic rice plants with expression of P5CS. *codA* introduced into transgenic rice plants promotes the synthesis of glycine betaine and enhancement of salt tolerance in plants (Su *et al*., 2006). Li *et al.* (2011) demonstrated that overexpression of *OsTPS1* in rice plants enhances salt tolerance by increasing the amount of trehalose and proline.

Tissue tolerance

Tissue tolerance is achieved by sequestration of Na^+ in the vacuole, synthesis of compatible solutes, production of enzymes catalyzing detoxification of reactive oxygen species. Reactive oxygen species (ROS) are dual functioning molecules which act both as toxic compounds as well as signal transduction molecules in response to abiotic as well as biotic stresses in plants (Miller *et al*., 2010). Under stress condition, there will be accumulation of plenty of ROS which will damage protein structure cause lipid peroxidation and mutation of DNA (Vaidyanathan *et al*., 2003). The antioxidant enzymes catalase (CAT) and ascorbate peroxidase (APX) are efficient scavengers of hydrogen peroxide (H_2O_2) and thus inhibit membrane lipid peroxidation, which is induced by high concentrations of H_2O_2. This is a key factor in reducing the effects of salinity stress (Mittler, 2007). Under saline condition, the capacity to scavenge reactive oxygen species decreased with age, and thus the apical region of the leaf blade suffered severe damage by Na^+ than the basal region in salt sensitive rice variety (Yamane *et al*., 2009). In salinity tolerant rice cultivar, gene expression and the activity of antioxidant enzymes (SOD, CAT, APX and GR) were increased and protected from oxidative stress. Under salinity stress, salt tolerant rice cultivar Pokkali was reported to have higher activity of ROS scavenging enzymes such as catalase (CAT), ascorbate (ASC) and glutathione (GSH), compared to the salt sensitive rice cultivar (Pusa Basmati). In rice during salinity stress, cell death advances in a well-regulated manner and causes PCD in the root tip of rice (Liu *et al*., 2007). Transgenic rice plants containting antioxidative genes *OsECS*, *OsVTE1* and *OsMSRA4.1* showed high salt tolerance (Choe *et al.,* 2013).

Role of membrane transporters for salt tolerance in rice

Transporting Na^+ out of the cytoplasm into apoplast or vacuole, which is driven, respectively, by Na^+/H^+ antiporters in plasma membrane or tonoplast and exclusion of Na^+ from sensitive shoot tissues are the basic strategies for the salt stressed plants to alleviate the Na^+ toxicity (Hamamoto *et al*., 2015). Transporters could be considered as enzymes where conformational changes of a protein molecule are required for a complete transport cycle of ions (Gadsby, 2009).

Turnover rate of the transporter is the number of complete transport cycles performed per second. There are three main types of transporters involved in salt tolerance mechanism of rice; Na^+ transporters, such as vacuolar Na^+/H^+ antiporters (*NHX*s), salt overly sensitive 1 transporter (*SOS*1) and high-affinity potassium transporters (*HKTs*).

Vacuolar NHX transporters

The coupled exchange of K^+ or Na^+ for H^+ is mediated by a family of transporters known as *NHXs (NHX*1-6). The cytosolic Na^+ is compartmentalize Na^+ into the vacuoles by *NHXs* (Bassil and Blumwald, 2014). The compartmentation of Na^+ into vacuoles not only reduces the toxicity of Na^+ in the cytosol but also contributes to osmotic regulation. First evidence of *NHX* transporter in tonoplast vesicles from rice roots was reported by Fukuda *et al*. (1998). The mechanism of compartmentalization are the pH gradient generated by proton-translocating *H^+-ATPase* and proton translocating inorganic pyrophosphatase (*H^+- PPase*) on the tonoplast and that the amount of antiporter in vacuolar membranes. Later, Fukuda *et al*. (2004) investigated the function and intracellular localization of the product of the *OsNHX1* cloned from rice. Their results indicate that the *OsNHX1* gene encodes a vacuolar (Na^+, K^+)/H^+ antiporter. In addition, overexpression of *OsNHX1* improved the salt tolerance of transgenic rice cells and plants, suggesting that *OsNHX1* on the tonoplast plays important roles in removing accumulated Na^+ and K^+ in the cytoplasm into the vacuoles. Up to now, four vacuolar NHXs (OsNHX1-4) have been reported in rice (Kumar *et al*., 2013).

***SOS* transporter**

SOS1 transporter was first isolated from rice (Martinez-Atienze *et al*., 2007). Based on genetic and biochemical criteria, they identified that *OsSOS*1 is a rice plasma membrane Na^+/H^+ antiporter and is functionally equivalent to the Arabidopsis *SOS1. H^+-ATPase* present in plasma membrane is responsible for Na^+ efflux from cytoplasm. The energy released from the hydrolysis of ATP is used to pump H^+ out of the cell, generating electrochemical H^+ gradient. The proton-motive-force generated by the H^+-ATP*ase* operates the Na^+/H^+ antiporters, which couple the movement of H^+ into the cell along its electrochemical gradient to the efflux of Na^+ against its electrochemical gradient.

In salt overly sensitive (*SOS*) pathway, salt stress elicited Ca^{2+} signals are perceived by Ca^{2+} binding protein *SOS*3 and activates *SOS*2, a serine/ threonine protein kinase (Halfter *et al*., 2000). Then the *SOS2-SOS3* complex phosphorylates *SOS1* which is a Na^+ /H^+ antiporter on cell membrane that extrudes Na^+ out of the cytosol (Guo *et al*., 2004). The transport level of *SOS1* is controlled by *SOS3-SOS2* kinase complex. When there is a higher accumulation of Na^+ in cytosol, *SOS*2 activates the tonoplast Na^+/H^+ antiporter that sequesters Na^+ into the vacuole (Blumwald, 2000). Elevation of salinity influences the Na^+ /H^+ antiporter activity and the activity is more significant in salt tolerant species than in salt sensitive ones.

***HKT*1 function in ion transport**

Sodium concentration in shoots is affected by the regulation of sodium transport from root to shoot by the family of *HIGH-AFFINITY POTASSIUM TRANSPORTERS (HKTs*). The members of this family can be divided into two classes according to their affinity for either sodium or potassium, caused by a single amino acid substitution in the pore region of the transporter. The *HKT*1 subclass transports sodium, while the HKT2 subclass transports potassium. *OsHKT*2; 2 is present in the salt-tolerant 'Pokkali' cultivar, but absent in 'Nipponare', can mediate both Na^+ and K^+ influx. This would explain why 'Pokkali' cutivar is salt tolerant (Garciadeblas *et al*., 2003). Identified three hybrid *HKT* gene (named *OsHKT2;2/1; OsHKT2*;1/2) in 'Nona Bokra' and *OsHKT*2;2

from 'Pokkali' (Oomen *et al*., 2012). These genes may contribute to the salt tolerance by maintaining K^+ uptake under salinity conditions.

HAK/KUP family transporters

Importance of K^+ retention ability in root and leaf tissues, role of HAK/KUP family of transporters in K^+ uptake and its possible linkage with Ca^{2+} and ROS signaling, control of xylem K^+ loading control of phloem K^+ recirculation in salt stressed plants (Wu *et al*., 2018).

Role of ion channel proteins for salt tolerance in rice

Ion channels in an open state/conformation allow passage of over 106-108 ions per second via a selective pore formed within a protein molecule. The diameter of the pore is determined by the molecular structure of ion channel. The diameter of the pore and amino acids lining it essentially determine the ion selectivity of ion channel and potential number of passing ions per unit of time. Some of the important channel proteins present in rice are: *OsAKT1* (K^+ inwardrectifying channel), *OsKCO*1 (K^+ outward-rectifying channel), *OsTPC*1 (Ca^{2+} permeable channel), *OsCLC*1 (Cl^- channel) and *OsNRT1*;2 (nitrate transporter).

Role of protein kinases for salt tolerance in rice

Important protein kinases involved in salt tolerance mechanism in rice are Mitogen-activated protein kinases (*MAPKs*), sucrose non-fermenting1-related kinase 2 (*SnRK2*) family members have been designated as stressactivated protein kinases (*OsSAPK*), receptor-like cytoplasmic kinases (*RLCKs*). Over-expression of *OsMAPK*5 (encoding an abscisic acid-inducible MAPK), enhances salt tolerance in rice (Xiong and Yang, 2003). *OsMAPK*44, reduced the salt injury by maintaining ion homeostasis in transgenic rice (Jeong *et al*., 2006). *OsMAPK*33 exhibits salinity tolerance in transgenic rice by altered expression of ion transport genes such as the K^+/H^+ antiporter or other downstream positive regulators of salt tolerance in the MAPK pathway (Lee *et al*., 2011). The expressions of all the ten *OsSAPK*s (stress activated protein kinases) are induced under salt stress conditions by osmotic stress and/or ABA dependent signaling (Cutler *et al*., 2010). Overexpression of *SAPK4* improved germination, growth and development in seedlings and mature rice plants under salinity following alteration in the expression of genes involved in ion homeostasis and oxidative stress responses (Diédhiou *et al*., 2008). Overexpression of *SAPK6* in rice up-regulates the expression of the enzymes involved in diverse salt stress-responsive metabolic pathways under salt stress conditions.

Role of transcription factors and miRNAs for salt tolerance in rice

Transcription factors (TFs) are protein that controls the rate of transcription of genetic information from DNA to messenger RNA, by binding to a specific DNA sequence. Several salt responsive TFs were identified in rice, *OsCOIN* (Liu *et al*., 2007), *OsNAC*6 (Nakashima *et al*., 2007), *OsNAC*5 (Takasaki *et al*., 2010), *OsDREB2A* (Mallikarjuna *et al*., 2011), *OsMYB*2 (Yang *et al*., 2012), *OsTZF*1 (Jan *et al*., 2013), *OsBZIP*71 (Liu *et al*., 2014) which improve salt tolerance by various mechanisms like the accumulation of osmoprotectants and antioxidants, increase transporter activity for Na^+ and K^+ ions, regulate the expression of other saltresponsive genes.


By modulating the gene expression at the post-transcriptional level, miRNAs can regulate plant development and respond to environmental stress (Table 1).

Approaches to improve salt tolerance in rice

Conventional breeding methods such as use of *in vitro* selection, interspecific hybridization and pooling physiological *etc*. were used to enhance salinity tolerance (Flowers, 2004). Superior structured design to breeding for salinity tolerance acquired when there was a better understanding of the function regulation and function of the responsive genes and their association to QTL regions (Negrao *et al*., 2012). Fig 1 shows the different approaches used for developing salt tolerant rice varieties. In India, there were many success stories in developing and releasing the salt tolerant cultivars including CSR10, CSR13, CSR27, Narendra usar 2 and Narendra usar 3 (Singh *et al*., 2004) and those varieties have been used to reclaim saline soils in India.

Table 1: List of salt stress-related rice miRNAs and their predicted target gene.

miRNA	Up-regulation	Down-regulation	Target gene	Reference
Osa-miR393	In leaves under salt and drought stress	In roots under salt and drought stress	*OsTIR1*, *OsAFB2*	Xia *et al*., 2012
Osa-miR408	Under heat, cold and salt stress		*OsUCL30*, *OsIAA6*	Sun *et al*., 2018
Osa-miR319a		Under drought and salt stress	*AsPCF5*, *AsPCF6*, *AsPCF8*, *AsTCP14*, *AsNAC60*	Zhou *et al*., 2013
Osa-miR820	In leaves under salt, heat and drought stress	In roots under salt, heat and drought stress	*OsDRM2*	Sharma *et al*., 2015
Osa-miR12477	In leaves and shoots under salt stress		*L-ascorbate oxidase*	Parmer *et al*., 2020



Platforms such as phenomics in tandem with combination of genomics-transcriptomics-proteomics-metabolomics need to be integrated for identifying key genes/regulators which can improve salinity tolerance in rice apart from general practices like usage of better agricultural practices and soil amelioration (Reddy *et al*., 2017).

Fig 1: Approaches for developing of salt tolerant rice plants.

CSR 2, CSR 3, CSR 13, CSR 22, CSR 23, CSR 26, CSR 27, CSR 30, Panvel 1, Panvel 2, Panvel 3, Vyttila 1 and Vyttila 2 are the varieties developed for the coastal saline soil conventionally (Shahbaz and Ashraf, 2013). Kerala Agricultural University has released 11 Vytilla series of rice verieties.

A major QTL *saltol* was identified by Gregorio and Senadhira in salinity tolerant Pokkali responsible for maintaining Na^+/K^+ homeostasis at the seedling stage (Gregorio *et al*., 2002). Marker-assisted backcross breeding (MABB) was attempted to incorporate the *Saltol* loci from Pokkali or its derivative FL478 facilitate the development of salinity-tolerant rice varieties (Valarmathi *et al.,* 2019). *Saltol* gene introgressed Pusa Basmati 1 (PB1) showed enhanced seeding stage salt tolerance (Singh *et al*., 2018). Under salinity stress, *saltol* introgressed backcross lines showed salt tolerance by maintaining good morphological and yield traits (Banumathy *et al*., 2018).

A salt-tolerant rice mutant *rst1 was* isolated and characterized which showed significantly lower lipid peroxidation and electrolyte leakage under salinity stress but higher shoot biomass and chlorophyll content (Deng *et al.* 2015). Microarray analysis of rice mutant *RGA1* disclosed its role in regulation of multiple abiotic stresses (Jangam *et al.* 2016). One mutant variety (Shua-92) and two mutants (IR8-202 and Pokkali-M) were developed from rice varieties IR8 and Pokkali through mutation breeding, evaluated to check the yield performance for two years under salinity stress and results showed that the mutant variety Shua-92 produced 40-49 per cent more paddy yield on saline soil than the famous salinity tolerant Nona Bokra and Pokkali varieties (Baloch *et al*., 2003).

Exploiting the potential of beneficial microorganisms is an alternative strategy for ameliorating plant stress tolerance. Stress alleviation is ensured by signaling events occurring during the plant–microbe interaction and these mechanisms are regulated by a complex network of signaling events (Smith *et al*., 2017). Recent studies showed that plant growth promoting bacteria act as growth promoter as well as elicitors of salinity tolerance (Tiwari *et al*., 2016).

There are many recent developments in the understanding of the mechanisms of salt tolerance such as RNA guided CRISPR endonuclease technology including base and prime editing or targeted mutagenesis which help in gene discovery, functional analysis transferring favorable alleles into elite breeding line to accelerate the breeding of salt-tolerant rice cultivars (Ganie, 2021). As a future line of study, the advanced technologies such as, genome editing can be used as an excellent tool, for improving crop traits, with salinity stress tolerance, as a target.

CONCLUSION

Na^+ exclusion from the root, modulation of root–shoot transport, and cellular compartmentalization of Na^+, as well as maintenance of cytoplasmic osmotic balance, are crucial aspects of salt tolerance. *HKT1* plays an important role in root stellar cells, modulating sodium transport to the shoot. Tissue-specific expression of *HKT*1 at specific developmental stages can contribute to plant salt tolerance. By targeting and phosphorylating downstream components, both ABA-independent *SUCROSE NONFERMENTING1-RELATED PROTEIN KINASE2 (SnRK2s*) and ABA-dependent *SnRK2s*

play pivotal roles in transcriptional regulation and posttranscriptional regulation during salt stress response.

Conflict of interest

All the authors declare that there is no conflict of interest for publishing this article.

REFERENCES

Almansouri, M., Kinet, J.M. and Lutts, S. (2001). Effect of salt and osmotic stresses on germination in durum wheat (*Triticum durum* Desf.). Plant and Soil. 231:243-254.

Baker, N.R. (2008). Chlorophyll fluorescence: A probe of photosynthesis *in vivo*. Annual Review of Plant Biology. 59: 89-113.

Baloch, A.W., Soomro, A.M., Javed, M.A., Bughio, H.U.R., Alam, S.M., Bughio, M.S., Mohammed, T. and Mastoi, N.U.N. (2003). Induction of salt tolerance in rice through mutation breeding. Asian Journal of Plant Science. 2(3): 273-276.

Banumathy, S., Veni, K., Anandhababu, R., Arunachalam, P., Raveendran, M., Vanniarajan, C. (2018). Character association and stress indices for yield components in *Saltol* introgressed backcross inbred lines of rice (*Oryza sativa* L.). Indian Journal of Agricultural Research. 52(1): 28-33. doi: 10.18805/IJARe.A-4926.

Bassil, E., Coku, A. and Blumwald, E. (2012). Cellular ion homeostasis: Emerging roles of intracellular NHX Na^+/H^+ antiporters in plant growth and development. Journal of Experimental Botany. 63: 5727-5740.

Beena, R. (2005). Studies on physio-morphological traits and genetic markers associated with drought tolerance in rice (*Oryza sativa* L.). Ph.D. thesis submitted to Tamil Nadu Agricultural University.

Beena, R., Praveenkumar, V.P., Vighneswaran, V. and Narayankutty, M.C. (2018c). Bulked line analysis: A useful tool to identify microsatellite markers linked to drought tolerance in rice. Indian Journal of Plant Physiology: 23(1)-7-15.

Blumwald, E. (2000). Sodium transport and salt tolerance in plants. Current Opinion in Cell Biology. 12(4): 431-434.

Cai, X., Chen, T., Zhou, Q.Y., Xu, L., Qu, L.Q., Hua, X.J., and Lin, J.X. (2011). Development of casparian strip in rice cultivars. Plant Signal Behavior. 6(1): 59-65.

Choe, Y.H., Kim, Y.S., Kim, I.S., Bae, M.J., Lee, E.J., Kim, Y.H., Park, H.M. and Yoon, H.S. (2013). Homologous expression of c-glutamylcysteine synthetase increases grain yield and tolerance of transgenic rice plants to environmental stresses. Journal of Plant Physiology. 170(6): 610-618.

Cutler, S.R., Rodriguez, P.L., Finkelstein, R.R. and Abrams, S.R. (2010). Abscisic acid: Emergence of a core signaling network. Annu Rev Plant Biol. 61: 651-679.

Deng, P., Jiang, D., Dong, Y., Shi, X., Jing, W. and Zhang, W. (2015). Physiological characterisation and fine mapping of a salt-tolerant mutant in rice (*Oryza sativa*). Functional Plant Biology. 42(11): pp.1026-1035.

Diédhiou, C.J., Popova, O.V., Dietz, K.J. and Golldack, D. (2008). The *SNF*1-type serine-threonine protein kinase *SAPK*4 regulates stress-responsive gene expression in rice. BMC Plant Biology. 8: 49.

Dubey, R.S. and Singh, A.K. (1999). Salinity induces accumulation of soluble sugars and alters the activity of sugar metabolising enzymes in rice plants. Biologia Plantarum. 42: 233-239.

FAO (2008). FAO Land and Plant Nutrition Management Service.

Flowers, T.J. (2004). Improving crop salt tolerance. J. Exp. Bot. 55(396): 307-319.

Flowers, T.J. and Yeo, A.R. (1981). Variability in the resistance of sodium chloride salinity within rice (*Oryza sativa* L.) varieties. New Phytologist. 88: 363-373.

Fukuda, A.A., Nakamura, A., Tanaka, T.H., Miyao, A., Hirochika, H., Tanaka,Y., Yazaki,Y.,Ishikawa,T., Koike,S. and Tanaka Y. (1998). Na^+/H^+ antiporter in tonoplast vesicles from rice roots. Plant Cell Physiol. 39: 196-201.

Gadsby, D.C. (2009). Ion channels versus ion pumps: The principal difference, in principle. Nat. Rev. Mol. Cell Biol. 10: 344-352.

Ganie, S.A., Wani, S.H., Henry, R. and Hensel, G. (2021). Improving rice salt tolerance by precision breeding in a new era. Current Opinion in Plant Biology. 60: 101996.

Garcia, M.S., Tellez, L.I., Merino, F.C.G., Caldana, C., Victoria, E.D., and Cabrera, H.E.B. (2012). Growth, photosynthetic activity and potassium and sodium concentration in rice plants under salt stress. Acta Scientiarum. 34: 317-324.

Garciadeblas, B., Senn, M.E., Banuelos, M.A. and Rodriguez-Navarro, A. (2003). Sodium transport and HKT transporters: The rice model. Plant Journal. 801-34: 788.

Golldack, D., Su, H., Quigley, F., Kamasani, U.R., Munoz-Garay, C., Balderas, E., Popova, O.V., Bennert, J., Bohnert, H.J. and Pantoja, O. (2002). Characterization of a HKT-type transporter in rice as a general alkali cation transporter. Plant Journal. 31: 542-529.

Grattan, S.R., Zeng, L., Shannon, M.C. and Roberts, S.R. (2002). Rice is more sensitive to salinity than previously thought. California Agriculture. 56(6): 189-195.

Gregorio, G.B., Senadhira, D., Mendoza, R.D., Manigbas, N.L., Roxas, J.P. and Guerta, C.Q. (2002). Progress in breeding for salinity tolerance and associated abiotic stresses in rice. Field Crops Research. 76(2-3): 91-101.

Guo, Y., Qiu, Q.S., Quintero, F.J., Pardo, J.M., Ohta, M., Zhang, C., Schumaker, K.S. and Zhu, J.K. (2004). Transgenic evaluation of activated mutant alleles of *SOS2* reveals a critical requirement for its kinase activity and C-terminal regulatory domain for salt tolerance in *Arabidopsis thaliana*. Plant Cell. 16(2): 435-449.

Halfter, U., Ishitani, M. and Zhu, J.K. (2000). The *Arabidopsis SOS*2 protein kinase physically interacts with and is activated by the calcium-binding protein *SOS*3. Proceedings of National Academy of Science. 97(7): 3735-3740.

Hamamoto, S., Horie, T., Hauser,F., Deinlein, U., Schroeder, J.I. and Uozumi,N. (2015). *HKT* transporters mediate salt stress resistance in plants: From structure and function to the field. Current Opinion in Biotechnology. 32: 113-120.

Jan, A., Maruyama, K., Todaka, D., Kidokoro, S., Abo, M., Yoshimura, E., Shinozaki, K., Nakashima, K., Yamaguchi-Shinozaki, K. (2013). *OsTZF1*, a CCCH-tandem zinc finger protein, confers delayed senescence and stress tolerance in rice by regulating stressrelated genes. Plant Physiology. 161(3): 1202-1216.

Jangam, A.P., Pathak, R.R. and Raghuram, N. (2016). Microarray analysis of rice d1 (RGA1) mutant reveals the potential role of G-protein alpha subunit in regulating multiple abiotic stresses such as drought, salinity, heat and cold. Frontiers in Plant Science. 7: 11.

Jeong, M.J., Lee, S.K., Kim, B.G., Kwon, T.R., Cho, W.S., Park, Y.T., Lee, J.O., Kwon, H.B., Byun, M.O. and Park, S.C. (2006). A rice (*Oryza sativa* L.) MAP kinase gene, *OsMAPK44*, is involved in response to abiotic stresses. Plant Cell Tissue Organ Culture. 85: 151-160

Karthikeyan, A., Pandian, S.K. and Ramesh, M. (2011). Transgenic indica rice cv. ADT43 expressing a å1-pyrroline-5-carboxylate synthetase (*P5CS*) gene from *Vigna aconitifolia* demonstrates salt tolerance. Plant Cell, Tissure and Organ Culture. 107(3): 383-395.

Khatun, S., Rizzo, C.A. and Flowers, T.J. (1995) Genotypic variation in the effect of salinity on fertility in rice. Plant and Soil. 173: 239-250.

Kumar, K., Kumar, M., Kim, F.R., Ryu, H. and Cho, Y.G. (2013). Insight into genomics of salt stress response in rice. Rice. 6: 27.

Lee, S.K., Kim, B.G., Kwon, T.R., Jeong, M.J., Park, S.R., Lee, J.W., Byun, M.O., Kwon, H.B., Matthews, B.F., Hong, C.B. and Park, S.C. (2011). Overexpression of the mitogen-activated protein kinase gene OsMAPK33 enhances sensitivity to salt stress in rice (*Oryza sativa* L.). J of Bioscience. 36: 139-151.

Li, H.W, Zang, B.S., Deng, X.W. and Wang, X.P. (2011). Over expression of the trehalose-6-phosphate synthase gene *OsTPS1* enhances abiotic stress tolerance in rice. Planta. 234(5): 1007-1018.

Liu, C.T., Mao, B.G., Ou, S.J., Wang, W., Liu, L.C., Wu, Y.B., Chu, C.C. and Wang, X.P. (2014). OsbZIP71, a bZIP transcription factor, confers salinity and drought tolerance in rice. Plant Molecular Biology. 84: 19-36.

Liu, K.M., Wang, L., Xu, Y.Y., Chen, N., Ma, Q.B., Li, F. and Chong, K. (2007). Overexpression of OsCOIN, a putative cold inducible zinc finger protein, increased tolerance to chilling, salt and drought, and enhanced proline level in rice. Planta. 226: 1007-1016.

Liu, S.K., Cheng, Y.X., Zhang, X.X., Guan, Q.J., Nishiuchi, S., Hase, K. and Takano, T. (2007). Expression of an NADP-malic enzyme gene in rice (*Oryza sativa* L.) is induced by environmental stresses; over-expression of the gene in *Arabidopsis* confers salt and osmotic stress tolerance. Plant Molecular Biolology. 64: 49-58.

Mallikarjuna, G., Mallikarjuna, K., Reddy, M.K. and Kaul, T. (2011). Expression of OsDREB2A transcription factor confers enhanced dehydration and salt stress tolerance in rice (*Oryza sativa* L.). Biotechnol Letters. 33: 1689-1697.

Marschner, H. (1995). Mineral Nutrition of Higher Plants, 2nd ed.; Academic Press: London, UK.

Martinez-Atienza, J., Jiang, X., Garciadeblas, B., Mendoza, I., Zhu, J.K., Pardo, J.M. and Quintero, F.J. (2007). Conservation of the salt overly sensitive pathway in rice. Plant Physiol. 143: 1001-1012.

Miller, G.A.D., Suzuki, N., Ciftci Yilmaz, S.U.L.T.A.N. and Mittler, R.O.N. (2010). Reactive oxygen species homeostasis and signalling during drought and salinity stresses. Plant Cell and Environment. 33(4): 453-467.

Mittler, R. (2007). Oxidative stress, antioxidants and stress tolerance. Trends Plant Sci. 7: 405-410.

Munns, R. (2002). Comparative physiology of salt and water stress. Plant Cell and Enviroment. 25: 239-250.

Munns, R. and Tester, M. (2008). Mechanisms of salinity tolerance. Annu. Rev. Plant Biol. 59: 651-681.

Nakashima, K., Tran, L.S.P., Van Nguyen, D., Fujita, M., Maruyama, K., Todaka, D., Ito, Y., Hayashi, N., Shinozaki, K. and Yamaguchi Shinozaki, K. (2007). Functional analysis of a NAC-type transcription factor OsNAC6 involved in abiotic and biotic stress-responsive gene expression in rice. Plant Journal. 51(4): 617-630.

Negrao, S., Almadanim, M.C., Pires, I.S., Abreu, I.A., Maroco, J, Courtois, B., Gregorio, G.B., McKnally, K.L., Oliviera, M.M. (2012). New allelic variants found in key rice salt-tolerant genes: An association study. Plant Biotechnology Journal. 1-14.

Nithya, N., Beena, R., Abida, P.S., Sreekumar, J., Roy, S., Jayalekshmi, V.G., Manju, R.V. and Viji, M.M. (2021). Genetic diversity and population structure analysis of bold type rice collection from Southern India. Cereal Research Communications. 49(2): 311-328.

Nounjan, N. and Theerakulpisut, P. (2012). Effect of exogenous proline and trehalose on physiological responses in rice seedlings during salt-stress and after recovery. Plant Soil and Environment. 58: 308-315.

Oomen, R.K.F.K., Benito, B., Sentenac, H., Rodriguez-Navarro, A., Talon, M., Very, A.A., Domingo, C. (2012) HKT2;2/1, a K+ permeable transporter identified in a salt-tolerant rice cultivar through surveys of natural genetic 762-polymorphism. Plant Journal. 71: 750.

Parmer, S., Garat, S.A., Tagirasa, R., Chandra, T., Behera, L., Dash, S.K., Shaw, B.P. (2020). Identification and expression analysis of miRNAs and elucidation of their role in salt tolerance in rice varieties susceptible and tolerant to salinity. PLOS ONE. 1-26.

Pearson, G.A. and Bernstein, L. (1959). Salinity effects at several growth stages of rice. Agronomy Journal. 51(11): 654-657.

Rahman, M.S., Matsumuro, T., Miyake, H. and Takeoka, Y. (2001). Effects of salinity stress on the seminal root tip ultra structures of rice seedlings. Plant Production Science. 4(2): 103-111.

Rajendran, K., Tester, M. and Roy, S.J. (2009). Quantifying the three main components of salinity tolerance in cereals. Plant Cell Environment. 32(3): 237-249.

Rao, P.S., Mishra, B. and Gupta, S.R. (2013) Effects of soil salinity and alkalinity on grain quality of tolerant, semi-tolerant and sensitive rice genotypes. Rice Science. 20: 284-291.

Reddy, I.N.B.L., Kim, B.K., Yoon, I. S., Kim, K.H. and, Kwon, T.R. (2017). Salt tolerance in rice: Focus on mechanisms and approaches. Rice Sci. 24(3): 123-144.

Rejeth, R., Manikanta, Ch.L.N., Beena, R., Roy, S., Manju,R.V. and Viji, M.M. (2020). Water stress mediated root trait dynamics and identification of microsatellite markers associated with root traits in rice (*Oryza sativa* L.). Physiology and Molecular Biology of Plants. 26(6): 1225-1236.

Sanchez, D.H., Siahpoosh, M.R., Roessner, U., Udvardi, M. and Kopka, J. (2008). Plant metabolomics reveals conserved and divergent metabolic responses to salinity. Physiologia Plantarumt. 132: 209-219.

Shahbaz, M. and Ashraf, M. (2013). Improving salinity tolerance in cereals. Crit Rev. Plant Sci. 32: 237-249.

Sharma, P., Jha, A.B., Dubey, R.S., Pessarakli, M. (2012). Reactive oxygen species, oxidative damage and antioxidative defense mechanism in plants under stressful conditions. Journal of Botany. 1-26.

Singh, R.K., Mishra, B. and Singh, K.N. (2004). Salt tolerant rice varieties and their role in reclamation programme in Uttar Pradesh. Indian Farming. 2: 6-10.

Singh, V.K., Singh, B.D., Kumar, A., Maurya, S., Krishnan, S.G., Vinod, K.K., Singh, M.P., Ellur, R.K., Bhowmick, P.K. and Singh, A.K. (2018) Marker-Assisted Introgression of *Saltol* QTL Enhances Seedling Stage Salt Tolerance in the Rice Variety "Pusa Basmati 1". International Journal of Genomics. Hindawi, 1-12. https://doi.org/10.1155/2018/8319879.

Smith, D.L., Gravel, V. and Yergeau, E. (2017). Editorial: Signaling in the phytomicrobiome. Frontiers in Plant Science. 8(611): 1-3.

Su, J., Hirji, R., Zhang, L., He, C.K., Selvaraj, G. and Wu, R. (2006). Evaluation of the stress-inducible production of choline oxidase in transgenic rice as a strategy for producing the stress-protectant glycinebetaine. Jornal Experimental Botany. 57(5): 1129-1135.

Sun, M.Z., Yang, J.K., Cai, X.X., Shen, Y., Cui, N., Zhu, Y.M., Jia, B.W. and Sun, X.L. (2018). The opposite roles of *OsmiR408* in cold and drought stress responses in *Oryza sativa*. Molecular Breeding. 38:120.

Takasaki, H., Maruyama, K., Kidokoro, S., Ito, Y., Fujita, Y., Shinozaki, K., Yamaguchi-Shinozaki, K., Nakashima, K. (2010). The abiotic stress-responsive NAC-type transcription factor OsNAC5 regulates stress-inducible genes and stress tolerance in rice. Molecular Genetics and Genome. 284: 173-183.

Tiwari, S., Lata, C., Chauhan, P.S. and Nautiyal, C.S. (2016). Pseudomonas putida attunes morphophysiological, biochemical and molecular responses in *Cicer arietinum* L. during drought stress and recovery. Plant Physiology and Biochemistry. 99: 108-117.

Vaidyanathan, H., Sivakumar, P., Chakrabarty, R. and Thomas, G. (2003). Scavenging of reactive oxygen species in NaCl-stressed rice (*Oryza sativa* L.) differential response in salt-tolerant and sensitive varieties. Plant Science. 165(6): 1411-1418.

Valarmathi, M., Sasikala, R., Rahman, H., Jagadeeshselvam, N., Kambale, R. and Raveendran, M. (2019). Development of salinity tolerant version of a popular rice variety improved white ponni through marker assisted back cross breeding. Plant Physiology Reports. 24(2): 262-271.

Wu, H., Zhang, X., Giraldo, J.P and Shabala, S. (2018). It is not about sodium: Revealing tissue specificity and signaling roles of potassium in plant responses to salt stress. Plant and Soil. 431: 1-17.

Xia, K.F., Wang, R., Ou, X.J., Fang, Z.M., Tian, C.G., Duan, J., Wang, Y.Q. and Zhang, M.Y. (2012). *OsTIR1* and *OsAFB2* downregulation *via OsmiR393* overexpression leads to more tillers, early flowering and less tolerance to salt and drought in rice. PLoS One. 7(1): e30039.

Xiong, L.Z. and Yang, Y.N. (2003). Disease resistance and abiotic stress tolerance in rice are inversely modulated by an abscisic acidinducible mitogen-activated protein kinase. Plant Cell. 15(3): 745-759.

Yamane, K., Mitsuya, S., Kawasaki, M., Taniguchi, M. and Miyake, H. (2009). Antioxidant capacity and damages caused by salinity stress in apical and basal regions of rice leaf. Plant Production Science. 12(3): 319-326.

Yang, A., Dai, X.Y., Zhang, W.H. (2012). A R2R3-type MYB gene, *OsMYB2*, is involved in salt, cold and dehydration tolerance in rice. Journal of Experimental Botany. 63(7): 2541-2556.

Yeo, A.R., Yeo, M.E., Flowers, S.A. and Flowers, T.J. (1990). Screening of rice (*Oryza sativa* L.) genotypes for physiological characters contributing to salinity resistance and their relationship to overall performance. Theortical and Applied Genetics. 79: 377-384.

Zelm, E., Zhang, Y. and Testerink, C. (2020). Salt Tolerance Mechanisms of Plants. Annual Review of Plant Biology. 71: 403-433.

Zeng, Y., Zhang, H., Li, Z., Shen, S., Sun, J., Wang, M., Liao, D., Liu, X., Wang, X., Xiao, F. and Wen, G. (2007) Evaluation of genetic diversity of rice landraces (*Oryza sativa* L.) in Yunnan. China. Breeding Science. 57: 91-99.

Zhang, J., Jia, W., Yang, J. and Ismail, A.M. (2006). Role of ABA in integrating plant responses to drought and salt stresses. Field Crops Research. 97: 111-119.

Zhang, Z., Liu, Q., Song, H., Rong, X. and Abdelbagi, M.I. (2012). Responses of different rice (*Oryza sativa* L.) genotypes to salt stress and relation to carbohydrate metabolism and chlorophyll content. African Journal of Agricultural Research. 7: 19-27.

Zhou, M., Li, D.Y., Li, Z.G., Hu, Q., Yang, C.H., Zhu, L.H. and Luo, H. (2013). Constitutive expression of a *miR319* gene alters plant development and enhances salt and drought tolerance in transgenic creeping bentgrass. Plant Physiology. 161(3): 1375-1391.