RESEARCH ARTICLE Indian Journal of Animal Research, Volume 58 Issue 2: 330-335 (February 2024)

Age and Sex Affect Carcass Traits and Meat Quality in Malpura Sheep

Shrawan Kumar Meel[1], Anurag Pandey[1], Y.P. Gadekar[2], Arvind Soni[2], Priyanka Meena[1], Asman Singh Gurjar[1], Vilshan Kumar Chauhan[1]

10.18805/IJAR.B-4864

ABSTRACT

Background: Small ruminants play a vital role in livelihood and nutritional security in India. The sheep population in the country is showing positive growth. Malpura sheep from the semi-arid region of India is an important mutton purpose breed.

Methods: Twenty-four animals *viz*., ram lamb (G-$_{Rl}$), ewe lamb (G-$_{El}$) (10-12 months each), ram (G-$_{Ra}$) and ewe (G-$_{Ea}$) (20-24 months each) were slaughtered to assess carcass traits and meat quality.

Result: The average pre-slaughter weight and dressing yield on an empty live weight basis for G-$_{Rl,}$ G-$_{El,}$ G-$_{Ra}$ and G-$_{Ea}$ groups were 23.83, 17.00, 35.82 and 28.08 kg; 52.35, 51.26, 53.70 and 52.93% respectively. The loin eye area (cm^2) differed significantly. The weights of edible and inedible offals were significantly higher (P<0.001) in males in the same age group. The average lean content in different cut-up parts was higher (P<0.05) in G-3(G-$_{Ra}$). The shear force values and other meat quality attributes were significantly different among the groups. It is concluded that age and sex had a significant effect on carcass traits and meat quality in Malpura sheep.

Key words: Age, Carcass traits, Malpura sheep, Meat quality, Sex.

INTRODUCTION

Meat is an awesome wellspring of supplements. Amino acids are significant for the growth and repair of body tissues in humans (Lawrie and Ledward, 2006). Of late, leaner carcasses are in demand due to better consumer awareness for healthy meat, with a focus on the quantity and quality of fat. Meat quality and carcass characteristics vary among animal species (Muela *et al*., 2012). With increasing slaughter body weight in lambs, the carcass becomes darker, fattier and less tender (Abdullah and Qudsieh, 2008). Purchaser favors meat with more lean and less fat and steady quality (Gadekar *et al*., 2015). Recently, global demand for low-fat, high-protein lean meat has increased to satisfy the protein requirements (Shauyenov *et al.,* 2016).

Malpura is one of the main sheep of the semi-arid region of Indian state Rajasthan, notable for its hardiness, versatility to unforgiving climate and considered as an incredible mutton breed (Gowane *et al*., 2015). The carcass traits of intensively fed lambs (3 to 6 months of age) were studied. The dressing yield (%) on live weight and empty body weight basis was 51 and 59, respectively (Prasad *et al.,* 1981). The Malpura lambs maintained on suckling with *ad-lib* creep ration and tree leaves accomplished 14.6 kg body weight at 3 months of age with ADG of 134 g and FCR of 2.00 (Tripathi *et al*., 2011). For improved weight gain and thus higher carcass weight in Malpura lambs, *Lactobacillus acidophilus* culture @ 1.5 ml/kg body weight can be used as a probiotic (Gadekar *et al.,* 2014). Malpura ram lambs with an adequate dressing yield, loin eye area, desirable commercial cuts and lean, fat and bone content when slaughtered at 6 months of age and weighed 24.49 kg body weight (Shinde *et al.,* 2018). The information about carcass attributes of Malpura sheep has been documented in the literature (Karim *et al.,* 2000; 2002; Sen and Karim, 2010; Sureshkumar *et al.,* 2010); However, the detailed studies on the effect of age and sex on meat qualities in Malpura sheep is lacking.

Therefore, the present study was undertaken to study the effect of age and sex on carcass traits and meat quality of Malpura sheep in depth.

[1]Department of Livestock Products Technology, Post Graduate Institute of Veterinary Education and Research, Jaipur-302 001, Rajasthan, India.

[2]ICAR-Central Sheep and Wool Research Institute, Avikanagar, Tonk-304 501, Rajasthan, India.

Corresponding Author: Y.P. Gadekar, ICAR-National Research Centre on Meat, Chengicherla, Hyderabad-500 092, Telangana, India. Email: yogirajlpt@gmail.com

How to cite this article: Meel, S.K., Pandey, A., Gadekar, Y.P., Soni, A., Meena, P., Gurjar, A.S. and Chauhan, V.K. (2024). Age and Sex Affect Carcass Traits and Meat Quality in Malpura Sheep. Indian Journal of Animal Research. 58(2): 330-335. doi: 10.18805/IJAR.B-4864.

Submitted: 14-01-2022 **Accepted:** 02-05-2022 **Online:** 28-05-2022

MATERIALS AND METHODS

The study was undertaken at ICAR-Central Sheep and Wool Research Institute, Avikanagar, Rajasthan during 2018-2019. Twenty-four Malpura sheep were divided into four groups *viz*., ram lamb (G-$_{Rl}$), ewe lamb (G-$_{El}$) (10-12 months each), ram (G-$_{Ra}$) and ewe (G-$_{Ea}$) (20-24 months each). The animals fasted overnight with free access to water were slaughtered by the Halal method. Sticking, legging, dressing

and evisceration were done. The carcass was portioned along the midline and the left half was split into the leg, loin, rack, neck and shoulder, breast and foreshank (ISI, 1963). The *Longissimus dorsi* muscle was collected from each carcass for meat quality evaluation. The meat pH was estimated according to Trout *et al.* (1992). The moisture, protein, fat and ash contents were determined as per AOAC (2000). Total meat pigments were determined by Hornsey (1956). The procedure of Knipe *et al.* (1985) was used for the estimation of salt soluble proteins. The muscle fibre diameter was evaluated according to Jeremiah and Martin (1982). The muscle fibre diameter was measured as the mean diameter of the middle and the two extremities of the 25 randomly selected muscle fibres and expressed in micrometer.

The hydroxyproline (HP) content was determined using a procedure by Neuman and Logan (1950). The collagen content was estimated using a conversion factor of 7.25 (Goll *et al*., 1963). A soluble fraction of hydroxyproline was prepared by slightly modifying the procedure of Okonkwo *et al*. (1992). The percentage of soluble hydroxyproline was calculated as follows:

$$\text{Per cent soluble hydroxyproline} = \frac{\text{Soluble hydroxyproline of supernatant fraction}}{\text{Soluble hydroxyproline of residue fraction + Insoluble hydroxyl proline of supernatant}} \times 100$$

$$\text{Collagen solubility (\%)} = \frac{\text{Soluble collagen}}{\text{Soluble collagen + Insoluble collagen}} \times 100$$

The shear force values (N) were estimated with a texture analyzer (TA.HD plus), Stable Micro System Ltd., England. The myofibrillar fragmentation index (MFI) was estimated as per Davis *et al.* (1980). The water holding capacity was determined according to Wardlaw *et al*. (1973). The method of Swift *et al*. (1961) was used for determining emulsifying capacity.

The data obtained were analyzed using the SPSS v. 16.0. The variations in carcass traits and meat quality parameters between age groups and sex were compared by one-way ANOVA and test of significance.

RESULTS AND DISCUSSION

Body length, height, heart girth, paunch girth, pre-slaughter weight varied significantly (P<0.001) with age and sex of animals (Table 1). It increased with the advancement of age and was higher for males than females for identical age groups. Pre-slaughter weight, empty live weight, hot carcass weight were highest in ram and lowest in ewe lamb. These might be ascribed to the size differences in males and females. Dressing percentage based on pre-slaughter weight, the weight of reproductive organs, weight of caul fat, kidney fat, the weight of total edible, the weight of different type fat and fat percentage were higher in female than male in same age groups. The hot carcass weight was in accordance with Kumar *et al*. (2017). The loin eye area differed significantly (P<0.001) in all the groups. Our study results are in agreement with earlier findings (Das *et al*., 2008).

Yields of edible and inedible offals (Table 1) differed significantly (P<0.001). It increased with the advancement of age and was higher for males than females. The weight of testis and testis fat increased with age. Weights of the spleen, caul fat, kidney fat, kidney, heart, liver, lungs, total edible offal, differed significantly (P<0.05).

The primal cut yield was affected significantly (P>0.05) by age and sex (Table 2). Shinde *et al.* (2018) reported that half carcass weight increased significantly with age in Malpura sheep. Leg cut was the major portion of half carcass followed by neck and shoulder cut (Fig 1). Pena *et al.* (2005) reported that sex affected the amount and type of fat deposited. The lean fat ratio was highest for G-1(G_{Rl}). A higher lean fat ratio is always desirable. The meat bone ratio was highest for G-4(G_{Ea}) while meat bone ratios in G-1 (G_{Rl}) and G-2(G_{El}) were comparable. With an increase in the animal age, the growth rate of muscles is unchanged while the growth rate of bone tissue reduces (Koyuncu *et al.*, 2007). Santos-Silva and Portugal (2001) observed that the meat/bone ratio increases with an increase in slaughter weight.

The meat quality was also affected due to age and sex in Malpura sheep. Markedly higher protein content was observed in the ewe lamb group. A significantly (P<0.001)



Fig 1: Effect of age and sex on commercial cuts in Malpura sheep.

higher fat content was recorded in the female group for the same age group. With the progression of slaughter age, fat and protein levels increased whereas moisture content declined (Madruga *et al.,* 1999). Tejeda *et al.* (2008) also found a non-significant effect of sex on moisture content. A muscle composition varies with increasing animal age irrespective of species, breed, or sex and in most cases; younger animals are considered to have less myoglobin, less intramuscular fat and more moisture (Lawrie, 1998). Fat is the last tissue to mature and older animals tend to be fatter. The inter-muscular fat content of *Longissimus dorsi* increased and moisture content decreased with increasing slaughter weight. Lipids are major components of the muscle structure that give the meat the sensory characteristics

Table 1: Effect of age and sex on pre-slaughter parameters in Malpura sheep (Mean±SE).

Parameters	$G\text{-}_{Rl}$	$G\text{-}_{El}$	$G\text{-}_{Ra}$	$G\text{-}_{Ea}$
Length (cm)	55.67 ± 0.67^{bc}	51.83 ± 1.54^{c}	63.50 ± 1.06^{a}	58.50 ± 2.35^{b}
Height (cm)	61.33 ± 0.88^{c}	56.00 ± 0.73^{d}	69.00 ± 0.93^{a}	64.50 ± 0.96^{b}
Heart girth (cm)	71.67 ± 2.59^{b}	65.50 ± 1.48^{c}	84.00 ± 1.24^{a}	79.17 ± 1.40^{a}
Paunch girth (cm)	86.33 ± 1.38^{b}	73.50 ± 2.69^{c}	98.17 ± 1.66^{a}	87.67 ± 1.38^{b}
Pre Slaughter weight (kg)	23.83 ± 1.07^{c}	17.00 ± 1.21^{d}	35.82 ± 1.24^{a}	28.08 ± 1.26^{b}
Empty live weight (kg)	19.43 ± 1.01^{c}	13.93 ± 1.02^{d}	29.38 ± 1.18^{a}	23.67 ± 0.93^{b}
Hot carcass weight (kg)	10.19 ± 0.63^{c}	7.15 ± 0.54^{d}	15.76 ± 0.60^{a}	12.63 ± 0.51^{b}
Dressing yield (%) PSW	42.61 ± 0.81^{b}	41.99 ± 0.83^{b}	43.99 ± 0.45^{ab}	44.98 ± 0.55^{a}
Dressing yield (%) ELW	52.35±0.71	51.26±0.57	53.70±0.88	52.93±0.62
Weight of forequarter (kg)	5.47 ± 0.36^{c}	3.75 ± 0.27^{d}	8.50 ± 0.31^{a}	6.49 ± 0.23^{b}
Weight hindquarter (kg)	4.59 ± 0.26^{c}	3.28 ± 0.27^{d}	7.01 ± 0.31^{a}	5.65 ± 0.24^{b}
Loin eye area (cm^2)	12.26 ± 0.79^{bc}	9.91 ± 0.66^{c}	17.81 ± 1.61^{a}	13.11 ± 0.36^{b}
Blood (kg)	1.31 ± 0.06^{b}	0.87 ± 0.06^{c}	1.76 ± 0.16^{a}	1.46 ± 0.05^{b}
Head (kg)	1.47 ± 0.04^{c}	1.13 ± 0.05^{d}	2.21 ± 0.08^{a}	1.73 ± 0.03^{b}
Skin (kg)	2.03 ± 0.14^{b}	1.43 ± 0.10^{c}	3.05 ± 0.15^{a}	2.17 ± 0.10^{b}
Fore canon (kg)	0.33 ± 0.01^{b}	0.23 ± 0.01^{c}	0.45 ± 0.02^{a}	0.30 ± 0.01^{b}
Hind canon (kg)	0.28 ± 0.01^{b}	0.20 ± 0.01^{c}	0.38 ± 0.01^{a}	0.25 ± 0.00^{b}
Gallbladder (kg)	0.03±0.00	0.02±0.00	0.04±0.01	0.03±0.01
Total inedible offal(kg)	5.49 ± 0.23^{b}	3.93 ± 0.23^{c}	7.95 ± 0.33^{a}	6.17 ± 0.29^{b}
Inedible Offal (%) on PSW	23.12±0.66	23.2±0.37	22.24±0.73	22.18±0.53
Gastrointestinal tract (filled) (kg)	6.63 ± 0.28^{b}	4.77 ± 0.37^{c}	9.64 ± 0.53^{a}	7.34 ± 0.43^{b}
Gastro-intestinal tract (empty) (kg)	2.22 ± 0.17^{bc}	1.71 ± 0.16^{c}	3.20 ± 0.39^{a}	2.80 ± 0.13^{ab}
Testis fat (kg)	0.03±0.00	NA	0.04±0.00	NA
Testis (kg)	0.16 ± 0.02^{b}	NA	0.35 ± 0.04^{a}	NA
Spleen (kg)	0.05 ± 0.00^{b}	0.04 ± 0.00^{b}	0.10 ± 0.01^{a}	0.06 ± 0.01^{b}
Pancreas (kg)	0.03±0.0	0.02±0.0	0.04±0.0	0.04±0.0
Caul fat (kg)	0.06 ± 0.03^{c}	0.10 ± 0.05^{bc}	0.20 ± 0.03^{b}	0.50 ± 0.06^{a}
Kidney fat (kg)	0.07 ± 0.02^{b}	0.07 ± 0.02^{b}	0.15 ± 0.03^{b}	0.37 ± 0.08^{a}
Kidney (kg)	0.06 ± 0.0^{b}	0.05 ± 0.0^{c}	0.10 ± 0.01^{a}	0.07 ± 0.0^{b}
Heart (kg)	0.10 ± 0.0^{c}	0.08 ± 0.01^{d}	0.15 ± 0.01^{a}	0.12 ± 0.01^{b}
Liver (kg)	0.44 ± 0.02^{b}	0.31 ± 0.01^{c}	0.67 ± 0.05^{a}	0.51 ± 0.01^{b}
Lungs (kg)	0.46 ± 0.02^{c}	0.38 ± 0.03^{d}	0.67 ± 0.02^{a}	0.56 ± 0.03^{b}
Total edible offal (kg)	1.27 ± 0.08^{b}	1.06 ± 0.11^{b}	2.09 ± 0.11^{a}	2.21 ± 0.15^{a}
Total inedible offal (%) on PSW	5.49 ± 0.23^{b}	3.93 ± 0.23^{c}	7.95 ± 0.33^{a}	6.17 ± 0.29^{b}
Chilling loss (%)	2.91 ± 0.28^{a}	2.07 ± 0.25^{b}	3.23 ± 0.16^{a}	2.67 ± 0.13^{ab}
Lean yield (%)	56.98 ± 1.52^{b}	56.24 ± 0.70^{b}	60.24 ± 0.54^{a}	57.68 ± 0.83^{ab}
Subcutaneous fat (%)	3.91 ± 0.31^{b}	4.44 ± 0.65^{b}	4.14 ± 0.46^{b}	6.47 ± 0.58^{a}
Intermuscular fat (%)	3.41±0.70	3.89±0.88	4.01±0.22	5.82±0.78
Separable fat (%)	7.32 ± 0.99^{b}	8.33 ± 1.5^{b}	8.15 ± 0.67^{b}	12.4 ± 1.35^{a}
Dissected bone (%)	31.64 ± 1.20^{a}	31.26 ± 1.26^{a}	28.94 ± 0.57^{ab}	26.48 ± 1.50^{b}
Lean fat ratio	8.45±1.02	7.96±1.46	7.66±0.64	5.29±0.57
Meat bone ratio	1.82 ± 0.11^{b}	1.82 ± 0.08^{b}	2.09 ± 0.05^{ab}	2.16 ± 0.12^{a}

N=6 in each group; Means bearing different superscripts differ significantly (*-P<0.05, **-P<0.01, ***-P<0.001), NS- Non-significant (P>0.05). $G\text{-}_{Rl}$ = Ram lamb, $G\text{-}_{El}$ = Ewe lamb, $G\text{-}_{Ra}$= Ram and $G\text{-}_{Ea}$= Ewe, NA= Not applicable.

Table 2: Effect of age and sex on meat quality in Malpura sheep (Mean±SE).

Parameter	$G\text{-}_{Rl}$	$G\text{-}_{El}$	$G\text{-}_{Ra}$	$G\text{-}_{Ea}$
Moisture (%)	76.80±0.92	75.14±1.30	74.94±1.01	73.11±1.18
Protein (%)	20.83±0.94	21.70±1.18	20.72±0.84	20.54±1.4
Fat (%)	$1.20±0.02^{d}$	$2.02±0.15^{c}$	$3.22±0.34^{b}$	$4.83±0.21^{a}$
Ash (%)	1.17±0.05	1.14±0.05	1.13±0.04	1.52±0.21
pH 45 min.	6.89±0.09	7.03±0.09	6.87±0.09	6.65±0.18
Ultimate pH	5.78±0.14	5.81±0.12	5.95±0.13	5.96±0.15
Cooking loss (%)	$26.87±1.67^{a}$	$26.24±0.65^{ab}$	$22.8±1.67^{b}$	$23.18±0.71^{ab}$
Firmness (N)	$51.31±2.06^{a}$	$41.56±0.52^{b}$	$53.67±1.42^{a}$	$44.2±0.54^{b}$
Work of shear (N*sec.)	$284.42±14.11^{b}$	$183.05±5.18^{d}$	$324.52±4.74^{a}$	$222.06±2.34^{c}$
Heme iron content (µg/gm)	$11.41±1.01^{b}$	$11.72±0.59^{b}$	$14.56±1.15^{ab}$	$17.27±1.25^{a}$
Total meat pigment (ppm)	$129.35±11.46^{b}$	$132.84±6.67^{b}$	$165.08±12.98^{ab}$	$195.77±14.18^{a}$
Salt soluble protein (%)	5.68±0.65	5.90±0.39	6.16±0.37	7.15±0.72
Collagen content (%)	$2.47±0.09^{c}$	$2.39±0.08^{c}$	$2.95±0.04^{b}$	$3.20±0.03^{a}$
Collagen solubility (%)	$24.70±0.49^{b}$	$28.06±0.76^{a}$	$20.84±0.52^{c}$	$22.23±0.62^{c}$
Water holding capacity (ml/100 gm)	52.86±4.88	50.67±3.63	51.47±1.57	47.71±7.85
Muscle fiber diameter (µm)	$30.78±0.33^{b}$	$31.35±1.04^{b}$	$39.78±0.85^{a}$	$38.79±0.78^{a}$
Myofibrillar fragmentation index (%)	76.76±0.89	77.34±0.81	74.67±3.03	75.54±1.14

N=12 in each group (N= 300 for muscle fibre diameter); Means bearing different superscripts differ significantly (*-P<0.05, **-P<0.01, ***-P<0.001), NS- Non-significant (P>0.05) $G\text{-}_{Rl}$ = Ram lamb, $G\text{-}_{El}$ = Ewe lamb, $G\text{-}_{Ra}$= Ram and $G\text{-}_{Ea}$= Ewe.

desired by the consumer such as juiciness, flavour and aroma (Zorzi *et al.*, 2013).

The pH of the meat is very important as it has a definite impact on meat quality and shelf life. The meat pH recorded after 45 minutes of slaughter was found to be comparatively higher in ewe lamb than others. The meat pH variation might be due to variations in animal age, pre-slaughter handling of animals, slaughter condition and handling of meat. Tejeda *et al*. (2008) reported that animal sex did not affect meat pH. The cooking loss differed (P<0.05) significantly between the groups with comparatively higher values in ram lambs. Yarali *et al.* (2014) recorded cooking losses of 26.80, 28.76 and 24.10 percent in *Longissimus thoracis et lumborum*, *Longissimus thoracis* and *Semitendinosus* muscle, respectively. The firmness (N) and work of shear (N*sec.) differed significantly (P<0.001) between the groups with significantly higher in ram. The fibre diameter, hydroxyproline content and toughness of the meat had a positive correlation with the shear force value of meat while it has a negative correlation with the sarcomere length (Biswas *et al.,* 1989). The value obtained in the present study is in concurrence with the reports of Sen *et al.* (2004) in yearling sheep reared under a semi-arid region. The heme iron content (µg/gm) and total meat pigment (ppm) differed significantly (P<0.001) with sex and age. A significantly higher total meat pigment was observed in ewes and increased with increasing age. The greater meat pigment concentration in aged sheep groups was attributed to the more heme pigment and myoglobin pigment (Mamino and Horn, 1996). As animal age increases the myoglobin concentration increases which in turn improves red color intensity (Warner *et al*., 2007). Advancement in sheep age leads to increased aerobic muscle fibres which raise myoglobin pigment leading to better red colour of meat. Nevertheless, the degree differs according to the breed and nutrition (Warner *et al*., 2007). The salt soluble protein content was comparable (P>0.05) between the groups. Ahmed *et al*. (2015) observed a non-significant effect of age on salt soluble protein content in goatmeat. The collagen content of meat in ewes was significantly higher compared to in other groups. Significantly higher collagen solubility was observed in ewe lambs ($G\text{-}_{El}$). Collagen content and muscle fiber diameter increased while collagen solubility decreased with age. Intra-muscular collagen and cross-link formation depends upon the age of the animal and is directly proportional to age. Polidori *et al*. (2017) observed a significant effect of age on collagen content in lambs. The collagen has a low rate of metabolic turnover which leads to the permanent formation of cross-linkages (Purslow *et al*., 2012). The water holding capacity in ram lambs was comparatively higher than in other groups. Stankov *et al*. (2002) observed a non-significant effect of age on water holding capacity in the young goat of Bulgarian breeds and crossbreeds of goats. A significantly larger muscle fibre diameter was observed in the ewes group. The myofibrillar fragmentation index was comparatively higher for ewe lambs. The myofibrillar fragmentation index is a measure of myofibrillar protein degradation (Seideman *et al*., 1987). This was highly related to shear force and sensory tenderness ratings (Calkins and Davis, 1980).

CONCLUSION

The findings of the study indicated that pre-slaughter weight, loin eye area, a yield of different edible offals and lean yield in primal cuts were higher in ewes and rams compared to

the ewe lambs and ram lambs, while the above parameters were higher in rams and ram lambs compared to their female counterparts in same age groups. Fat content and total meat pigment were found higher in females compared to males in the same age groups. Collagen content and muscle fiber diameter were higher in the adult animal while collagen solubility was higher in the young animal. The age and sex of animal affected carcass traits and meat quality.

ACKNOWLEDGEMENT

The first author thanks to the Director, ICAR-CSWRI, Avikanagar and Incharge LPT section for providing facilities to carry out work. Author is also grateful to RAJUVAS, Bikaner for providing financial support in form of stipend.

Conflict of interest

All authors declared that there is no conflict of interest.

REFERENCES

Abdullah, A.Y., Qudsieh, R.I. (2008). Carcass characteristics of Awassi ram lambs slaughtered at different weights. Livestock Science. 11: 165-175.

Ahmed, M.H., Elghandour, M.M.Y., Salem, A.Z.M., Zeweil, H.S., Kholif, A.E., Klieve, A.V., Abdelrassol, A.M.A. (2015). Influence of *Trichoderma reesei* or *Saccharomyces cerevisiae* on performance, ruminal fermentation, carcass characteristics and blood biochemistry of lambs fed *Atriplex nummularia* and *Acacia saligna* mixture. Livestock Science. 180: 90-97.

AOAC (Association of Official Analytical Chemists) (2000). Official Methods of Analysis of AOAC International, 17th ed. AOAC International, Arlington, VA, USA.

Biswas, S., Radhakrishnan, K.T., Arumugam, M.P., Ramamurthi, R. (1989). Studies on the relationship between tenderness measures on myofibrillar and connective tissue components of heated and unheated mutton. Cheiron. 18: 154-157.

Calkins, C.R., Davis, G.W. (1980). Fragmentation index of raw muscle as a tenderness predictor of steaks from US Good and US Standard steer and bullock carcasses. Journal of Animal Science. 50: 1067-1072.

Das, A.K., Dass, G., Singh, N.P. (2008). Growth, carcass characteristics and meat quality of Muzaffarnagari lambs at various stages under intensive and semi-intensive management Ind. Journal of Animal Science. 78: 541-546.

Davis, G.W., Dutson, T.R., Smith, G.C., Carpenter, Z.L. (1980). Fragmentation procedure for bovine longissimus muscle as an index of cooked steak tenderness. Journal of Food Science. 45: 880-884.

Gadekar, Y.P., Shinde, A.K., Sahoo, A., Karim, S.A. (2015). Effect of probiotic supplementation on carcass traits and meat quality of Malpura Lambs. Indian Journal of Small Ruminants. 21: 306-310.

Gadekar, Y.P., Shinde, A.K., Soren, N.M., Karim, S.A. (2014). Effect of different levels of *Lactobacillus acidophilus* culture on carcass traits and meat quality of Malpura lambs. Ruminant Science. 3(2): 229-234.

Goll, D.E., Bray, R.W., Hoekstra, W.G. (1963). Age associated changes in muscle composition. The isolation and properties of a collagenous residue from bovine muscle. Journal of Food Science. 28: 503-509.

Gowane, G.R., Prince, L.L.L., Lopes, F.B., Paswan, C., Sharma, R.C. (2015). Genetic and phenotypic parameter estimates of live weight and daily gain traits in Malpura sheep using Bayesian approach. Small Ruminant Research. 128: 10-18.

Hornsey, H.C. (1956). The colour of cooked pork. I. Estimation of the nitric oxide haeme pigments. Journal of the Science of Food and Agriculture. 7: 534-540.

ISI. (1963). Indian Standard Specification for Mutton and Goat Flesh: Fresh Chilled and Frozen, IS-2536. Indian Standard Institution, New Delhi.

Jeremiah, L.E., Martin, A.H. (1982). Effect of prerigor chilling and freezing and subcutaneous fat cover upon the histological and shear properties of bovine *Longissimus dorsi* muscle. Journal of Animal Science. 62: 353-361.

Karim, S.A., Santra, A. and Verma, D.L. (2002). Growth, feed conversion efficiency and carcass characteristics of Malpura and Malpura × Awassi crossbred lambs in hot semi-arid environment. Asian-Australasian Journal of Animal Sciences. 13(3): 377-381.

Karim, S.A., Santra, A. (2000). Growth performance of Malpura and crossbred lambs under intensive feeding. Small Ruminant Research. 37: 287-291.

Knipe, C.L., Olson, D.G., Rust, R.E. (1985). Effects of selected inorganic phosphates, phosphate levels and reduced chloride levels on protein solubility, stability and pH of meat emulsions. Journal of Food Science. 50: 1010.

Koyuncu, M., Duru, S., Uzun, Ş.K., Özişş, Ş., Tuncel, E. (2007). Effect of castration on growth and carcass traits in hair goat kids under a semi-intensive system in the south-Marmara region of Turkey. Small Ruminant Research. 72: 38-44.

Kumar, D.A.P., Prakash, M.G., Gupta, B.R., Raghunandan, T., Chandra, A.S. (2017). Carcass traits in Deccani sheep. The Pharma Innovation. 6: 198-201.

Lawrie, R.A. (1998) Lawrie's Meat Science (6th ed.). Cambridge England: Woodhead Publishing Limited.

Lawrie, R.A., Ledward, D.A. (2006). Lawrie's Meat Science, Abington. http://dx.doi.org/10.1533/9781845691615.

Madruga, M.S., Arruda, S.G.B., Nascimento, J.A. (1999). Castration and slaughter age effects on nutritive value of the "mestiço" goat meat. Meat Science. 52: 119-125.

Mamino, E.N., Horn, S.P. (1996). Chemical, Processing and Sensory Characteristics of Carabeef Ham from Phil-Murrah × Phil-Murrah crossbred. In 2. Asian Buffalo Association Congress, Makati City (Philippines), 9-12 Oct 1996. PSAS Foundation,Inc.

Muela, E., Sañudo, C., Campo, M.M., Medel, I., Beltrán, J.A. (2012). Effect of freezing method and frozen storage duration on lamb sensory quality. Meat Science. 90: 209-215.

Neuman, R.E., Logan, M.A. (1950). The determination of hydroxyproline. Journal of Biological Chemistry. 184: 299-306.

Nowak, D. (2011). Enzymes in tenderization of meat-The system of calpains and other systems-A review. Polish Journal of Food and Nutrition Sciences. 61: 231-237.

Okonkwo, T.M., Obanu, Z.A., Ledward, D.A. (1992). Characteristics of some intermediate moisture smoked meats. Meat Science. 31: 135-145.

Peña, F., Cano, T., Domenech, V., Alcalde, M.J., Martos, J., Garcia-Martinez, A., Rodero, E. (2005). Influence of sex, slaughter weight and carcass weight on "non-carcass" and carcass quality in Segureña lambs. Small Ruminant Research. 60: 247-254.

Polidori, P., Pucciarelli, S., Cammertoni, N., Polzonetti, V., Vincenzetti, S. (2017). The effects of slaughter age on carcass and meat quality of Fabrianese lambs. Small Ruminant Research. 155: 12-15.

Prasad, V.S.S., Arora, C.L., Singh, R.N., Bohra, S.D.J. (1981). Assessment of carcass and mutton quality in native and cross breed sheep. Indian Journal of Animal Sciences. 51: 638-642.

Purslow, P.P., Archile-Contreras, A.C., Cha, M.C. (2012). Meat Science and Muscle Biology Symposium: Manipulating meat tenderness by increasing the turnover of intramuscular connective tissue. Journal of Animal Sciences. 90: 950-1959.

Santos-Silva, J., Portugal, A.V. (2001). The effect of weight on carcass and meat quality of Serra da Estrela and Merino Branco lambs fattened with dehydrated lucerne. Animal Research. 50: 289-1298.

Seideman, S.C., Koohmaraie, M., Crouse, J.D. (1987). Factors associated with tenderness in young beef. Meat Science. 20: 281-291.

Sen A., Karim, S. (2010). Carcass and meat quality characteristics of designated indigenous sheep breeds of India. Indian Journal of Animal Sciences. 80(4): 362-365.

Sen, A.R., Santra, A. and Karim, S.A. (2004). Carcass yield, composition and meat quality attributes of sheep and goat under semiarid conditions. Meat Science. 66: 757-763.

Shauyenov, S., Islamov, Y., Narbayev, S., Ibrayev, D. (2016). Effect of sire breed on the productivity of Kazakh mutton-semifine-wool sheep. Indian Journal of Animal Research. 50: 418-424. doi: 10.18805/ijar.9493.

Shinde, A.K., Gadekar, Y.P., Gauri, J. (2018). Effect of slaughter age on carcass traits of Malpura lambs. Indian Journal of Small Ruminants. 24: 308-314.

Stankov, I.K., Todorov, N.A., Mitev, J.E., Miteva, T. (2002). Study on some qualitative features of meat from young goat of Bulgarian breeds and crossbreeds of goats slaughtered at various ages. Asian-Australasian Journal of Animal Sciences. 15: 283-289.

Sureshkumar, S., Chopra A., Gowane, G.R. (2010). Carcass characteristics of Malpura lambs maintained under feedlot condition. Indian Journal of Small Ruminants. 16(1): 280-283.

Swift, C.E., Fryar, A.J., Lockett, C. (1961). Comminuted meat emulsions-Capacity of meats for emulsifying fat. Food Technology. 15: 468.

Tejeda, J.F., Peña, R.E. Andrés, A.I. (2008). Effect of live weight and sex on physico-chemical and sensorial characteristics of Merino lamb meat. Meat Science. 80: 1061-1067.

Tripathi, M.K, Tripathi, P., Karim, S.A. (2011). Nutrition and feeding of lambs for mutton production under tropical climates: A review. Indian Journal of Small Ruminants. 17: 135-150.

Trout, E.S., Hunt, M.C., Johnson, D.E., Clauss, J.R., Kastner, C.L., Kropf, D.H., Stroda, S. (1992). Chemical, physical and sensory characterization of ground beef containing 5 to 10 per cent fat. Journal of Food Science. 57: 25-29.

Wardlaw, F.B., Maccaskill, L.H., Acton, J.C. (1973). Effect of post mortem muscle changes in poultry meat loaf properties. Journal of Food Science. 38: 421-424.

Warner, R.D., Pethick, D.W., Greenwood, P.L., Ponnampalam, E.N., Banks, R.G., Hopkins, D.L. (2007). Unravelling the complex interactions between genetics, animal age and nutrition as they impact on tissue deposition, muscle characteristics and quality of Australian sheep meat. Australian Journal of Experimental Agriculture. 47: 1229-1238.

Yarali, E., Yilmaz, O., Cemal, I., Karaca, O., Taşkin, T. (2014). Meat quality characteristics in Kıvırcık lambs. Turkish Journal of Veterinary and Animal Sciences. 38: 452-458.

Zorzi, K., Bonilha, S.F.M., Queiroz, A.C., Branco, R.H., Sobrinho, T.L., Duarte, M.S. (2013). Meat quality of young Nellore bulls with low and high residual feed intake. Meat Science. 93: 593-599.