REVIEW ARTICLE Indian Journal of Animal Research, Volume 60 Issue 1: 01-13 (January 2026)

Phytotherapeutic Control of Coccidiosis in Livestock and Poultry Animals: Harnessing Plant Heritage for Novel Drug Discovery- Ethnobotanical Insights and Modern Validation: A Review

Vijay Kumar Dubey[1], Deen Dayal[2], Maya Datt Joshi[2], Alok Bhardwaj[2], Jyotika Vats[3], Ravindra Kumar[1], S.Y. Mukartal[4], Gaya Prasad Jatav[5], Shailendra Thapliyal[6], Kundan Kumar Chaubey[7]

10.18805/IJAR.B-5682

ABSTRACT

Coccidiosis, caused by *Eimeria* species, is a major parasitic disease affecting livestock and poultry, leading to significant economic losses. Conventional treatments are increasingly limited due to drug resistance and concerns over chemical in the livestock products. This review aims to identify and evaluate novel medicinal plants with established ethnobotanical relevance as potential alternatives for the treatment of coccidiosis. Drawing from traditional medicinal practices, plant species historically used for various are explored and assessed their phytochemical constituents, efficacy and safety profiles based on modern scientific evidence. Additionally, recent advances in drug delivery and formulation to improve bioavailability and therapeutic outcomes of plant-based are examined in the present review. The present study also addresses key challenges including standardization, formulation, validation method, mode of action and integration into current veterinary practices. The objective of this review is to provide a comprehensive, evidence-based assessment of plant-derived therapeutics for sustainable and effective coccidiosis management, bridging traditional knowledge with modern veterinary science.

Key words: Antidiarrheal, Anti-inflammatory, Anti-ulcer, Coccidia, Immunomodulatory.

Coccidiosis, caused by *Eimeria* protozoa, is a major parasitic disease impacting livestock and poultry worldwide (Saeed and Alkheraije, 2023). Infection occurs when animals ingest sporulated oocysts from contaminated feed or water, with each *Eimeria* species targeting specific intestinal epithelial cells (Haug *et al.,* 2008). After ingestion, sporozoites are released in the duodenum and invade the intestinal epithelium, starting a life cycle with asexual (merogony) and sexual (gametogony) phases (Conway and McKenzie, 2007). Asexual reproduction produces merozoites that spread the infection, while sexual reproduction forms gametes that fuse into zygotes and develop into oocysts. This rapid cycle allows a single oocyst to produce millions of parasites within 4 to 5 days (Pattison *et al.,* 2007). Pathogenic *Eimeria* infections cause hemorrhagic enteritis, bloody diarrhea, malabsorption, emaciation and stunted growth (Awais *et al.,* 2012; Ram *et al.*, 2019; Kaur *et al.*, 2019). These pathological changes reduce feed efficiency, delay sexual maturity and lower production, leading to economic losses through both mortality and subclinical effects. Disease severity depends on host, parasite and environmental factors (Reid, 1990). Its prevalence varies globally, reaching up to 98% in goats and 86% in calves and sheep in England and the USA (Keeton and Navarre, 2018), nearly 100% in Malaysian goats (Paul *et al.,* 2020), 78 to 79% in Ethiopian and Iranian goats and sheep (Dau *et al.,* 2021; Barre *et al.,* 2023; Hatam-Nahavandi *et al.,* 2023), 28.6% in India (Bangoura and Bardsley, 2020) and 39.8% in Amazonian livestock (Paul *et al.,* 2020). Management traditionally relies on synthetic anticoccidials, vaccination and plant-derived

[1]Division of Animal Nutrition Management and Products Technology, ICAR-Central Institute for Research on Goats, Makhdoom, Mathura-281 122, Uttar Pradesh, India.
[2]Department of Biotechnology, GLA University, Mathura-281 406, Uttar Pradesh, India.
[3]Department of Pathology, Santosh Institute of Allied Health Sciences, Santosh Nagar-201009, Uttar Pradesh, India.
[4]Department of Veterinary Microbiology, Veterinary College, Karnataka Veterinary, Animal and Fisheries Sciences University , Athani-591 230 Belagavi, Karnataka, India.
[5]Department of Veterinary Pathology, College of Veterinary Science and Animal Husbandry, Nanaji Deshmukh Veterinary Science University, Mhow-453 446, Madhya Pradesh, India.
[6]Uttaranchal Institute of Technology, Uttaranchal University, Dehradun-248 007, Uttarakhand, India.
[7]Department of Biotechnology, School of Basic and Applied Sciences, Sanskriti University, Mathura-281401, Uttar Pradesh, India.

Corresponding Author: Kundan Kumar Chaubey, Department of Biotechnology, School of Basic and Applied Sciences, Sanskriti University, Mathura-281401, Uttar Pradesh, India.
Email: kundan2006chaubey@gmail.com
ORCIDs: 0000-0001-7069-080X, 0000-0002-7275-3189, 0000-0002-6028-475X, 0000-0002-7383-188X, 0009-0007-9890-9149, 0000-0002-4954-4514, 0009-0005-6168-0489, 0000-0003-0555-6617, 0009-0002-6212-2057, 0000-0002-5940-4638

How to cite this article: Dubey, V.K., Dayal, D., Joshi, M.D., Bhardwaj, A., Vats, J., Kumar, R., Mukartal, S.Y., Jatav, G.P., Thapliyal, S. and Chaubey, K.K. (2026). Phytotherapeutic Control of Coccidiosis in Livestock and Poultry Animals: Harnessing Plant Heritage for Novel Drug Discovery- Ethnobotanical Insights and Modern Validation: A Review. *Indian Journal of Animal Research.* **60(1):** 01-13. doi: 10.18805/IJAR.B-5682.

Submitted: 13-09-2025 **Accepted:** 02-12-2025 **Online:** 24-12-2025

products. While coccidiocides and ionophores are cost-effective and widely used, rising drug resistance and regulatory restrictions, such as the Europian Union's ban on prophylactic additives (Council Directive 2011/50/EU), have prompted alternative approaches (Chapman *et al.,* 2010). Vaccination with wild-type or attenuated *Eimeria* strains offers partial protection and plant-based natural products are gaining attention as sustainable feed additives, though challenges remain in standardization, safety, mechanism elucidation and cost-effectiveness (Gokila *et al.,* 2014; Yang *et al.,* 2015; Murshed *et al.*, 2023).

The rising global prevalence of coccidiosis and increased interest in natural therapies have driven research into plant-derived anticoccidial agents. Phytochemicals show promise as supplements to conventional treatments by providing immunomodulatory, anti-inflammatory and direct antimicrobial effects. The present review explores herb-based alternatives targeting coccidial infections, emphasizing anti-inflammatory, antidiarrheal, anti-ulcer and immunomodulatory actions. Evaluating traditional ethnobotanical knowledge offers valuable insights for developing plant-based strategies, potentially reducing dependence on synthetic anticoccidial drugs.

Source of literature and selection process

Coccidia infections cause diarrhea, ulcers, inflammation and immunosuppression in animals. The present study reviews plants with immunomodulatory, anti-inflammatory, anti-ulcer and anti-diarrheal properties, focusing on those cited in traditional Indian literature and validated by *in vitro* and *in vivo* studies. Although not yet tested directly against coccidiosis, these plants show potential as future phytotherapeutic agents. The present review integrates traditional Ayurvedic knowledge with modern research, considering only relevant English-language studies and excluding repetitive or unrelated data.

Sources of evidence

This review draws on both traditional and modern scientific sources. Key references include *The Ayurvedic Pharmacopoeia of India*, which details validated formulations and therapeutic uses, alongside contemporary studies sourced from Google Scholar, PubMed and Scopus. By combining traditional knowledge with evidence-based research, 48 medicinal plants were identified as promising candidates for potential treatment of coccidial infections.

Screening process and evidence mapping

After screening and removing duplicates, 48 medicinal plants from Indian traditional texts previously untested for anticoccidial activity were identified for their anti-inflammatory, antidiarrheal, anti-ulcer and immunomodulatory properties (Table 1). Their scientific relevance was verified through databases like Google Scholar, PubMed and Scopus, including only English-language articles with supportive evidence. Plants without data on these properties were excluded. Fig 1 shows the selection and screening workflow, while targeted keywords ensured retrieval of relevant literature for each activity.

Development of coccidial infection

Coccidiosis starts when animals ingest sporulated oocysts containing sporocysts and sporozoites. In the gut, the oocyst wall breaks, releasing sporozoites that invade intestinal cells. These sporozoites multiply asexually, producing many merozoites and repeat this cycle several times. Later, sexual stages form male and female gametes, which combine to create new oocysts that are passed in the feces (Saif, 2009). The life cycle of coccidia with environmental transmission and host-dependent developmental stages is shown in Fig 2.

Conceptual Framework for anti-coccidial activity of selected plants

Based on current literature, the intestinal epithelium and gut-associated immune components play a pivotal role in host-pathogen interactions during *Eimeria* infection. The major sites and mechanisms involved in the pathogenesis and potential intervention points are summarized and the mechanistic flowchart of plants exhibiting anti-coccidial effects is shown in Fig 3.

Invasion of intestinal epithelium and nutrient absorption

The intestinal epithelium, responsible for nutrient absorption, is also the site where *Eimeria* parasites invade and reproduce (Shivaramaiah *et al.,* 2014). The invasive sporozoites enter gut cells, multiply asexually (merogony) producing around 1,000 merozoites per sporozoite, repeating 2 to 4 cycles depending on the species. These merozoites then enter the sexual phase (gametogony), forming gametes that fuse into zygotes and develop into unsporulated oocysts, which are shed in feces (Lillehoj, 1998). This cycle damages the gut lining, weakens the host and aids parasite transmission.

Mucosal and cellular immune mechanisms

After *Eimeria* infection, the intestinal epithelium not only absorbs nutrients but also activates immune defenses (Shivaramaiah *et al.,* 2014). Gut-associated lymphoid tissue, including Peyer's patches, cecal tonsils and lamina propria lymphocytes, processes antigens, produces antibodies and stimulates B cells, T cells, Natural killer (NK) cells and dendritic cells (Lillehoj, 1998; Dalloul and Lillehoj, 2006). Additional defenses include mucus, antimicrobial peptides and a healthy gut microbiota (Zhao *et al.,* 2001). Cell-mediated immunity, especially $CD4^+$ and $CD8^+$ T cells producing Interferon-gamma (IFN-γ), is crucial for parasite elimination (Kim *et al.,* 2019; Arstila *et al.,* 1994; Lillehoj and Choi, 1998). Intraepithelial lymphocytes and NK cells provide early local defense, while dendritic cells bridge innate and adaptive immunity, activating T cells, reducing oocyst shedding and enhancing protection (Lillehoj, 1989; Cornelissen *et al.,* 2009; Schnitzer *et al.,* 2010;

Table 1: Potential anti-coccidial plants, major phytoconstituents and associated mechanisms.

Botanical name	Ayurvedic name	Utilized part	Key phyto-chemicals	Formulation method	Mode of action	Observed effects	Validation model	Reference
Coleus forskohlii	*Gandhira*	Root, Stem	Coleonol, coleosol, naphthopyrone and coleoforsin	Aquous, alcoholic, methanolic, ethanolic and chloroform extract	Protects cell membranes, prevents protein damage and inhibits COX-2 enzyme activity.	Anti-inflammatory	*In vivo*	(Menon and Latha, 2011; Kundur *et al*., 2012; Philomina mary and Malarvizhi, 2022)
Erythrina indica Lam.	*Paribhadra*	Stem, bark	Apigenin and beta-sitosterol	Aquous and hydroalcoholic extract	Blocks neutrophil and monocyte recruitment and prevents protein denaturation.	Anti-inflammatory	*In vivo*	(Shah *et al.,* 2017; Wankhede *et al.,* 2009)
Nerium indicum Mill.	*Karavira*	Leaf, flower	Oleandrin	Alcoholic and hydroalcoholic extract	Inhibits nitric oxide production and PEG2 synthesis.	Anti-inflammatory	*In vivo*	(Dey and Chaudhuri, 2015; and Balkan *et al.,* 2018)
Swertia chirata Buch. Ham	*Kiratatikta*	Whole plant	Xanthones, xanthone and glycoside	Ethanolic, alcoholic and hydroalcoholic extract	Inhibits tissue swelling and downregulates COX-2, NF-κB and MAPK pathways, leading to diminished serum proinflammatory responses	Anti-inflammatory	*In vivo*	(Das *et al.,* 2012; Hu *et al.,* 2019)
Cannabis sativa linn.	*Vijaya*	Leaf, stem	Cannabinols and tetrahydrocannabinol	Ointment, aquous and ethanolic extract	Suppresses NO and pro-inflammatory cytokines, blocks NF-κB/MAPK, reduces ROS and TNF-α, restores apoptosis balance, relieves edema, stabilizes mast cells, lowers oxidative stress, modulates ECS and protects epithelium.	Anti-inflammatory	*In vivo*	(Kim *et al.,* 2022; Woliñska *et al.,* 2025; Ambrose and Simmons, 2019; Cocetta *et al.,* 2021)
Angelica archangelica Linn.	Chanda	Root	Angelica lactone and linalool	Essential oil and aquous extract	Reduces IL-6, IL-4 and IL-13 levels and prevents protein denaturation.	Anti-inflammatory	*In vitro*	(Fraternale *et al.,* 2018)
Cedrus deodara	*Devadaru*	Stem bark,Heart wood	Thujopsene, α-cedrene and cedrol	Aquous extract and essential oil	Shows anti-inflammatory action by blocking COX-2, TNF-α and NF-κB, reducing swelling, stabilizing mast cells and inhibiting LOX.	Anti-inflammatory	*In vivo*	(Chen *et al.,* 2020; Rathor and Goyal, 1973; Shinde *et al.,* 1999a; Shinde *et al.,* 1999b)
Euphorbia neriifolia Linn.	*Snuhi*	Stem, leaves	Nerrifolin, Neriifoliene and euphol	Ethanolic and hydroalcoholic extract	Inhibits LPS-induced IL-6, TNF-α, IFN-γ, NO, COX and LOX production.	Anti-inflammatory	*In vitro*	(Chang *et al.,* 2023; Palit *et al.,* 2016; Palit *et al.,* 2018)
Plumbago zeylanica	*Citraka*	Root	Plumbagin	Hydroalcoholic and	Stabilizes red blood cell	Anti-	*In vitro*	(Thanigavelan *et al.,* 2014;

Table 1: Continue..

Table 1: Continue..

linn.		bark.		dichloromethane extract	membranes and reduces paw edema by inhibiting prostaglandins.	inflammatory	and *In vivo*	Subramaniyan and Paramasivam, 2017)
Ficus lacor Buch.	*Plaksa*	Root	Beta-sitosterol and arabinose	Ethanol, petroleum ether, ethyl acetate and chloroform extract	Reduces inflammation	Anti-inflammatory	*In vivo*	(Sindhu and Arora, 2014)
Trianthema portulacastrum Linn.	*Varsabhu*	Whole plant	β-sitosterol, stigmasterol and their β-glucopyranosides, β-cyanin	Ethanolic extract	Inhibits inflammation by blocking COX and LOX, reducing prostaglandin and leukotriene production.	Anti-inflammatory	*In vivo*	(Kendri and Wari, 2015)
Saraca asoca (Roxb.) De. Willd	*Asoka*	Seeds	Apigenin, gallic acid, linolenic acid and tannins	Aqueous, methanolic and acetone extract	Reduces granuloma tissue weight.	Anti-inflammatory	*In vivo*	(Gupta *et al.,* 2013)
Achyranthes aspera Linn.	*Apamarga*	Leaves	D-Glucuronic acid	Ethanolic and alcoholic extract	Reduces edema by inhibiting inflammatory mediators in both early and late inflammation phases.	Anti-inflammatory	*In vivo*	(Semuel and Shabaraya, 2011)
Anacyclus pyrethrum DC.	*Akarakarabha*	*R*oots, seeds, leaves and capitula	Sarcosine, malonic acid and palmitic acid	Hydroalcoholic extract	Reduces release of serotonin, prostaglandins, bradykinin and histamine.	Anti-inflammatory	*In vivo*	(Jawhari *et al.,* 2020)
Aegle marmelos Corr.	*Bilva*	Leaves	Tannins, skimmianin and caryophyllene	Aquous, methanol acetone and chloroform extract	Reduces inflammation	Anti-inflammatory	*In vivo*	(Arul *et al.,* 2005)
Leucas cephalotes Spreng.	*Dronapuspi*	Whole plant	β-sitosterol, stigmasterol and lupeol	Methanolic, hexanes, dichloromethane, ethyl acetate methanol extract.	Reduces LPS-induced TNF-α and IL-1β production.	Anti-inflammatory	*In vivo*	(Baburao *et al.,* 2010; Patel *et al.,* 2015)
Asteracantha longifolia Nees	*Kokilaksa*	Whole plant	Myristic acid, palmitic acid and linoleic acid	Alcoholic, aquous, chloroform, and petroleum ether extract	Reduces edema	Anti-inflammatory	*In vivo*	(Muthulingam and Veerakumar, 2021)
Clitoria ternatea	*Aparajita*	Root	Rutin, quercetin	Ethanolic extract	Lowers inflammatory	Anti-	*In vivo*	(Swathi *et al.,* 2020)

Table 1: Continue..

Table 1: Continue..

			and malvidin.		mediators.	inflammatory		
Withania somnifera Dunal	*Asvagandha*	Root	Alkaloids and withanolides	Hexane, chloroform, ethyl acetate and methanolic extract	Reduces inflammation	Anti-inflammatory	*In vivo*	(Uthirapathy, 2021)
Boerhaavia diffusa Linn	*Punarnava*	Whole plant	Punarnavine and rotenoids	Hexane, benzene, chloroform, acetone, ethanol, methanol and aquous extract.	Inhibits secretory phospholipase A2.	Anti-inflammatory	*In vitro* and *In vivo*	(Giresha *et al.,* 2017)
Symplocos racemosa Roxb	*Lodhra*	Bark	Phenols, flavonoids, saponins, tannins, glycosides and steroids	Ethylacetate, chloroform, n-butanol and aqueous extract.	Protects membranes and stabilizes cells	Anti-inflammatory	*In vitro* and *In vivo*	(Maitra *et al.,*2017; Ahmad *et al.,* 2014)
Cleome gynandra Linn.	*Ajagandha*	Leaf	Linolenic and arachidic acid	Methanolic extract	Reduces wet feces and total feces volume.	Antidiarrheal	*In vivo*	(Reddy *et al.,* 2015)
Momordica charantia Linn	*Karavallaka*	Leaf	Glycosides and momoridicine	Aquous extract	Reduces diarrheal stool count, slows gut movement and prevents intestinal fluid buildup.	Antidiarrheal	*In vivo*	(BR *et al.,* 2011)
Adhatoda zeylanica Medic	*Vasa*	Leaf	Vasicinol and vasicine	Leaf juice	Increases reabsorption of water.	Antidiarrheal	*In vivo*	(Sutare and Kareppa, 2009)
Pterocarpus marsupium	*Asana*	Stem bark, heartwood	Alkaloids, resin, gum kino and tannins	Ethanolic extract	Increases electrolyte and water absorption, preventing intestinal fluid buildup.	Antidiarrheal	*In vivo*	(Shridhar *et al.,* 2011)
Alstonia scholaris	*Saptaparna*	Aerial part, leaf	Alstoscholarisine K and ditamine	Methanolic extract	Inhibits high K-induced contractions with a spasmolytic effect similar to calcium channel blockers.	Antidiarrheal	*In vivo*	(Shah *et al.,* 2010)
Pongamia pinnata Merr	*Karanja*	Seed	Pongachromene, pinnatin and gamatin	Petroleum ether, ethanolic and aquous extract	Acts against microbial enterotoxins that cause diarrhea.	Antidiarrheal	*In vivo*	(Dahikar and Bhutada, 2017)
Butea	*Palasa*	Stem,	Coumarins, butrin	Ethanolic extract	Inhibits castor oil-induced	Antidiarrheal	*In vivo*	(Gunakkunru *et al.,* 2005)

Table 1: Continue..

Table 1: Continue..

monosperma Kuntze		Bark	and isocoreopsin		diarrhea, PGE2-induced enteropooling and reduces gut motility.			
Cuminum cyminum Linn.	*Svetajiraka*	Seed	Eugenol and limonene	Aquous extract	Inhibits chloride channels, relaxes muscles and enhances water absorption.	Antidiarrheal	*In vivo*	(Sahoo *et al.*, 2014)
Cyperus rotundus	*Nagarmotha*	Whole plant	Dihydroatisine and heteratisine	Methanolic extract	Reduces diarrheal frequency and delays its onset.	Antidiarrheal	*In vivo*	(Uddin *et al.*, 2006)
Bauhinia variegata Blume	*Kanchanara*	Root, seed, stem, bark, leaf	Quercetin and apigenin	Alcoholic extract	Reduces gastric secretion, total acidity and free acidity.	Antiulcer	*In vivo*	(Rajkapoor *et al.*, 2003)
Indigofera tinctoria	*Nili*	Leaf	Indican	Ethanolic extract	Protects gastric mucosa from oxidative damage.	Antiulcer	*In vivo*	(Manimekalai *et al.*, 2018)
Sphaeranthus indicus Linn.	*Munditika*	Whole plant	Sterols	Ethyl acetate, alcoholic and chloroform extract	Reduces total acidity through antisecretory action.	Antiulcer	*In vivo*	(Malairajan *et al.*, 2013)
Panax ginseng Wall	*Ninjin*	Fruit, root	Panaxdiol, panaxtriole and oleanolic acid	Liquid ginseng extract	Enhances immunity by boosting neutrophil migration, phagocytosis and T-cell proliferation.	Immunomodulatory	*In vivo*	(Khare, 2008; *Panax ginseng*, 2009)
Murraya koenigii	*Surabhini-nimba, Nagalinga*	Leaf, fruit, flower	Coumarin, carbazole, alkaloids and flavonoids	Methanolic extract	Boosts phagocytic activity.	Immunomodulatory	*In vivo*	(Shah *et al.*, 2007)
Lagenaria siceraria Mol.	*Katu-tumbi*	Leaf, fruit	Cucurbitacin	Ethanolic extract	Enhances delayed-type hypersensitivity response.	Immunomodulatory	*In vivo*	(Deshpande *et al.*, 2008)
Terminalia arjuna Roxb.	*Arjuna*	Leaf, bark	Flavonoids and oligomeric proanthocyanidins	Aquous and Alcoholic extract	Boosts secondary immune response.	Immunomodulatory	*In vivo*	(Halder *et al.*, 2009)
Bauhinia variegata Linn.	*Kanchanara*	Root, bark, buds	Flavonoids, beta-sitosterol and lupeol	Ethanolic extract	Stimulates macrophages, supports neutrophil adhesion and increases antibody levels.	Immunomodulatory	*In vivo*	(Ghaisas *et al.*,2009)
Urena	*Naagabala*	Root,	Flavanoids	Methanolic extract	Enhances phagocytic	Immunomodul	*In vivo*	(Rinku *et al.*, 2009)

Table 1: Continue..

Table 1: Continue..

lobata linn		flower			function.	atory		
Gymnema sylvestre R.Br	*Gurmaar*	Leaf	Sapogenins	Aquous extract	Stimulates phagocytosis, neutrophil movement and chemotaxis.	Immunomodulatory	*In vivo*	(Malik *et al.,* 2009)
Cordia superba Cham. and C. rufescens	*Shleshmaataka*	Leaf, fruit, bark	Alpha-amyrin	Methanolic extract	Activates nitric oxide production, IFN-γ secretion and LPS response.	Immunomodulatory	*In vivo*	(Costa *et al.,* 2008)
Picrorhiza scrophulariiflora Benth.	*Kutki*	Root	Iridoid and glycosides	Diethyl ether, ethyl acetate, Aqueous and methanolic extract	Inhibits classical complement activation, chemiluminescence and T-cell proliferation.	Immunomodulatory	*In vivo*	(Smit, 2000)
Heracleum persicum Desf. (Apiaceae)	*Golpar*	Fruit	Flavonoids and furanocoumarins	Aquous extract	Stimulates both antibody and cellular immunity and increases IL-6 secretion.	Immunomodulatory	*In vivo*	(Sharififar *et al.,* 2009)
Cissampelos pareira linn.	*Paatha*	Root	Hayatine and alkaloids	Aqueous and ethanolic extract	Enhances humoral and T-cell immunity, boosting antibody levels and immunoglobulin titers.	Immunomodulatory	*In vivo*	(Bafna *et al.,* 2009)
Abutilon indicum linn.	*Atibaala*	Whole plant	Flavonoids and triterpenoids	Aqueous and ethanolic extract	Boosts humoral immunity by increasing immunoglobulins and strengthens innate immunity *via* neutrophil adhesion.	Immunomodulatory	*In vivo*	(Dashputre and Naikwade, 2010)
Chlorophytum borivilianum Sant. F	*Safed musli*	Root	Sapogenins	Ethanolic extract	Enhances DTH response, neutrophil adhesion, phagocytosis and carbon clearance.	Immunomodulatory	*In vivo*	(Thakur *et al.,* 2006)
Hibiscus rosa sinensis linn.	*Japaa*	Flower	Cyclopropanoids	Hydroalcoholic and ethanolic extract	Stimulates both humoral and cell-mediated immunity by enhancing phagocytic activity.	Antidiarrheal, anti-inflammatory and immunomodulatory	*In vivo*	(Gaur *et al.,* 2009)
Asparagus racemosus willd	*Satavari*	Root leaf	Beta-sitosterol, glycosides and saponin	Ethanolic extract	Inhibits TNF-α production in THP-1 cells.	Antiulcer, immunomodulatory, antidiarrheal	*In vitro* and *in vivo*	(Plangsombat *et al.,* 2016; Battu and Kumar, 2010)

COX-2- Cyclooxygenase-2; NF-κB- Nuclear factor kappa-light-chain-enhancer of activated B cells; MAPK- Mitogen-activated protein kinase; ROS- Reactive oxygen species; TNF-α- Tumor necrosis factor-alpha; ECS- Endocannabinoid system; IL-6- Interleukin-6; IL-4- Interleukin-4; IL-13- Interleukin-13; TNF-α- Tumor Necrosis Factor-alpha; LOX- Lipoxygenase; LPS-

del Cacho *et al.,* 2012; Sheridan *et al.,* 2013; Shoaib *et al.,* 2017).

Cytokine and chemokine regulation of immune dynamics

Cytokines and chemokines are crucial for immunity against *Eimeria*, affecting both protection and disease severity (Hong *et al.,* 2006). Early responses of IFN-γ and IL-10 enhance resistance, while delayed responses increase susceptibility (Bremner *et al.,* 2021). IFN-γ inhibits parasite growth and promotes Th1 type response (Dimier *et al.,* 1998) and pro-inflammatory cytokines like IL-1β and IL-18 trigger inflammation and further IFN-γ production




Fig 1: A stepwise flow outlining major steps of the review process.




Fig 2: Life cycle of coccidia: Environmental transmission and host-dependent developmental stages.




Fig 3: Mechanistic flowchart of plants exhibiting anti-coccidial effects.

(Hong *et al.,* 2006; Dalloul *et al.,* 2007). IL-2 activates T and NK cells against *E. acervulina* (Ding *et al.,* 2004) and IL-6 supports B cell differentiation and antibody production (Hong *et al.,* 2006). IL-10 controls inflammation but can suppress Th1 responses, aiding parasite survival (Rothwell *et al.,* 2004). TNF-α recruits neutrophils and drives tissue inflammation during primary infection (Zhang *et al.,* 1995), while chemokines such as MIP-1β and K203 guide immune cells to infection sites for effective gut defense (Laurent *et al.,* 2001).

Microbial dysbiosis and compromised gut integrity

The gut microbiota is vital for immune balance, metabolism and protection against pathogens. *Eimeria tenella* infection disrupts the cecal microbial community, causing dysbiosis (Huang *et al.,* 2018; Macdonald *et al.,* 2017). Beneficial bacteria like *Lactobacillus* and *Faecalibacterium* decline, while harmful bacteria such as *Clostridium and Escherichia-Shigella* increase (Chen *et al.,* 2020; Cui *et al.,* 2017). Loss of *Faecalibacterium*, a major butyrate producer, weakens mucosal integrity and anti-inflammatory responses, worsening gut damage and delaying recovery (Chen *et al.,* 2020).

CONCLUSION

The present review highlights medicinal plants with anti-inflammatory, antidiarrheal, anti-ulcer and immunomodulatory properties, emphasizing their potential as natural treatments against coccidial infections. These “phyto-coccidial” agents, noted in traditional Indian texts, may target multiple stages of the parasite’s life cycle. Although largely unexplored for anticoccidial use, they offer promising candidates for future research aimed at developing safe, plant-based therapies for coccidiosis, presenting a novel direction for drug discovery and disease management.

ACKNOWLEDEGMENT

None.

Disclaimers

The views and conclusions expressed in this article are solely those of the authors and do not necessarily represent the views of their affiliated institutions. The authors are responsible for the accuracy and completeness of the information provided but do not accept any liability for any direct or indirect losses resulting from the use of this content.

Informed consent

Not applicable.

Conflict of interest

The authors declare that there are no conflicts of interest regarding the publication of this article. No funding or sponsorship influenced the design of the study, data collection, analysis, decision to publish, or preparation of the manuscript.

REFERENCES

Ahmad, M., Jahan, N., Rehman, A.B. and Muhammad, S. (2014). Antidiarrhoeal, anti-inflammatory and analgesic activities of symplocos racemesa roxb. Bark. *Pakistan J. of Pharmaceutical Sciences*. **27(6):** 2221-2226.

Ambrose, T. and Simmons, A. (2019). Cannabis, cannabinoids and the endocannabinoid system-is there therapeutic potential for inflammatory bowel disease? *Journal of Crohn’s and Colitis*. **13(4):** 525-535.

Arstila, T.P., Vainio, O. and Lassila, O. (1994). Central role of CD4+T cells in avian immune response. *Poultry Science*. **73(7):** 1019-1026.

Arul, V., Miyazaki, S. and Dhananjayan, R. (2005). Studies on the anti-inflammatory, antipyretic and analgesic properties of the leaves of *Aegle marmelos* Corr. *Journal of Ethnopharmacology*. **96(1-2):** 159-163.

Balkan, İ.A., Goren, A.C., Kirmizibekmez, H. and Yesilada, E. (2018). Evaluation of the *in vitro* anti-inflammatory activity of *Nerium oleander* L. flower extracts and activity-guided isolation of the active constituents. *Records of Natural Products.* **12(2):** 128-141.

Awais, M.M., Akhtar, M., Iqbal, Z., Muhammad, F. and Anwar, M.I. (2012). Seasonal prevalence of coccidiosis in industrial broiler chickens in Faisalabad, Punjab, Pakistan. *Tropical Animal Health and Production*. **44:** 323-328.

Baburao, B., Reddy, A.R.N., Kiran, G., Reddy, Y.N. and Mohan, G.K. (2010). Antioxidant, analgesic and anti-inflammatory activities of *Leucas cephalotes* (Roxb. ex Roth) Spreng. *Brazilian Journal of Pharmaceutical Sciences*. **46:** 525-529.

Bafna, A. and Mishra, S. (2009). Antioxidant and immunomodulatory activity of the alkaloidal fraction of *Cissampelos pareira* Linn. *Scientia Pharmaceutica*. **78(1):** 21.

Bangoura, B. and Bardsley, K.D. (2020). Ruminant coccidiosis. *Veterinary Clinics: Food Animal Practice.* **36(1):** 187-203.

Barre, A., Hirabe, A.M., Mohamed, A.A., Ibrahim, A.A., Mohamed, H.E., Adan, M.A. and Mohamed, N.A. (2023). Prevalence, public health and associated risk factors of coccidiosis in small ruminants at deyniile sub-districtin mogadishu, somalia. *Journal of Veterinary Medicineand Animal Sciences*. **6(1):** 112.

Battu, G.R. and Kumar, B.M. (2010). Anti-inflammatory activity of leaf extract of *Asparagus racemosus* willd. *International Journal of Chemical Sciences.* 20103270294, English, Journal article, India, 0972-768X, Udaipur, Sadguru Publications. **8(2):** 1329-1338.

BR, I., MO, A., AA, I. and OO, W. (2011). Antidiarrhoeal activity of aqueous leaf extract of momordica charantia in rats. *J. Pharmacogn Phyther*. **3(1):** 1-7.

Bremner, A., Kim, S., Morris, K.M., Nolan, M.J., Borowska, D., Wu, Z. and Vervelde, L. (2021). Kinetics of the cellular and transcriptomic response to *Eimeria maxima* in relatively resistant and susceptible chicken lines. *Frontiers in Immunology*. **12:** 653085.

Chang, S.S., Huang, H.T., Wei, W.C., Lo, I.W., Lin, Y.C., Chao, C.H. and Liaw, C.C. (2023). Anti-inflammatory effect of euphane-and tirucallane-type triterpenes isolated from the traditional herb *Euphorbia neriifolia* L. *Frontiers in Chemistry*. **11:** 1223335.

Chapman, H.D., Jeffers, T.K. and Williams, R.B. (2010). Forty years of monensin for the control of coccidiosis in poultry. *Poultry Science*. **89(9):** 1788-1801.

Chen, H.L., Zhao, X.Y., Zhao, G.X., Huang, H.B., Li, H.R., Shi, C.W. and Yang, G.L. (2020). Dissection of the cecal microbial community in chickens after *Eimeriatenella* infection. *Parasites and Vectors*. **13:** 1-15.

Chen, Y., Chi, L., Liang, X., Shi, Y., Wu, T., Ye, M., Du, Z. (2020). Essential oils of *Cedrus deodara* leaves exerting anti-inflammation on TPA-induced ear edema by inhibiting COX-2/TNF-α/NF-κB Activation. *Journal of Essential Oil Bearing Plants*. **23(3):** 422-431.

Cocetta, V., Governa, P., Borgonetti, V., Tinazzi, M., Peron, G., Catanzaro, D., Berretta, M., Biagi, M., Manetti, F., Dall'Acqua, S., Montopoli, M. (2021). Cannabidiol isolated from *Cannabis sativa* L. protects intestinal barrier from *in vitro* inflammation and oxidative stress. *Front. Pharmacol.* **12:** 641210.

Conway, D.P. and McKenzie, M.E. (2007). Poultry Coccidiosis: Diagnostic and Testing Procedures. Blackwell Publishing (3rd ed.). pp. 7-16.

Cornelissen, J.B.W.J., Swinkels, W.J.C., Boersma, W.A. and Rebel, J.M.J. (2009). Host response to simultaneous infections with *Eimeria acervulina, maxima* and *tenella*: A cumulation of single responses. *Veterinary Parasitology*. **162(1-2):** 58-66.

Costa, J.F.O., David, J.P.L., David, J.M., Giulietti, A.M., Queiroz, L.P., Santos, R.R., Soares, M.B.P. (2008). Immunomodulatory activity of extracts from *Cordia superba* cham and *Cordia rufescens* A. DC. (*Boraginaceae*), plant species native from Brazilian semiarid. *Rev. Bras. Farmaacogn.* **18(1):** 11e5.

Cui, N., Wang, X., Wang, Q., Li, H., Wang, F. and Zhao, X. (2017). Effect of dual infection with *Eimeriatenella* and subgroup javian leukosis virus on the cecal microbiome in specific-pathogen-free chicks. *Frontiers in Veterinary Science*. **4:** 177.

Dahikar, S.B., Bhutada, S.A. (2017). Evaluation of anti-diarrheal activity in seed extracts of *Pongamia pinnata* (Fabaceae). *Indian Journal of Applied Research.* **7(5).** 2249-555X.

Dalloul, R.A. and Lillehoj, H.S. (2006). Poultry coccidiosis: Recent advancements in control measures and vaccine development. *Expert Review of Vaccines*. **5(1):** 143-163.

Dalloul, R.A., Bliss, T.W., Hong, Y.H., Ben-Chouikha, I., Park, D.W., Keeler, C.L. and Lillehoj, H.S. (2007). Unique responses of the avian macrophage to different species of *Eimeria*. *Molecular Immunology*. **44(4):** 558-566.

Das, S.C., Bhadra, S., Roy, S., Saha, S.K., Islam, M.S. and Bachar, S.C. (2012). Analgesic and anti-inflammatory activities of ethanolic root extract of *Swertia chirata* (*Gentianaceae*). *Biological*. **5(1):** 131.

Dashputre, N.L., Naikwade, N.S. (2010). Immunomodulatory activity of *Abutilon indicum* Linn on albino mice. *International Journal of Pharma Sciences and Research.* **1(3):** 178e84.

Dau, N.T., Thu, H.T.V., Trang, N.T.M. and Linh, N.T. (2021). Genetic characteristics of antibiotic resistance gene of vibrio cholerae isolated. *International Journal of Veterinary Science.* **10(2):** 89-95.

del Cacho, E., Gallego, M., Lee, S.H., Lillehoj, H.S., Quilez, J., Lillehoj, E.P. and Sánchez-Acedo, C. (2012). Induction of protective immunity against *Eimeria tenella, Eimeria maxima and Eimeria acervulina* infections using dendritic cell-derived exosomes. *Infection and Immunity*. **80(5):** 1909-1916.

Deshpande, J.R., Choudhary, A.A., Mirsha, M.R., Meghre, V.S., Wadokar, S.G., Dorle, A.K. (2008). Benefical effects of *Lagenaria siceraria* mol. fruit epicarp in animal models. *Indian J. Exp. Biol.* **46:** 234e42.

Dey, P., Chaudhuri, T.K. (2015). Anti-inflammatory activity of Nerium indicum by inhibition of prostaglandin E2 in murine splenic lymphocytes. *Indian J. Pharmacol.* **47(4):** 447-450.

Dimier, I.H., Quéré, P., Naciri, M. and Bout, D.T. (1998). Inhibition of *Eimeria tenella* development *in vitro* mediated by chicken macrophages and fibroblasts treated with chicken cell supernatants with IFN-γ activity. *Avian Diseases*. pp. 239-247.

Ding, X., Lillehoj, H.S., Quiroz, M.A., Bevensee, E. and Lillehoj, E.P. (2004). Protective immunity against *Eimeriaacervulina* following in ovo immunization with a recombinant subunit vaccine and cytokine genes. *Infection and Immunity.* **72(12):** 6939-6944.

Fraternale, D., Teodori, L., Rudov, A., Prattichizzo, F., Olivieri, F., Guidarelli, A., Albertini, M.C. (2018). The *in vitro* Activity of *Angelica archangelica* L. essential oil on inflammation. *J. Med Food.* **21(12):** 1238-1243.

Gaur, K., Kori, M.L. and Nema, R.K. (2009). Comparative screening of immunomodulatory activity of hydro-alcoholic extract of hibiscus rosa sinensis Linn. and ethanolic extract of *Cleome gynandra* Linn. *Global J. Pharmacol.* **3(2):** 85-89.

Ghaisas, M.M., Shaikh, S.A. and Deshpande, A.D. (2009). Evaluation of the immunomodulatory activity of ethanolic extract of the stem bark of *Bauhinia variegata* Linn. *International Journal of Green Pharmacy.* **3(1).** doi: 10.4103/0973-8258.49379.

Giresha, A.S., Pramod, S.N., Sathisha, A.D., Dharmappa, K.K. (2017). Neutralization of inflammation by inhibiting *in vitro* and *in vivo* secretory phospholipase A_2 by ethanol extract of *Boerhaavia diffusa* L. *Pharmacognosy Res.* **9(2):** 174-181.

Gokila, T., Rajalekshmi, M., Haridasan, C., Hannah, K. (2014). Plant Parts and Extracts Having Anticoccidial Activity. World Intellectual Property Organization. no. WO2014004761.

Gunakkunru, A., Padmanaban, K., Thirumal, P., Pritila, J., Parimala, G., Vengatesan, N., Gnanasekar, N., Perianayagam, J.B., Sharma, S.K., Pillai, K.K. (2005). Anti-diarrhoeal activity of *Butea monosperma* in experimental animals. *J. Ethnopharmacol.* **98(3):** 241-244.

Gupta, M., Sasmal, S. and Mukherjee, A. (2013). Evaluation of anti-inflammatory activity of *Saraca asoca* seeds using cotton pellet induced granuloma method in rats. *Int. J. Pharm. Phytopharm Res*. **2(4):** 268-271.

Halder, S., Bharal, N., Mediratta, P.K., Kaur, I., Sharma, K.K. (2009). Antiinflammatory, immunomodulatory and anti-nociceptic activity of *Terminalia arjuna* Roxb. bark powder in mice and rats. *Indian J. Exp Biol.* **47:** 577e83.

Hatam-Nahavandi, K., Carmena, D., Rezaeian, M., Mirjalali, H., Rahimi, H.M., Badri, M., Eslahi, A.V., Shahrivar, F.F., Oliveira, S.M.R. and Pereira, M.D.L. (2023). Gastrointestinal parasites of domestic mammalian hosts in Southeastern Iran. *Veterinary Sciences*. **10(4):** 261.

Haug, A., Gjevre, A.G., Thebo, P., Mattsson, J.G. and Kaldhusdal, M. (2008). Coccidial infections in commercial broilers: Epidemiological aspects and comparison of *Eimeria* species identification by morphometric and polymerase chain reaction techniques. *Avian Pathology*. **37(2):** 161-170.

Hong, Y.H., Lillehoj, H.S., Lee, S.H., Dalloul, R.A. and Lillehoj, E.P. (2006). Analysis of chicken cytokine and chemokine gene expression following *Eimeria acervulina* and *Eimeria tenella* infections. *Veterinary Immunology and Immunopathology*. **114(3-4):** 209-223.

Hong, Y.H., Lillehoj, H.S., Lillehoj, E.P. and Lee, S.H. (2006). Changes in immune-related gene expression and intestinal lymphocyte subpopulations following *Eimeria maxima* infection of chickens. *Veterinary Immunology and Immunopathology*. **114(3-4):** 259-272.

Hu, T.Y., Ju, J.M., Mo, L.H., Ma, L., Hu, W.H., You, R.R., Chen, X.Q., Chen, Y.Y., Liu, Z.Q., Qiu, S.Q., Fan, J.T., Cheng, B.H. (2019). Anti-inflammation action of xanthones from *Swertia chirayita* by regulating COX-2/NF-κB/MAPKs/Akt signaling pathways in RAW 264.7 macrophage cells. *Phytomedicine.* **55:** 214-221.

Huang, G., Tang, X., Bi, F., Hao, Z., Han, Z., Suo, J. and Liu, X. (2018). *Eimeria tenella* infection perturbs the chicken gut microbiota from the onset of oocyst shedding. *Veterinary Parasitology.* **258:** 30-37.

Jawhari, F.Z., El Moussaoui, A., Bourhia, M., Imtara, H., Mechchate, H., Es-Safi, I., Ullah, R., Ezzeldin, E., Mostafa, G.A., Grafov, A., Ibenmoussa, S., Bousta, D., Bari, A. (2020). *Anacyclus pyrethrum* (L): Chemical composition, analgesic, anti-inflammatory and wound healing properties. *Molecules*. **25(22):** 5469.

Kaur, S., Singla, L.D., Sandhu, B.S., Bal, M.S. and Kaur, P. (2019). Coccidiosis in goats: Pathological observations on intestinal developmental stages and anticoccidial efficacy of amprolim. *Indian Journal of Animal Research*. **53(2):** 245-249. doi: 10.18805/ijar.B-3471.

Keeton, S.T.N. and Navarre, C.B. (2018). Coccidiosis in largeand small ruminants. *Veterinary Clinics: Food Animal Practice*. **34(1):** 201-208.

Kendri, S.S. and Wari, U.G. (2015). Screening of the antiinflammatory activity of "*Trianthema portulacastrum*" in acute models of inflammation. *Journal of Evolution of Medical and Dental Sciences*. **4(30):** 5185-5190.

Khare, C.P. (2008). Indian Medicinal Plants: An Illustrated Dictionary. Springer Science and Business Media.

Kim, W.H., Chaudhari, A.A. and Lillehoj, H.S. (2019). Involvement of T cell immunity in avian coccidiosis. *Frontiers in Immunology*. **10:** 2732.

Kim, Y.N., Sim, K.S., Park, S., Sohn, H.Y., Kim, T. and Kim, J.H. (2022). *In vitro* and *in vivo* anti-inflammatory effects of cannabis sativa stem extract. *Journal of Medicinal Food*. **25(4):** 408-417.

Kundur, S., Karnati, P.R. and Daggu, M.R. (2012). Anti-inflammatory and analgesic activity of leaf and callus extracts of *Coleus forskohlii*. *International Journal of Phytomedicine*. **4(4):** 496.

Laurent, F., Mancassola, R., Lacroix, S., Menezes, R. and Naciri, M. (2001). Analysis of chicken mucosal immune response to *Eimeria tenella* and *Eimeria maxima* infection by quantitative reverse transcription-PCR. *Infection and Immunity.* **69(4):** 2527-2534.

Lillehoj, H.S. (1989). Intestinal intraepithelial and splenic natural killer cell responses to eimerian infections in inbred chickens. *Infection and Immunity*. **57(7):** 1879-1884.

Lillehoj, H.S. (1998). Role of T lymphocytes and cytokines in coccidiosis. *International Journal for Parasitology*. **28(7):** 1071-1081.

Lillehoj, H.S. and Choi, K.D. (1998). Recombinant chicken interferon-gamma-mediated inhibition of *Eimeria tenella* development *in vitro* and reduction of oocyst production and body weight loss following *Eimeria acervulina* challenge infection. *Avian Diseases*. pp. 307-314.

Macdonald, S.E., Nolan, M.J., Harman, K., Boulton, K., Hume, D.A., Tomley, F.M. and Blake, D.P. (2017). Effects of *Eimeria tenella* infection on chicken caecal microbiome diversity, exploring variation associated with severity of pathology. *Plos One*. **12(9):** e0184890.

Maitra, S. and Satardekar, K. (2017). Pharmacological study of symplocos racemose Roxb. *International Journal of Pharmacognosy and Phytochemical Research. Human.* **9(1):** 106-113.

Malairajan, P., Venu Babu, G., Saral, A. and Mahesh, S. (2013). Gitanjali anti-ulcer activities of *Sphaeranthus indicus* Linn. *Int. J. Drug Dev. and Res*. **5(1):** 43-46.

Malik, J.K., Manvi, F.V., Nanjwade, B.K., Alagawadi, K.R., Sinsh, S. (2009). Immunomodulatory activity of Gymnema sylvestre R. Br. leaves on *in vitro* human neutrophils. *J. Pharm. Res.* **2(8):** 1284e6.

Manisha, S.M.S. and Kareppa, B.M. (2009). Studies on utility of *Adhatoda zeylanica* medic. *Asian Journal of Bio Science.* **4(2):** 304-306.

Menon, D.B., Latha, K. (2011). Phytochemical screening and *in vitro* anti-inflammatory activity of the stem of coleus forskohlii. *Pharm. J.* **3(23):** 75-79.

Muthulingam, M. and Veerakumar, D. (2021). A precious remedial plant *Asteracantha longifolia* (Nees.). *Comprehensive Review.* **10:** 37-58.

Manimekalai, P., Maheshwari, P., Velmurugan, R., Gurumoorthy, M., Kumar, S.H. and Vijayakumar, G. (2018). Gastro protective effect of standardized ethanolic leaf extract of indigofera tinctoriao on experimental gastric ulcers in rats. *Research Journal of Pharmacy and Technology*. **11(2):** 527-531.

Murshed, M., Aljawdah, H.M., Mares, M.M. and Al-Quraishy, S. (2023). *In vivo* anticoccidial efficacy of *Calotropis procera* leaf extract as an alternative treatment against *Eimeria papillate* infection in mice. *Indian Journal of Animal Research.* **59(1):** 85-92. doi: 10.18805/IJAR.BF-1607.

Palit, P., Mandal, S.C., Bhunia, B. (2016). Total steroid and terpenoid enriched fraction from *Euphorbia neriifolia* Linn offers protection against nociceptive-pain, inflammation and *in vitro* arthritis model: An insight of mechanistic study. *Int. Immunopharm.* **41:** 106-115.

Palit, P., Mukherjee, D., Mahanta, P., Shadab, M., Ali, N., Roychoudhury, S., Asad, M., Mandal, S.C. (2018). Attenuation of nociceptive pain and inflammatory disorders by total steroid and terpenoid fraction of *Euphorbia tirucalli* Linn root in experimental *in vitro* and *in vivo* model. *In Inflammopharmacology.* pp. 235-250.

Panax ginseng. (2009). Monograph. *Altern. Med. Rev.* **14:** 172e6.

Patel, N.K., Khan, M.S., Bhutani, K.K. (2015). Investigations on *Leucas cephalotes* (Roth.) spreng for inhibition of LPS-induced pro-inflammatory mediators in murine macrophages and in rat model. *EXCLI J.* **10(14):** 508-516.

Pattison, M., McMullin, P., Bradbury, J.M. and Alexander, D. (2007). *Poultry diseases*. Elsevier Health Sciences.

Paul, B.T., Jesse, F.F.A., Chung, E.L.T., Che'Amat, A. and Lila, M.A.M. (2020). Risk factors and severity of gastrointestinal parasites in selected small ruminantsfrom Malaysia. *Veterinary Sciences*. **7(4):** 2.

Philomina, M.P. and Malarvizhi, A. (2022). Anti-inflammatory and antioxidant effect of various extracts of Coleus forskohlii root. *Intern. J. Zool. Invest.* **8(Special Issue):** 106-112.

Plangsombat, N., Rungsardthong, K., Kongkaneramit, L., Waranuch, N., Sarisuta, N. (2016). Anti-inflammatory activity of liposomes of *Asparagus racemosus* root extracts prepared by various methods. *Exp. Ther. Med*. **12(4):** 2790-2796.

Rajkapoor, B., Jayakar, B., Anandan, R., Kavimani, S. (2003). Anti-ulcer effect of *Bauhinia variegata* Linn. in rats. *Journal of Naturalremedies.* **3(2):** 21.

Ram, H., Garg, R. and Banerjee, P.S. (2019). Coccidiosis diseases, symptoms and prevention in chickens. *Bhartiya Krishi Anusandhan Patrika*. **34(3 and 4):** 263-266. doi: 10.18805/BKAP179.

Rathor, S.R. and Goyal, H.R. (1973). Studies on the anti-inflammatory and anti-arthritic activity of an indian medicinal plant, *Cedrus deodara. Indian Journal of Pharmacology*. **5(2):** 334-343.

Reddy, N.S., Santhosha, D., Reddy, P.V. and Venu, C. (2015). Evaluation of antidiarrhoeal activity cleome gynandra leaf extracts on magnesium sulphate-and castor oil-induced diarrhoea in wistar rats. *Research Journal of Pharmacology and Pharmacodynamics*. **7(4):** 184-186.

Reid, W.M. (1990). History of avian medicine in the United States. X. Control of coccidiosis. *Avian Diseases.* **34(3):** 509-525.

Rinku, M., Prasanth, V.V., Parthasarathy, G. (2009). Immunomodulatory activity of the methanolic extract of *Urena lobata* Linn. *Int. J. Pharmacol.* **7:** http://www.ispub.com/journal/the_internet_ journal_of_pharmacology/volume_7_number_1_27/article/ immunomodulatory-activity-of-the-methanolic-extract-ofurena-lobata-linn.html,1 [accessed 2009].

Rothwell, L., Young, J.R., Zoorob, R., Whittaker, C.A., Hesketh, P., Archer, A. and Kaiser, P. (2004). Cloning and characterization of chicken IL-10 and its role in the immune response to *Eimeria maxima*. *The Journal of Immunology.* **173(4):** 2675-2682.

Saeed, Z. and Alkheraije, K.A. (2023). Botanicals: A promising approach for controlling cecal coccidiosis in poultry. *Frontiers in Veterinary Science*. **10:** 1157633.

Sahoo, H.B., Sahoo, S.K., Sarangi, S.P., Sagar, R., Kori, M.L. (2014). Anti-diarrhoeal investigation from aqueous extract of *Cuminum cyminum* Linn. *Seed in Albino Rats. Pharmacognosy Res.* **6(3):** 204-209.

Saif, Y.M. and Fadley, A.M. (2009). Diseases of Poultry (12th ed). Blackwell Publishing.

Schnitzer, J.K., Berzel, S., Fajardo-Moser, M., Remer, K.A. and Moll, H. (2010). Fragments of antigen-loaded dendritic cells (DC) and DC-derived exosomes induce protective immunity against *Leishmania major*. *Vaccine*. **28(36):** 5785-5793.

Semuel, M.R. and Shabaraya, A.R. (2011). Antiinflammatory activity of fractionated extracts of *Achyranthes aspera* Linn leaves. *Journal of Applied Pharmaceutical Science*. **01(08):** 188-190.

Shah, S.A., Wakade, A.S., Juvekr, A.R. (2007). Immunomodulatory activity methanolic extract of *Murraya koenigii* (L.) spreng leaves. *Indian J. Exp. Biol.* **46:** 505e9.

Shah, A.J., Gowani, S.A., Zuberi, A.J., Ghayur, M.N. and Gilani, A.H. (2010). Antidiarrhoeal and spasmolytic activities of the methanolic crude extract of *Alstonia scholaris* L. are mediated through calcium channel blockade. *Phytotherapy Research: An International Journal Devoted to Pharmacological and Toxicological Evaluation of Natural Product Derivatives*. **24(1):** 28-32.

Shah, V.V., Shah, V.K., Sheth, N.R. and Patel, M.M. (2017). Hepatoprotective and anti-inflammatory activities of Hydroalcoholic extract of Bark of *Erythrina indica*. *Research Journal of Pharmacology and Pharmacodynamics.* **9:** 189-194.

Sharififar, F., Pournourmohammadi, S., Arabnejad, M., Rastegarianzadeh, R., Ranjbaran, O., Purhemmaty, A. (2009). Immunomodulatory activity of aqueous extract of *Heracleum persicum* Desf. in mice. *Iran J. Pharm. Res.* **8(4):** 287e92.

Sheridan, B.S., Romagnoli, P.A., Pham, Q.M., Fu, H.H., Alonzo, F., Schubert, W.D. and Lefrançois, L. (2013). γδ T cells exhibit multifunctional and protective memory in intestinal tissues. *Immunity.* **39(1):** 184-195.

Shinde, U.A., Kulkarni, K.R., Phadke, A.S., Nair, A.M., Mungantiwar, A.A., Dikshit, V.J., Saraf, M.N., (1999a). Mast cell stabilizing and lipoxygenase inhibitory activity of *Cedrus deodara* (roxb.) Loud. wood oil. *Indian J. Exp. Biol.* **37(3):** 258-261.

Shinde, U.A., Phadke, A.S., Nair, A.M., Mungantiwar, A.A., Dikshit, V.J., Saraf, M.N. (1999b). Membrane stabilizing activity-A possible mechanism of action for the anti-inflammatory activity of *Cedrus deodara* wood oil. *Fitoterapia.* **70(3):** 251-257.

Shivaramaiah, C., Barta, J.R., Hernandez-Velasco, X., Téllez, G. and Hargis, B.M. (2014). Coccidiosis: Recent advancements in the immunobiology of *Eimeria* species, preventive measures and the importance of vaccination as a control tool against these Apicomplexan parasites. *Veterinary Medicine: Research and Reports*. pp. 23-34.

Shoaib, M., Xiaokai, S., Ul-Hasan, M., Zafar, A., Riaz, A., Umar, S. and Xiangrui, L. (2017). Role of dendritic cells in immunity against avian coccidiosis. *World's Poultry Science Journal*. **73(4):** 737-746.

Shridhar, B., Mani, M., Kumar, P., Yadav, S. and Mahaur, K.K. (2011). Anti-diarrhoeal activity of ethanolic extract of heartwood of *Pterocarpus marsupium* roxb. *Journal of Drug Delivery and Therapeutics.* **8(6-s):** 294-297.

Sindhu, R.K. and Arora, S. (2014). Anti-inflammatory potential of different extracts isolated from the roots of ficus lacor Buch. Hum and *Murraya koenigii* L. Spreng. *Archives of Biological Sciences*. **66(3):** 1261-1270.

Smit, H.F. (2000). Picrorhiza scrophularii flora from traditional use to immunomodulatory activity [doctoral thesis]. Utrecht, Netherlands: University of Utrecht.

Subramaniyan, V. and Paramasivam, V. (2017). Potential anti-inflammatory activity of Plumbago zeylanica. *Asian J. Pharm. Clin. Res*. **10(10):** 372-375.

Swathi, K.P., Jayaram, S., Sugumar, D., Rymbai, E. (2020). Evaluation of anti-inflammatory and anti-arthritic property of ethanolic extract of *Clitoria ternatea*. *Chin Herb Med.* **13(2):** 243-249.

Thakur, M., Bhargava, S., Dixit, V.K. (2006). Immunomodulatory activity of *Chlorophytum borivilianum Sant. F. Evid Based Complement Alternat Med*. **4(4):** 419e23.

Thanigavelan, V., Venkatachalam, K., Venkatachalam, L., Natarajan, S., Murugan, P.K. and Savarimuthu, J.A. (2014). Hydroalcoholic extract of Plumbago zeylanica Linn root bark exhibit analgesic and anti-inflammatory activities in experimental rat models. *Am. J. Pharm Health Res*. **2(4):** 209-221.

Uddin, S.J., Mondal, K., Shilpi, J.A. and Rahman, M.T. (2006). Antidiarrhoeal activity of *Cyperus rotundus*. *Fitoterapia*. **77(2):** 134-136.

Uthirapathy, S. (2021). Analgesic and Anti-inflammatory activity of *withania somnifera* root extract. *Journal of Pharmaceutical Research International*. **33(41A):** 75-84.

Wankhede, S., Juvekar, M., Juvekar, A., Sakat, S., Gambhire, M. (2009). Study of *in vitro* and *in vivo* anti-inflammatory activity of aqueous extract of leaves *Erythrina indica*. *Planta Med.* **75(9):** PJ74.

Woliñska, R., Zalewska, M., Poznañski, P., Nawrocka, A., Kowalczyk, A., Sacharczuk, M., Bujalska-Zadrożny, M. (2025). Anti-inflammatory activity of *Cannabis sativa* L. extract in 2,4-dinitrochlorobenzene-induced dermatitis in rats. *Pharmaceuticals (Basel).* **18(3):** 370.

Yang, W.C., Tien, Y.J., Chung, C.Y., Chen, Y.C., Chiou, W.H., Hsu, S.Y. and Chang, C.L.T. (2015). Effect of *Bidens pilosa* on infection and drug resistance of *Eimeria* in chickens. *Research in Veterinary Science*. **98:** 74-81.

Zhang, S., Lillehoj, H.S. and Ruff, M.D. (1995). Chicken tumor necrosis-like factor: 1. *in vitro* production by macrophages stimulated with *Eimeria tenella* or bacterial lipopoly saccharide. *Poultry Science*. **74(8):** 1304-1310.

Zhao, C., Nguyen, T., Liu, L., Sacco, R.E., Brogden, K.A. and Lehrer, R.I. (2001). Gallinacin-3, an inducible epithelial β-defensin in the chicken. *Infection and Immunity*. **69(4):** 2684-2691.