RESEARCH ARTICLE Indian Journal of Animal Research, Volume 57 Issue 6: 806-810 (June 2023)

# Comparative Evaluation of Hatching Traits in Four Phenotypes of Naked Neck Chicken

M. Shafiq[1], M.T. Khan[2], M. Rauf[3], F. Raziq[4], E. Bughio[5], M.A. Gondal[6], M.M. Jalees[7], S. Liaqat[8]

10.18805/IJAR.BF-1449

## ABSTRACT

**Background:** In Pakistan, the naked neck chickens are characterized on the base of plumage colors such as black, white black, light brown and dark brown. There is scanty of information on hatching traits in different phenotypes of naked neck chicken. The current study aimed to investigate and compare the hatching traits in four phenotypes of naked neck chicken (black, white black, light brown and dark brown).

**Methods:** In total, 300 females (20 weeks old) comprising four plumage colors (black, white black, light brown and dark brown), 75 from each, were randomly assigned to 20 replicates (15/replicate) under a completely randomized design (CRD). Number of settable eggs, fertility, hatch of fertile, hatchability, embryonic mortality and chick quality parameters were evaluated.

**Result:** The results showed higher settable eggs, fertility and hatchability in black, dark brown and light brown phenotypes than that of the white black whereas better hatch of fertile and reduced embryonic mortality were recorded in black and dark brown phenotypes as compared to white black. Better chick quality was observed in dark brown phenotype followed by light brown, black and then white black. Phenotypes other than white black had better hatching performance, hence could be used to revive backyard poultry farming.

**Key words:** Hatching traits, Naked neck, Performance, Phenotype.

## INTRODUCTION

There has been rapid increase in the number of farmers keeping chicken parent and grandparent stock leading to increase in the population of meat and egg type chickens in the world (Kathleen, 2002). Among other factors, poor fertility and hatchability rates constitute the major threat to performance of the industry (Adebambo, 2005). Fertility and hatchability characteristics of egg type chickens are the two main traits of assessing the potentials of hens (Allanah *et al.,* 2014). Fertility is an important parameter in chicken that reflects the total actual reproductive capacity of females. An egg is said to be infertile when it fails to display any evidence of developing embryo (Miazi *et al.,* 2012). Hatchability is referred as the ability of the embryo to successfully escape from the shell (Tarek, 1992). Good hatchability of eggs is to some extent heritable, but is determined by a complicated genetic constitution and the environment. Hatchability is a trait of economic importance in the chicken industry because it has a strong effect on chick output (Wolc *et al.,* 2010). Number of factors such as egg weight, turning of eggs, storage, humidity, shell strength, egg size and genetic factors influence hatchability. According to Peters *et al.* (2008) the genetic make-up of an individual chicken is fixed at fertilization. Hence fertility and hatchability are generally considered as traits of two parents.

Indigenous chicken are widely distributed in the rural areas of tropical and sub-tropical countries where they are kept by the majority of rural poor. Indigenous chicken from Pakistan are in general hardy, adoptive to rural environments (Khan *et al*., 2017), survive on little or no inputs and adjust of fluctuations in feed availability. They are self-reliant and hardy birds with capacity to withstand harsh weather conditions and adaptation to adverse environment (Khan *et al.,* 2017). They are known to possess qualities such as the ability to hatch on their own, brood and scavenge for

[1]Department of Livestock and Dairy Development, Poultry Research Institute, Rawalpindi-46300, Pakistan.
[2]Department of Poultry Science, Cholistan University of Veterinary and Animal Sciences, Bahawalpur-63100, Pakistan.
[3]Department of Pathology, Cholistan University of Veterinary and Animal Sciences, Bahawalpur-63100, Pakistan.
[4]Department of Livestock and Dairy Development (Extension), Khyber Pakhtunkhwa, Pakistan.
[5]Department of Poultry Production, Shaheed Benazir Bhutto University of Veterinary and Animal Sciences, Sakrand-67210, Pakistan.
[6]Institute of Continuing Education and Extension, Cholistan University of Veterinary and Animal Sciences, Bahawalpur-63100, Pakistan.
[7]Department of Microbiology, Cholistan University of Veterinary and Animal Sciences, Bahawalpur-63100, Pakistan.
[8]Faculty of Veterinary and Animal Sciences, The Islamia University of Bahawalpur-63100, Pakistan.

**Corresponding Author:** M.T. Khan, Department of Poultry Science, Cholistan University of Veterinary and Animal Sciences, Bahawalpur-63100, Pakistan. Email: mtahirkhan@cuvas.edu.pk

**How to cite this article:** Shafiq, M., Khan, M.T., Rauf, M., Raziq, F., Bughio, E., Gondal, M.A., Jalees, M.M. and Liaqat, S. (2023). Comparative Evaluation of Hatching Traits in Four Phenotypes of Naked Neck Chicken. Indian Journal of Animal Research. 57(6): 806-810. doi: 10.18805/IJAR.BF-1449.

**Submitted:** 05-10-2021 **Accepted:** 26-04-2022 **Online:** 26-05-2022

major parts of their food and possess appreciated immunity against endemic diseases. Additionally, the indigenous poultry species represent valuable resources for livestock development because their extensive genetic diversity allows for rearing of poultry under varied environmental conditions, providing a range of products and functions. Thus, great genetic resources embedded in indigenous poultry await full exploitation (Sonaiya *et al.,* 1999). In Pakistan, the naked neck chickens are characterized on the base of plumage colors such as black, white black, light brown and dark brown. There is scanty of information on hatching traits in different phenotypes of naked neck. The present study was designed to investigate and compare hatching traits in four phenotypes (black, white black, light brown and dark brown) of naked neck chicken.

## MATERIALS AND METHODS

### Experimental site, birds and husbandry

The study was executed at the Indigenous Chicken Genetic Resource Centre (ICGRC), Department of Poultry Production, University of Veterinary and Animal Sciences, Lahore, Pakistan, in 2016 for the duration of 22 weeks. In total, 340 birds (300 females and 40 males) comprising four phenotypic groups (black, white black, light brown and dark brown birds), 85 from each (75 female and 10 male), were selected from the breeding stock maintained at (ICGRC) and randomly assigned to 4 experimental groups under a completely randomized design. These birds were placed in 20 deep litter pens with 17 (15 female, 2 male) in each pen to get the fertile eggs. Same managerial conditions were applied at all experimental units. Vaccination and medication was done according to the standard practices under the supervision of a qualified veterinarian. The birds were fed iso-caloric and iso-nitrogenous layer mash diet (Table 1) from 20 weeks of age to the end of the experiment (42 weeks). The feed and water was supplied *ad libitum* through trough feeder and nipple drinking system.

### Collection, storage and setting of eggs

Collection of fertile eggs began seven days after introducing males. In total, 600 eggs (30 from each pen) were properly tagged according to the batch number, phenotype and the date of lay before sending them to the hatchery each week. After collection, the eggs were checked for cracks, morphological deformities and dirt stains (soiled). Thereafter, these eggs were stored for one to seven days in an environmentally controlled room (15 to 18°C and 75 to 80% RH) and then incubated in automatic incubator at 37.5°C (99.5°F) and 65% RH in trays identified per replicate. Proper cleaning, disinfection and fumigation practices were ensured before setting the eggs. Candling was done on the 18$^{th}$ day of incubation for the identification of fertile eggs. After candling, the fertile eggs were shifted to the hatching tray, according to the phenotypes and then into the hatcher part of the incubator. After the hatch, chicks were left in the hatchery until they were 90% dried. On the 21$^{st}$ day, the numbers of hatched chicks (including the normal, weak and abnormal) and dead-in-shell embryos were recorded separately. Chicks were weighed after being pulled out of the incubator using a 500 grams digital scale. Normal and healthy chicks having above 30 grams weight were put into the A-grade chick category.

### Data collection

Settable egg percentage was calculated as the ratio between number of settable eggs and the total number of eggs multiplied by 100. Fertility percentage was calculated as the ratio between number of fertile eggs and the total number of eggs set multiplied by 100. Hatchability percentage was calculated as the ratio between number of hatched chicks and the total number of eggs set multiplied by 100. Similarly, the hatch of fertile percentage was calculated as the ratio between number of hatched chicks and the total number fertile eggs multiplied by 100. Embryonic mortality percentage was calculated as the ratio between number of dead embryos and the total number of hatched chicks multiplied by 100. A-grade chick percentage was calculated as the ratio between number A-grade chicks and total number of chicks multiplied by 100. The data were analyzed through ANOVA technique by using the GLM procedure of SAS (SAS Institute Inc., 2002-03). Post-hoc comparison among treatment means was made through Duncan's Multiple Range (DMR) test (Duncans, 1955) at a 5% probability level.

## RESULTS AND DISCUSSION

### Settable eggs and fertility

Overall means of settable eggs showed significant differences among different phenotypes (Table 2). Black, light brown and dark brown phenotypes demonstrated higher settable eggs as compared to white black. Settable egg percentage is an economically important parameter. In the present study, variation in settable eggs might be attributed to the genetic differences among phenotypes. It has already been reported that settable egg varies due to strain (Renema *et al.,* 2001) or genotype effect (Dunga, 2013). Similarly, studies on different Aseel varieties (Khan *et al.,* 2017) and genotypes of poultry (Adedeji, 2015) displayed variations in settable eggs. Abudabos (2010), likewise, reported significant differences in hatching traits among different local and imported stocks of Japanese quails.

Means of fertility indicated variations with respect to different phenotypes (Table 2). Black, light brown and dark brown phenotypes illustrated remarkably higher fertility than that of white black. Fertility is an important parameter in chicken that shows the total actual reproductive capacity of females (Miazi *et al.,* 2012). Higher fertility was observed in black, light brown and dark brown phenotypes, which may be due to the genetic variations among the phenotypes. Similarly, it has been reported that genotype of different strains (Hussnain *et al.,* 2013) or varieties (Khan *et al.,* 2017) responds differently for fertility. Likewise, Peters *et al.* (2008)

reported a significant strain effect on fertility of eggs. Similarly, Moreki *et al.* (2014) observed fluctuation in fertility among different genotypes, strengthening the argument that genotype is the major cause of variation in fertility (Abudabos, 2010; Adedeji, 2015). Likewise, Dunga (2013) also reported significant difference in percent fertile eggs among different phenotypes of naked neck. Significant differences in fertility among different strains (Hussnain *et al.,* 2013), ecotypes (Fayeye *et al.,* 2005) and local breeds (Ensaf *et al.,* 2005) have previously been reported. In contrast, Shafik *et al.* (2013) reported no genotype effect on fertility. In the current study, the overall F values were not optimum because the experiment was conducted under very high ambient temperatures conditions, which might have reduced bird mating activity resulting in reduced fertility. A previous study also claimed negative effect of heat stress on birds' mating activity (Ernst *et al.,* 2004).

**Hatch of fertile eggs and hatchability**

Different phenotypes manifested significant differences in hatch of fertile (Table 2). Higher hatch of fertile values were observed in black and dark brown phenotype as compared to white black. It might be due to the greater fertility in hens of respective phenotypic group. In line with results, Dunga *et al.* (2013) indicated variations in hatch of fertile among different phenotypes of naked neck. Similarly, it was reported that hatching traits were different (P<0.05) among different local and imported stocks of Japanese quails (Abudabos, 2010). Similarly, Hussnain *et al.* (2013) reported improved hatch of fertile in Cobb breeder than other breeds and lower hatch of fertile in Hubbard breeders than other strains. Likewise, Adedeji (2015) revealed significant differences in hatch of fertile among different genotypes of chicken. Similarly, significant strain (Yousria *et al.,* 2010) or variety (Khan *et al.,* 2017) effect on hatch of fertile has also been reported. In contrast, Shafik *et al.* (2013) reported no correlation between hatches of fertile and genotype.

Different phenotypes exhibited pronounced effects on hatchability (Table 3). Black, light brown and dark brown phenotypes showed better hatchability than that of white black that may be as a result of high fertility in these phenotypes since only fertile eggs can hatch to chicks. In the present study, differences in hatchability may also be linked to the genetic differences among different phenotypes. Similar to these findings, Khan *et al.* (2017) reported a significant genotype effect on hatchability. Likewise, Peters *et al.* (2008) reported a significant effect of strain on hatchability of eggs. Similarly, Abudabos (2010) reported that different local and imported stocks of Japanese quails influenced hatchability significantly; indicating that hatchability varies among different strains (Hussnain *et al.,* 2013) or genotypes (Heier and Jerp, 2001) of chicken. Likewise, Elibol *et al.* (2002) reported that strain is a major factor that affects the hatchability. Moreki *et al.* (2014) also observed variations in hatchability among naked neck and normal feathered chickens, highlighting the breed effect on hatchability (Adeleke *et al.,* 2012). In contrast, Ahmad (2013) claimed no effect of genotype on hatchability.

**Embryonic mortality and A-grade chick**

In the current study, overall means of embryonic mortality indicated marked differences among different phenotypes (Table 3). Chick embryonic mortality has long been the subject of economic interest. In the present study, black and dark brown phenotypes indicated reduced embryonic mortality. It is reported that the entire process of embryonic development and successful hatching rely highly on egg external and internal quality (Narushin, 2001, Narushin and Romanov, 2002) and many factors, including breed, strain, variety, temperature, relative humidity, rearing practices and seasons, influence egg quality (Washburn, 1990). Thus, the variations in embryonic mortality in the current study may be due to the phenotypic or genetic variations. Similarly, it was reported that breed (Adedeji, 2015) or variety (Ahmad, 2013) had a significant effect on embryonic mortality. Abudabos (2010) also reported disparities in embryonic mortality among different local and imported stocks of Japanese quails, endorsing the above view that genotype is the main source of variation in embryonic mortality (Moreki *et al.,* 2014). Non-significant differences in embryonic

**Table 1:** Ingredient and nutrient composition of basal diet.

| Ingredient (%) | Quantity |
|---|---|
| Corn | 42.61 |
| SBM[1] | 15.62 |
| Corn Gluten (60%) | 1 |
| Rice tips | 19 |
| Wheat bran | 13 |
| DCP[2] | 1.2 |
| CaCO | 7.42 |
| DL-Methionine | 0.15 |
| **Nutrient** | |
| CP (%) | 15.00 |
| ME (Kcal/kg) | 2,750 |
| Calcium (%) | 1.30 |
| Phosphorous (%) | 0.40 |
| Lysine (%) | 0.70 |
| Methionine (%) | 0.30 |
| NA | 0.16 |

[1]Soybean meal.

[2]Dicalcium phosphate.

**Table 2:** Settable egg (SE), fertility (F) and hatch of fertile (HF) in four phenotypes of naked neck chicken.

| Treatment | Parameters | | |
|---|---|---|---|
| | SE (%) | F (%) | HF (%) |
| Black | 96.00±0.8[a] | 73.96±0.7[a] | 76.06±1.0[a] |
| White black | 92.33±0.8[b] | 70.74±0.7[b] | 71.39±0.7[b] |
| Light brown | 96.66±0.5[a] | 73.11±0.8[a] | 74.55±1.4[ab] |
| Dark brown | 96.00±1.0[a] | 75.32±0.7[a] | 75.05±1.2[a] |
| P-value | 0.0075 | 0.0048 | 0.0490 |

[a-c]Means within a column lacking a common superscript differ (P≤0.05).

**Table 3:** Hatchability (H), embryonic mortality (EM) and A-grade chicks (AC) in four phenotypes of naked neck chicken.

| Treatment | Parameters | | |
|---|---|---|---|
| | H (%) | EM (%) | AC (%) |
| Black | 56.25±0.7$^{a}$ | 23.93±1.0$^{b}$ | 24.10±1.2$^{bc}$ |
| White black | 50.51±0.9$^{b}$ | 28.60±0.7$^{a}$ | 20.68±1.1$^{c}$ |
| Light brown | 54.49±1.0$^{a}$ | 25.44±1.4$^{ab}$ | 27.82±0.9$^{ab}$ |
| Dark brown | 56.55±1.3$^{a}$ | 24.94±1.2$^{b}$ | 28.91±.16$^{a}$ |
| P-value | 0.0024 | 0.0490 | 0.0011 |

$^{a-c}$Means within a column lacking a common superscript differ (P≤0.05).

mortality or dead germ among different genotypes (Hussnain *et al.,* 2013; Khan *et al.,* 2017) have also been reported.

In the current study, A-grade chick quality was significantly influenced by different phenotypes (Table 3). Highest A-grade chicks were observed in dark brown followed by light brown, black and white black phenotypes. Initial chick quality has direct effect on subsequent poultry performance. Chick quality majorly depends on the type of responsible genes or gene combination. Difference in plumage color among different phenotypes is also the manifestation of genetic differences, which might have reflected variation in chick quality. Similar to the current study, discrepancies in A-grade chicks were observed in different Aseel varieties (Khan *et al.,* 2017), reflecting that chick quality depends on variety or genotype.

## CONCLUSION

Settable eggs, fertility, hatchability and hatch of fertile percentages have no significant differences among black, light brown and dark brown phenotypes whereas lesser embryonic mortality was observed in eggs of black phenotype and A-grade chicks were hatched maximum in dark brown phenotype. It was concluded that all naked neck phenotypes except white black indicated better hatching characteristics.

**Conflict of interest:** None.

## REFERENCES


Abudabos, A. (2010). The effect of broiler breeder strain and parent flock age on hatchability and fertile hatchability. International Journal of Poultry Science. 9: 231-235.

Adebambo, A.O. (2005). Physiological response of poultry to the environmental condition. Online Research Poultry Science. 8: 17-19.

Adedeji, T.A., Amao, S.R., Popoola, A.D., Ogundipe, R.I. (2015). Fertility, hatchability and eggs quality traits of Nigerian locally adapted chickens in the derived savanna environment of Nigeria. Journal of Biology, Agriculture and Healthcare. 5: 36-42.

Adeleke, M.A., Peters, S.O., Ozoje, M.O., Ikeobi, C.O., Bamgbose, A.M., Adebambo, O.A. (2012). Effect of crossbreeding on fertility, hatchability and embryonic mortality of Nigerian local chickens. Tropical Animal Health and Production. 44: 505-510.

Ahmad, Z. (2013). Pre and post-moult productive and reproductive performance, egg geometry, quality and meat composition of four varieties of native Aseel chicken. Ph.D Thesis, Department of Poultry Production, University of Veterinary and Animal Sciences, Lahore, Pakistan.

Allanah, T.O., Okonkwo, J.C., Omeje, S.I. (2014). Fertility and hatchability characterization of three strains of egg type chickens. Scientific Journal of Biological Sciences. 3: 59-68.

Duncan, D.B. (1955). Multiple ranges and multiple F tests. Biometrics. 11: 1-42.

Dunga, G.T. (2013). The effect of the Naked-Neck (*na*) and frizzling (*f*) genes on the fertility, hatchability, egg quality and pterylosis of locally developed commercial layer parent lines. M.Sc. Thesis, Department of Animal Science, Kwame Nkrumah University of Science and Technology.

Elibol, O., Peak, S.D., Brake, J. (2002). Effect of flock age, length of egg storage and frequency of turning during storage on hatchability of broiler hatching eggs. Poultry Science. 81: 945-950.

Ensaf, A.E., Abd, E., Whed, H.M., Namra, M.M., Osma, A.M.R., Hataba, N.A. (2005). Results of random sample test laying performance of nine Egyptian strains of chicken. Egyptian Poultry Science. 25: 195-208.

Ernst, R.A., Bradely, F.A., Delany, M.E., Abbot, U.K. (2004). Common Incubation Problems: Causes and Remedies. Retrieved 18 March 2013, from http//anrcatalog.ucdavis.edu.

Fayeye, T.R., Adeshiyan, A.B., Olugbami, A.A. (2005). Egg traits, hatchability and early growth performance of the Fulani-ecotype chicken. Livestock Research for Rural Development. 17: 48. http://www.lrrd.org/lrrd17/8/faye17094.htm.

Heier, B.T. and Jarp, J. (2001). An epidemiology study of the hatchability in broiler breeds flocks. Poultry Science. 80: 1132-1138.

Hussnain, F., Ishaq, H.M., Akram. M., Sahota, A.W., Baber, M.E., Hussain, J., Mehmood, S., Iqbal, A. (2013). Hatching Performance of Eggs Kept for 3 Different Storage Periods at 4 Production Phases in Three Genetic Groups of Broiler Breeder. Abstracts of 33$^{rd}$ Pakistan Congress of Zoology: 352.

Kathleen, M.M. (2002). Effect of egg storage on hatchability. Journal of Poultry Science. 2: 81-86.

Khan, M.T., Mahmud, A., Zahoor, I., Javed, K. (2017). Organic and inorganic selenium in Aseel chicken diets: Effect on hatching traits. Poultry Science. 96: 1466-1472

Miazi, O.F., Miah, G., Miazi, M.M., Uddin, M.M., Hassan, M.M., Faridahsan, M. (2012). Fertility and hatchability of Fayoumi and Sonali chicks. Scholarly Journal of Agricultural Science. 2: 83-86.

Moreki, J.C, Mmopelwa, G.M., Nthoiwa, G.P. (2014). Hatchability traits of normal feathered and Naked-Neck *Tswana* chickens reared under intensive system. International Journal of Current Microbiology and Applied Sciences. 3: 395-401.

Narushin, V.G. (2001). What Egg Parameters Predict Best its Shell Strength? In: IX European Symposium on the Quality of Eggs and Egg Products. 2001; Kusadasi, Turkey. Pages 349-355.

Narushin, V.G. and Romanov, M.N. (2002). Egg physical characteristics and hatchability. World's Poultry Science Journal. 58: 297-303.

Peters, S.O., Ilori, B.M., Ozoje, M.O., Ikeobi, C.O.N., Adebambo, O.A. (2008). Gene segregation effects on fertility and hatchability of pure and crossbred chicken genotypes in the humid tropics. International Journal of Poultry Science. 7: 954-958.

Renema, R.A., Robinson, F.E., Feddes, J.J.R., Fasenko, G.M., Zuidhoft, M.J. (2001). Effects of light intensity from photostimulation in four strains of commercial egg layers: 2. Egg production parameters. Poultry Science. 80: 1121-1131.

SAS Institute (2002-2003). SAS/STAT User's Guide: Statistics. Version 9.1.SAS Inst. Inc., Cary, NC.

Shafik, B.M.N., El-Bayomi, K.M., Sosa, G.A., Osman, A.M.R. (2013). Effect of crossing Fayoumi and Rhode Island Red on growth performance, egg and reproductive traits under Egyptian conditions. Benha Veterinary Medical Journal. 24: 11-18.

Sonaiya, E.B., Branckaert, R.D.S., Gueye, E.F. (1999). Research and Development Options for Family Poultry. First INFPD/FAO Electronic Conference on Family Poultry. December 7 1998 - March 5 1999.

Tarek, M.K. (1992). The performance of exotic breeds under scavenging cum supplementary feeding in rural condition of rearing. Unpublished M.Sc. Thesis. Department of Poultry Science. Bangladesh Agricultural University, Mymensingh.

Washburn, K.W. (1990). Genetic Variation in Egg Composition. In: Poultry Breeding and Genetics. [Crawford, R.D. (Ed.)], Elsevier Science Publisher, B V, Amsterdam. pp 781-804.

Wolc, A., White, I.M.S., Hill, W.G., Olori, V.E. (2010). Inheritance of hatchability in broiler chickens and its relationship to egg quality traits. Poultry Science. 89: 2334-2340.

Yousria, A.K.M., Aly, O.M., Abou El-Ella, N.Y. (2010). Effect of crossing on the performance of local chicken strain. 4-Effect of strain and laying age on egg quality characteristics. Egyptian Poultry Science. 30: 1171-1188.