RESEARCH ARTICLE Legume Research- An International Journal, Volume 46 Issue 7: 927-933 (July 2023)

Insecticide Resistance Development and Detoxification Enzyme Activities of *Spodoptera litura* (Fabricius) in Soybean from Kumaon Himalayas

Rashmi Joshi[1], Neeta Gaur[2], Sudha Mathpal[2]

10.18805/LR-5101

ABSTRACT

Background: Soybean is an important crop of Uttarakhand both in hills as well as in plains and *Spodoptera litura* is an important pest of soybean in Uttarakhand. Several insecticides have been in use for its management and resistance has been observed against different group of insecticides. Thus, to study the level of infestation and insecticide resistance from soybean fields of hill farmers of Kumaon Himalaya, survey and experiments were conducted.

Methods: Survey was conducted and scale of infestation was determined. Chemical and biochemical assay were conducted to study insecticide resistance. For chemical assay IRAC Method No. 7 using insecticides indoxacarb 14.5% SC, chlorantraniliprole 18.5% Sc, fipronil 5% SC, chlorpyriphos 50%+cypermethrin 5% EC and profenofos 40%+cypermethrin 4% EC and for biochemical assay the specific activity of Carboxylesterase, Acetyl choline esterase, Mono-oxygenase and Glutathione-S-transferase was studied.

Result: The infestation level from different regions showed the presence of *S. litura*, ranging from mild to severe infestation. The resistance ratio of different insecticides used ranged from a low to an extremely high resistance level with LC_{50} ranging from 0.37 µg/ml to 1648.84 µg/ml. The biochemical analysis of carboxylesterase, acetylcholinesterase, monooxygenase and gluthione-s-transferase showed an overproduction of these detoxification enzymes in the resistant population.

Key words: Detoxification enzymes, Insecticide-resistance, *Spodoptera litura*.

INTRODUCTION

Spodoptera litura is one of the most important polyphagous lepidopteran pests in agriculture and belongs to the Noctuidae family (Wang *et al*., 2018; EFSA PLH Panel, 2018). This pest has enormous economic importance for field and horticultural crops (Murthy *et al*., 2006) and is widespread throughout tropical and temperate Asia (Kranz *et al*., 1977). Severe outbreaks of *S. litura* have been reported in Kota (Rajasthan) and Marathwada and Vidarbha (Maharashtra) in India, causing total damage of about US$ 4.5 crores and US$22.5 crores, respectively (CROPSAP, 2012). Soybean, which is an important crop of India and the country ranks fifth in world soybean production (SOPA, 2020), suffered about 24.7% damage from *S. litura* (Higuchi *et al*., 1991). Uttarakhand is a Himalayan state located in the north-western part of India (Sati, 2020). *S. litura* is one of the major defoliator moths that occur in different regions of Uttarakhand and cause significant crop losses. Singh and Sachan (1992) reported that defoliating moths play an important role at the pod stage and thereafter in soybean. Several insecticides were recommended for use against this pest, which exerted high selection pressure on populations of *S. litura*. This selection pressure was responsible for the development of resistance of *S. litura* to many insecticides (Shi *et al*., 2019). In the 2017 IRAC Newsletter, it was documented that *S. litura* was ranked 7th among top 20 arthropods in terms of number of resistance cases. Resistance to insecticides is a complex process and may result mainly from four mechanisms: i) increased metabolism to non-toxic products, ii) decreased sensitivity of the target site, iii) decreased rate of penetration of the insecticide, iv) increased rate of excretion of the insecticide (Sakine, 2012). The simplest and most conventional detection of resistance is dose-mortality experiments conducted in the laboratory under controlled conditions (Brown, 1976). Another approach is the biochemical mechanism, which, when used in addition to basic toxicity studies, gives the best results. Resistance to insecticides has been found to be essentially due to two mechanisms, *i.e*., enhanced enzymatic detoxification of an insecticide or reduced sensitivity of a

[1]Government Beekeeping Centre, Jeolikote-263 127, Nainital, Uttarakhand, India.
[2]Department of Entomology, College of Agriculture, G.B. Pant University of Agriculture and Technology, Pantnagar-263 145, Udham Singh Nagar, Uttarakhand, India.

Corresponding Author: Rashmi Joshi, Government Beekeeping Centre, Jeolikote-263 127, Nainital, Uttarakhand, India.
Email: rashmipandey.hdw@gmail.com

How to cite this article: Joshi, R., Gaur, N. and Mathpal, S. (2023). Insecticide Resistance Development and Detoxification Enzyme Activities of *Spodoptera litura* (Fabricius) in Soybean from Kumaon Himalayas. Legume Research. 46(7): 927-933. doi: 10.18805/LR-5101.

Submitted: 23-01-2023 **Accepted:** 28-03-2023 **Online**: 30-05-2023

target enzyme to inhibition by the insecticide; out of the four mechanisms discussed (Brown, 1987). The farmers from locations surveyed were using insecticides such as Profenofos 40%+Cypermethrin 4% EC, Indoxacarb 14.5% SC, Chlorantraniliprole 18.5% SC, Carbofuran 3G, Chlorpyriphos 50%+ Cypermethrin 5% EC, Fipronil 3.5%+Chlorpyriphos 35% *etc*. which were provided by government agencies involved in agricultural work as per information provided by farmers themselves. Therefore, the present study investigated the resistance level of five different insecticides and five *S. litura* field populations collected during 2019 and 2020 from soybean fields of Kumaon, Himalayas. The results of this study can help in developing effective and timely management practises.

MATERIALS AND METHODS

Field populations of *S. litura* egg masses and larvae in soybean fields were collected from four hill districts, namely Ramgarh (Nainital district), Bhujan (Almora district), Kameri, Bageshwar and Ganai (Pithoragarh district) in Kumaon, Uttarakhand in 2019 and 2020. Other crops observed during survey in farmers field were Soybean, Cabbage, Cauliflower, Colocacia, Capsicum and French bean. Larvae were reared on artificial diet (Ballal, 2003) and adults were fed on 10% honey solution under laboratory conditions (27±1°C and 65±5% relative humidity) with a photoperiod of 16:8 hours light: dark. Adults were kept in jars with butter paper attached for oviposition by females and reared to F1 generation. A susceptible population of *S. litura* was obtained from NBAIR, Bangaluru, for comparative studies. Based on the survey information on insecticides used by farmers in the selected regions, five insecticides were used in the study, namely indoxacarb 14.5% SC, chlorantraniliprole 18.5% Sc, fipronil 5% SC, chlorpyriphos 50%+cypermethrin 5% EC and profenofos 40%+cypermethrin 4% EC. The bioassay method of IRAC Method No. 7 (IRAC, 2010) was performed on third instar larvae of *S. litura* in Pulse Entomology Lab, Deptt of Entomology, College of Agriculture, GBPUAT, Pantnagar. The insecticide solution was prepared from a 1% stock solution by serial dilution. Fresh castor leaves of uniform size *i.e*., 5 × 5 cm were taken and dipped in the insecticide solution for 10 seconds. Excess liquid was drained off and the leaves were then air dried for half an hour. Ten third instar larvae from the F1 progeny (after being starved for 6 hours) were transferred to each petri dish and mortality was monitored at 24, 48 and 72 hours post exposure. Tests were conducted at a controlled temperature of 27± 1°C, 65±5% relative humidity and a 16:8 L:D photoperiod. To investigate the potential of the detoxification enzymes, biochemical analysis was performed to test the total protein and estimate the activity of Carboxylesterase (CarE) (Devonshire,1977 and Van, 1962); Acetyl choline esterase (AChE) (Kranthi, 2005); Mono-oxygenase (P450) (Kranthi, 2005) and Glutathione-S-transferase (GST) (Van,1962 and Booth *et al*., 1961). Polo Suite Leora Software LLC was used to estimate the LC_{50} values of the chemical bioassays and the specific activities of the biochemical studies were analysed using analysis of variance (ANOVA).

RESULTS AND DISCUSSION

The survey to study the distribution and collection of *S. litura* was conducted in 2019 and 2020 in different districts of Kumaon region of Uttarakhand (Table 1, Fig 1). Visual estimation was used to determine the infestation level.




Fig 1: Map of Uttarakhand and districts surveyed shown in shaded colour *viz*., Pithoragarh, Bageshwar, Almora and Nainital (on left side) and image from farmer filed surveyed Ramgarh, District Nainital (on right side).

Table 1: Geographical dimensions and meteorological data of areas from where *S. litura* population was collected for study.

Name of place	Latitude	Longitude	Altitude [m above mean sea level (msl)]	Average temperature °C		RH (%)		Scale of infestation of *S. litura*
				Previous month	Collection month	Previous month	Collection month	
Ramgarh (Nainital district)	29.4347N	79.5521E	1518	30.93	27.38	38.25	75.70	4
Bhujan (Almora district)	29.51N	79.4776E	1604	24.76	23.51	75.48	84.93	4
Kameri (Bageshwar district)	29.8514N	79.7996E	1004	22.42	20.76	85.96	87.16	3
Ganai (Pithoragarh district)	26.6629N	80.075E	1760	19.43	17.90	85.81	87.85	3

The scales for determining infestation were given by Vennila *et al*. (2010) and used by Singh and Gandhi (2012) in their study on *S. littoralis* on cotton, cabbage, radish and other insect pests. Four scales were given for infestation: no insect/scares appearance (scale 0); scattered appearance of few (scale 1); severe incidence on only one branch (scale 2); severe incidence on more than one branch (scale 3) and complete severe incidence (scale 4). The study found that severe to complete severe incidence of the pest was observed in the regions studied. The *S. litura* samples were collected from different locations and brought to the laboratory for further rearing of the F1 generation, where bioassay tests were carried out (Table 2). Insecticide resistance results according to insecticide used and locations visited were classified according to Shen *et al*., (1991) insecticide resistance level classification, *i.e*., RR <3.0 (susceptible), 3.0<RR<5.0 (Decreased Susceptibility); 5.0<RR<10.0 (Low level of resistance); 10.0<RR<40.0 (Moderate level of resistance); 40.0<RR<160.0 (High level of resistance) and RR>160.0 (Extremely high level of resistance).

S. litura populations tested with indoxacarb 14.5% SC were found to be susceptible for Bhujan and Pithoragarh populations; low level of resistance was observed for Ramgarh; moderate level for Bageshwar. Wang *et al*. (2018) also reported low to moderate level of resistance between 2-31-fold of *S. litura* population in Sichuan, China, to indoxacarb 14.5% SC from 2014-2016. Chlorantraniliprole 18.5% SC showed moderate to high resistance at almost all sites. Muthuswamy *et al*. (2014) reported a resistance ratio of 80.07 with respect to the susceptible NBAIR population, which is in insecticide resistance class 4. The bioassay results showed that Fipronil 5% SC showed resistance at almost all sites, from low to very high levels of resistance. Ahmad and Mehmood (2015) found that Pakistani populations developed moderate resistance every 8 years, increasing from 5.5-5.6-fold in 1998 to 28-35-fold in 2006. Ahmad *et al*. (2008) also reported that 22-fold resistance was observed to Fipronil 5% SC and this high level of resistance was explained by a multiple resistance mechanism. For combination insecticides *i.e*., profenofos 40%+cypermethrin 4% EC and chlorpyriphos 50%+ cypermethrin 5% EC, almost similar results were observed with both insecticides at all locations except Bhujan and the possible reason could be the selection pressure of profenofos 40%+cypermethrin 4% EC since the farmers used only this insecticide in their fields. It was also observed that the population of Ramgarh showed sensitivity to the combination of insecticides, which may have been due to the strict control and regulations of some private companies with which the farmers collaborate for organic crop production in the area as mentioned by farmers in the area and due to the low relative humidity (RH) of the sampling site, as RH is an important factor determining the life span of *S. litura* (Kumar *et al*., 2013; Khan and Talukder, 2017). Both the profenofos 40%+cypermethrin 4% EC and chlorpyriphos 50%+cypermethrin 5% EC have an OP to pyrethroid ratio of 1:10 and have good resistance, which Ahmadi (2009), El-Guindy *et al*. (1983), Goebel and Jacquemard (1990) and Forrester *et al*. (1993) reported that profenofos and chlorpyrifos antagonize cypermethrin in a ratio of 1:10.

Populations from different districts of Uttarakhand were subjected to biochemical analysis to identify their levels of detoxification enzymes compared to susceptible populations from NBAIR, Bangaluru (Table 3 and 4). Almora Bhujaan population showed the highest specific activity for both acetylcholinesterase enzyme (6.770±0.140 nmol/min/ml enzyme) and monooxygenase P450 enzymes (2.780±0.630 nmol/min/ml enzyme) and resistance ratio 8 .63 and 111.20 were found in the reference laboratory population, respectively. For carboxylesterase enzyme, the highest specific activity was observed in the population of Kameri, Bageshwar, *i.e*., the formation of 0.505±0.035 µmoles 1-naphthol/min/mg protein and RR (resistance ratio) of 33.67; For glutathione-S-transferase enzyme, the highest specific activity was observed in Ganai, Pithoragarh population *i.e*. 1.910±0.050 µmoles /min/mg protein with a RR of 23.87.

This study was conducted to investigate the level of infection and resistance to five different pesticides preferred by farmers in the region in four different areas of the Kumaon hills, Uttarakhand. The level of *S. litura* infestation was found to be on a scale of 3 and 4, meaning that it ranged from severe presence in more than one branch to full severe presence. According to Chattopadhyay *et al*., (2019) outbreak of *S. litura* was observed at temperature between 21-27°C and RH above 90%. The infestation scale was related with temperature, humidity and availability of abundant host since the favourable conditions generated due to these factors are important for the growth and development of *S. litura* (Joshi *et al*., 2022; Fand *et al*., 2015). The data of previous and collection month showed that Ramgarh and Bhujaan had a constant favourable climatic condition in addition to abundance of host. Also, the chemical bioassay studies showed that the population is resistant against some of the insecticides used implying better survival of insect population in region, in addition to favourable environmental factors. It was observed in the studies that the resistance in Fipronil 5% SC and Chlorantraniliprole 18.5% SC could not be directly correlated with activity of any detoxification enzymes. Arain *et al*. (2018) reported that detoxification of Fipronil 5% SC is not strongly related to the activity of these detoxification enzymes, as well as Su *et al*. (2012) mentioned similar results for 18.5% SC resistance to chlorantranilprole and suggested that another mechanism may be responsible for detoxification of this insecticide. The level of resistance to the combination insecticides can be related to high activity of esterase enzymes *i.e*. AChE and CarE, but no such direct relationship with activity was observed for GST and P450 enzymes, also shown by Muthusamy *et al*. (2011) in their studies on insecticide detoxification mechanisms in *S. litura*.

Table 2: Dosage mortality showing susceptibility of *S. litura* against different insecticides.

Insecticides	Population	LC_{50} (95%CI)	Chi square	Slope±SE	Resistance ratio
Indoxacarb	Ramgarh, Nainital	2.48 (1.31-8.12)	0.499	Y=0.48+1.23x	7.09 (Low level of resistance)
14.5%SC	Bhujaan, Almora	0.37 (0.16-0.66)	2.651	Y=0.61+1.44x	1.06 (Susceptible)
	Kameri, Bageshwar	9.28 (5.32-17.94)	0.138	Y=1.39+1.44x	26.51(Moderate level of resistance)
	Ganai, Pithoragarh	0.79 (0.47-1.32)	1.157	Y=0.25+1.75x	2.26 (Susceptible)
	Control, Lab population	0.35 (0.14-0.64)	0.435	Y=0.60+1.33x	
Chlorantraniliprole	Ramgarh, Nainital	12.51(5.48-25.98)	0.238	Y=1.22+1.11x	12.64 (Moderate level of resistance)
18.5%SC	Bhujaan, Almora	146.39 (69.09-621.18)	2.05	Y=2.1+0.98x	147.87 (High level of resistance)
	Kameri, Bageshwar	21.79 (12.15-44.01)	0.307	Y=1.77+1.34x	22.01 (Moderate level of resistance)
	Ganai, Pithoragarh	17.55 (7.95-42.76)	0.303	Y=1.28+1.03x	17.73 (Moderate level of resistance)
	Control, Lab Population	0.99 (0.62-1.64)	0.342	Y=0.05+1.84x	-
Fipronil 5% SC	Ramgarh, Nainital	1648.84 (868.62-5010.72)	0.126	Y= 3.8+1.8x	355.35 (Extremely high level of resistance)
	Bhujaan, Almora	36.34 (21.54-61.26)	0.54	Y=2.55+1.64x	7.83 (Low level of resistance)
	Kameri, Bageshwar	55.73 (31.94-107.68)	0.138	Y= 2.5+1.44x	12.01 (Moderate level of resistance)
	Ganai, Pithoragarh	74.84 (44.29-146.09)	1.347	Y=2.94+1.55x	16.13 (Moderate level of resistance)
	Control, Lab Population	4.64 (2.45-8.78)	0.391	Y=0.89+1.37x	
Chlorpyriphos 50% +	Ramgarh, Nainital	17.19 (8.55-30.77)	0.517	Y=1.7+1.37x	1.81 (Susceptible)
Cypermethrin 5%EC	Bhujaan, Almora	18.99 (10.23-33.33)	0.457	Y=1.91+1.51x	2.00 (Susceptible)
	Kameri, Bageshwar	55.32 (31.69-123.62)	1.634	Y=2.36+1.37x	5.82 (Low level of resistance)
	Ganai, Pithoragarh	115.12 (63.45-215.82)	0.259	Y=2.93+1.41x	12.12 (Moderate level of resistance)
	Control, Lab Population	9.50 (5.48-18.27)	0.075	Y=1.41+1.45x	-
Profenofos 40% +	Ramgarh, Nainital	5.20 (3.07-9.46)	0.087	Y=1.09+1.54x	1.21 (Susceptible)
Cypermethrin 4% EC	Bhujaan, Almora	96.04 (51.75-167.62)	0.142	Y=2.89+1.46x	22.28 (Moderate level of resistance)
	Kameri, Bageshwar	206.25 (119.88-400.93)	0.267	Y=3.52+1.52x	47.85 (High level of resistance)
	Ganai, Pithoragarh	57.99 (21.77-109.45)	0.810	Y=2.26+1.28x	13.45 (Low level of resistance)
	Control, Lab Population	4.31(2.32-8.29)	0.440	Y=0.83+1.31x	-

Table 3: Specific activity of AChE and CarE for *S. litura* collected from different locations.

Places	AChE (nmoles/min/ml of enzyme)		CarE (µmoles of 1-naphthol formed/min/mg of protein)	
	Specific activity	Ratio w.r.t susceptible population	Specific activity	Ratio w.r.t susceptible population
Bhujaan, Almora	6.770±0.140	84.63	0.070±0.010	4.67
Ramgarh, Nainital	1.195±0.015	14.94	0.175±0.015	11.67
Ganai, Pithoragarh	1.790±0.100	22.38	0.265±0.035	17.67
Kameri, Bageshwar	0.985±0.025	12.31	0.505±0.035	33.67
Control, Lab population	0.080±0.010	1	0.015±0.005	1
CD	0.291		0.08	
P value	<0.05		<0.05	

Table 4: Specific activity of GST and P450 for *S. litura* collected from different locations.

Places	GST (µmoles/min/mg of protein)		P450 (nmoles/min/ml of enzyme)	
	Specific activity	Ratio w.r.t susceptible population	Specific activity	Ratio w.r.t susceptible population
Bhujaan, Almora	1.045±0.065	13.06	2.780±0.630	111.20
Ramgarh, Nainital	0.375±0.075	4.69	0.285±0.035	11.40
Ganai, Pithoragarh	1.910±0.050	23.87	0.260±0.030	10.40
Kameri, Bageshwar	1.475±0.055	18.44	0.070±0.010	2.80
Control, Lab population	0.080±0.010	1.00	0.025±0.005	1.00
CD	0.207		1.050	
P value	<0.05		<0.05	

Similar results were obtained comparing Indoxacarb 14.5%SC resistance with detoxification enzyme activity, as no correlation was observed between enzyme activity and insecticide resistance, suggesting a possible role for other detoxification mechanisms (Shi *et al*., 2019). Enzymatic activity indicated that overproduction of detoxification enzymes may be one of the main factors contributing to insecticide resistance in the studied *S. litura* populations (Despres *et al*., 2007).

CONCLUSION

Soybean is an important crop in Uttarakhand and is important pulse in the diet of people from hills of Kumoan. Development of insecticide resistance in *S. litura* which is one of the major insect-pests of soybean has already been reported from different parts of world. This study found that insecticide resistance could become a major concern for hill farmers in Uttarakhand, India. Multiple mechanisms are responsible for the development of insecticide resistance in insects, which requires further investigation, making it a promising area of research for developing integrated insect control strategies.

ACKNOWLEDGEMENT

We are thankful to G.B. Pant University of Agriculture and Technology, Pantnagar for providing all the support to conduct survey and lab research work.

Conflict of interest: None.

REFERENCES

Ahmadi, M. (2009). Observed potentiation between pyrethroid and organophosphorus insecticides for the management of *Spodoptera litura* (Lepidoptera: Noctuidae). Crop Protection. 28: 264-268.

Ahmad, M. and Mehmood, R. (2015). Monitoring of resistance to new chemistry insecticides in *Spodoptera litura* (Lepidoptera: Noctuidae) in Pakistan. Journal of Economic Entomology. 108(3): 1279-1288.

Ahmad, M., Sayyed, A.H., Saleem, M.A. and Mushtaq, A. (2008). Evidence for field evolved resistance to newer insecticides in *Spodoptera litura* (Lepidoptera: Noctuidae) from Pakistan. Crop Protection. 27: 1367-1372.

Arain, M.S., Shakeel, M., Elzaki, M.E.A. *et al*., (2018). Association of detoxification enzymes with butene-fipronil in larvae and adults of *Drosophila melanogaster*. Environmental Science and Pollution Research. 25: 10006-10013.

Ballal, C.R. (2003). Mass Production of *Helicoverpa armigera* and *Spodoptera litura* and their Parasitoids. In: Biological Control of Crop Pests in India. [Kumari, S. (Eds.)], First Edition. Ludhiana, India, Kalyani Publisher. pp. 190-197.

Booth, J., Boyland, E. and Sims, P. (1961). An enzyme from rat liver catalysing conjugations with glutathione. Biochemical Journal. 79: 516-524.

Brown, A.W.A. (1976). How have entomologists dealt with resistance? American Phytopathological Society. 3: 67-74.

Brown, M.T. (1987). Improved detection of insecticide resistance through conventional and molecular techniques. Annual Review of Entomology. 32: 145-162.

Chattopadhyay, N., Balasubramaniam, R., Attri, S.D., Ray, K., Gracy, J., Khedikar, S. and Karmakar, C. (2019). Forewarning of incidence of *Spodoptera litura* (Tobacco caterpillar) in soybean and cotton using statistical and synoptic approach. Journal of Agricultural Meteorology. 21: 68-75.

CROPSAP, (2012). Crop Pest Surveillance and Advisory Project in Maharashtra (2011-12), Sponsored by Commissionerate of Agriculture, Government of Maharashtra under Rashtriya Krishi Vikas Yojana. http://www.ncipm.org.in/cropsap2013/CROPSAP.PDF/Project.Background.pdf (accessed 22.04.2020).

Despres, L., David, J.L. and Gallet, C. (2007). The evolutionary ecology of insect resistance to plant chemicals. Trends in Ecology and Evolution. 22(6): 298-307.

Devonshire, A.L. (1977). The properties of a carboxylesterase from the peach potato aphid, *Myzus persicae* (Sulz.) and its role in conferring insecticide resistance. Biochemical Journal. 167: 675-683.

EFSA PLH Panel (EFSA Panel on Plant Health), Jeger, M., Bragard, C., Caffier, D. *et al*. (2018). Guidance on quantitative pest risk assessment. EFSA Journal. 16(8): 535. https://doi.org/10.2903/j.efsa.2018.5350 (accessed 22.04.2020).

El-Guindy, M.A., Rehman, A., El-Refai, M. and Abdel-Sattar, M.M. (1983). The joint action of mixtures of insecticides, or of insect growth regulators and insecticides, on susceptible and diflubenzuron-resistant strains of *Spodoptera littoralis* Boisd. Journal of Pesticide Science. 14: 246-252.

Fand, B.B., Sul, T.N., Bal, K.S. and Minhas, P.S. (2015). Temperature Impacts the Development and Survival of Common Cutworm (*Spodoptera litura*): Simulation and Visualization of Potential Population Growth in India under Warmer Temperatures through Life Cycle Modelling and Spatial Mapping. https://Journals.plos.org/plosone/article?id=10.1371/journal.pone.0124682 (accessed 09.04.2019).

Forrester, N.W., Cahill, M., Bird, L.J. and Layland, J.K. (1993). Management of pyrethroid and endosulfan resistance in *Helicoverpa armigera* (Lepidoptera: Noctuidae) in Australia. Bulletin of Entomological Research. 1: 1-132.

Goebel, R. and Jacquemard, P. (1990). Evaluation of the sensitivity of the cotton boll pest *Heliothis armigera* Hbn. to cypermethrin-chlorpyrifos and cypermethrin-methylparathion mixtures: Study of possible interactions between these insecticides. Coton et Fibres Tropicales. 45: 137-143.

Higuchi, H. (1991). Analysis of damage to soybeans infested by the common cutworm, *Spodoptera litura* Fabricius (Lepidoptera: Noctuidae). Effects of infestation at different growth stages of the soybean plant. Jpanese Journal of Applied Entomology and Zoology. 35(2): 131-135.

IRAC Insecticide Resistance Action Committee, (2010). Insecticide Resistance Action Committee, Method No: 007: Leaf Eating Lepidoptera and Coleoptera. http://www.irac-online.org/wpcontent/uploads/2009/09/Method007v3june09.pdf (accessed 09.04.2019).

Joshi, R., Gaur, N. and Mathpan S. (2022). Distribution of *Spodoptera litura* (F.) in Uttarakhand. Indian Journal of Entomology. e22485, accessed on 13.03.2023 DoI. No.: 10.55446/IJE.2022.796.

Khan, M.M.H. and Talukder, S. (2017). Influence of weather factors on the abundance and population dynamics of *Spodoptera litura* F. and *Pieris brassicae* L. on cabbage. SAARC Journal of Agriculture. 15(1): 13-21.

Kranthi, K.R. (2005). Insecticide Resistance-Monitoring, Mechanisms and Management Manual, Central Institute for Cotton Research. Nagpur, 49-87 pp.

Kranz, J., Schumutterer, H. and Koch, W. (1977). Diseases Pests and Weeds in Tropical Crops. Berlin and Hamburg, Germany, Verlag Paul Parley.

Kumar, S., Jaspreet, K.S., Jason, C.H., Jagdev, S.K. and Manjit, S.M. (2013). Effects of temperature and relative humidity on the life table of *Phenacoccus solenopsis* tinsley (Hemiptera: Pseudococcidae) on cotton. Florida Entomologist. 96(1): 19-28.

Murthy, M.S., Thippaiah, M. and Kitturmmath, M.S. (2006). Effect of neem formulations on larvae of tobacco cutworm, *Spodoptera litura* (Fab.). Insect Environment. 12: 84-85.

Muthusamy, M., Shivakumar, S., Karthi, K. and Ramkumar, R. (2011). Pesticide detoxifying mechanism in field population of *Spodoptera litura* (Lepidoptera: Noctuidae) from South India. Egyptian Academic Journal of Biological Sciences, F. Toxicology and Pest Control. 3(1): 51-57.

Muthuswamy, R.M., Vishnupriya, M.S. and Shivakumar. (2014). Biochemical mechanism of chlorantraniliprole resistance in *Spodoptera litura* (Fab) (Lepidoptera: Noctuidae). Journal of Asia-Pacific Entomology. 17: 865-869.

Sakine, U.K. (2012). Insecticide Resistance. In: Insecticides-Advances in Integrated Pest Management. [Perveen, F. (Ed.)], InTech. http://www.intechopen.com/books/insecticides-advances-in-integrated-pest management/insecticidere sistance (accessed 12.01.2020).

Sati, V.P. (2020). Geography and Geology. In: Himalaya on the Threshold of Change. [Sati, V.P. (Ed.)], Advances in Global Change Research. Switzerland, Springer, Cham. Vol 66, pp. 3-19.

Shen, J.L., Tan, J.G., Xiao, B., Tan, F.J. and You, Z.P. (1991). Monitoring and forecasting of *Helicoverpa armigera* (Lepidoptera: Noctuidae) resistance to pyrethroid pesticides in China. Chinese Journal of Applied Entomology. 28: 337-341.

Shi, L., Shi, Y., Zhang, Y. *et al*., (2019). A systemic study of indoxacarb resistance in *Spodoptera litura* revealed complex expression profiles and regulatory mechanism. Scientific Reports. 9: 14997. https://doi.org/10.1038/s41598-019-51234-5. (accessed 24.09.2020).

Singh, A. and Gandhi, S. (2012). Agricultural Insect Pest: Occurrence and infestation level in agricultural fields of Vadodara, Gujarat. International Journal of Scientific and Research Publications. 2(4): 1-5.

Singh, K.N. and Sachan, G.C. (1992). Assessment of yield loss due to insect pests at different growth stages of groundnut in Pantnagar, Uttar Pradesh, India. Crop Protection. 11: 414-418.

SOPA, (2020). World Soybean Production by the soybean processors association of India. http://www.sopa.org/statistics/world-soybean-production/ (accessed 12.12.2021).

Su, J., Lai, T. and Li, J. (2012). Susceptibility of field populations of *Spodoptera litura* (Fabricius) (Lepidoptera: Noctuidae) in China to chlorantraniliprole and the activities of detoxification enzymes. Crop Protection. 42: 217-222.

Van, A.K. (1962). A study of housefly esterases by means of a sensitive calorimetric method. Journal of Insect Physiology. 8: 401-416.

Vennila, S., Ramamurthy, V.V., Deshmukh, A., Pinjarkar, D.B., Agarwal, M., Pagar, P.C., Prasad, Y.G., Prabhakar, M., Kranthi, K.R. and Bambawale, O.M. (2010). A Treatise on Mealybugs of Central Indian Cotton Production System. National Centre for Integrated Pest Management, New Delhi, Tec. Bull. 24: 1-39.

Wang, X., Huang, Q., Hao, Q., Ran, S., Wu, Y., Cui, P., Yang, J., Jiang, C. and Yang, Q. (2018). Insecticide resistance and enhanced cytochrome P450 monooxygenase activity in field populations of *Spodoptera litura* from Sichuan, China. Crop Protection. 106: 110-116.