RESEARCH ARTICLE Legume Research- An International Journal, Volume 48 Issue 1: 143-148 (January 2025)

Hybrid Pigeonpea (ICPH 2740) Nodular Activity and Biological Nitrogen Fixation as Influenced by Agronomic Practices

Bathula Venkatesh[1], M. Malla Reddy[1], G.L. Sawargaonkar[2], Ch. Sarada[3], B. Padmaja[1], M. Yakadri[1], Nallagatla Vinod Kumar[1]

10.18805/LR-5167

ABSTRACT

Background: To explore the impact of agronomic practices on nodulation, nitrogenase activity and biological nitrogen fixation in hybrid pigeonpea, a field investigation was under taken at BW5 block ICRISAT development center (IDC), International Crops Research Institute for Semi-Arid Tropics (ICRISAT), Patancheru, Hyderabad, during *kharif,* 2021 and 2022.

Methods: The seedlings were raised in portrays in advance to onset of monsoon during last week of May (28th May in 2021 *kharif* and 30th May in 2022 *kharif*) in both the years by filling portrays with cocopeat and vermicompost in 1:1 ratio. Transplanting of 24 days and 25 days old seedlings were done on 20th June in 2021 *kharif* and 23th June in 2022 *kharif*, respectively, on the same day of dibbling in the main field at 100 cm × 100 cm ,120 cm × 120 cm square geometry and 150 cm × 60 cm. Nutrient management treatments were imposed one week after sowing.

Result: Among the planting methods higher nodule number and nitrogenase activity with transplanting compared to dibbling. 120 cm × 120 cm found higher nodule number plant^{-1} and nitrogenase activity but biological nitrogen fixation was higher with 100 cm × 100 cm ha^{-1} basis. With respect on nutrient management practices, control performed well in terms of higher nitrogenase activity and biological nitrogen fixation and on par with integrated approach of 100% soil test based NPK + vermicompost + PSB + seed treatment with *Rhizobium*. So, with transplanting method of establishment, 100 × 100 cm square plant geometry and 100% soil test based NPK + vermicompost + PSB + seed treatment with *Rhizobium* was recommended agronomic practices for hybrid pigeonpea.

Key words: Acetylene reduction assay, Biological nitrogen fixation, Hybrid pigeonpea, Nitrogenase activity, Nutrient management.

INTRODUCTION

Pigeonpea (*Cajanus cajan* L.) is the second most important pulse crop after chickpea in India. Pigeonpea is grown in an area of about 4.5 m ha with production of 3.3 MT in the country. The national productivity of pigeonpea is only 729 kg ha^{-1} give comparison to world productivity (Directorate of Economics and Statistics, GOI 2022). In Telangana, it is cultivated in an area of 2.96 lakh ha with production of 1.91 lakh tones and productivity of 647 kg ha^{-1} (Directorate of Economics and Statistics, GOI 2022).

In India more than 90 per cent of pigeonpea area is under rainfed conditions (Tiwari and Namrata, 2020). Now a day's due monsoon aberrations farmers unable to sow the pigeonpea in correct window as a result the length of vegetative period reduced (Susithra *et al*., 2019) so, to avoid this the concept of transplanting is came into picture. But with normal conventional method of crop geometry like 150 × 30 cm leads to severe interplant competition, improper expression and higher flower drops due to low ventilation (Pradeep *et al*., 2018). Transplanting hybrid pigeonpea under square system the distance between rows and plants is same which facilitates good branching, pod and seed set when compared to varieties. Due to higher branching ability of hybrid under transplanting conditions, more yields were obtained compared to normal dibbling (Ramanjaneyulu *et al*., 2017; Mallikarjun *et al*., 2015).

Biological nitrogen fixation is a process of great environmental and economic importance, performed by *Rhizobium* that will reduce atmospheric nitrogen (N_2) to ammonia (NH_3) and when in symbiotic association with plants, they provide NH_3, which is readily convertible to other forms, such as amides and ureides (Rufini *et al*., 2016). Seed treatment with rhizobium before sowing in pigeonpea is suitable alternative for increasing yields. ICPH 2740 is the first commercial pigeonpea hybrid released in Telangana state as Mannemkonda kandi in 2015 and yield potential of

[1]Department of Agronomy, Professor Jayashankar Telangana State Agricultural University, Rajendranagar, Hyderabad-500 030, Telangana, India.
[2]International Crops Research Institute for the Semi-Arid Tropics, Patancheru, Hyderabad-502 324, Telangana, India.
[3]ICAR-Indian Institute of Oilseeds Research, Rajendranagar, Hyderabad-500 030, Telangana, India.

Corresponding Author: Bathula Venkatesh, Department of Agronomy, Professor Jayashankar Telangana State Agricultural University, Rajendranagar, Hyderabad-500 030, Telangana, India. Email: vihaanvarma179@gmail.com

How to cite this article: Venkatesh, B., Reddy, M.M., Sawargaonkar, G.L., Sarada, C., Padmaja, B., Yakadri, M. and Kumar, N.V. (2025). Hybrid Pigeonpea (ICPH 2740) Nodular Activity and Biological Nitrogen Fixation as Influenced by Agronomic Practices. Legume Research. 48(1): 143-148. doi: 10.18805/LR-5167.

Submitted: 03-05-2023 **Accepted:** 16-09-2023 **Online:** 30-09-2023

this hybrid is ranges from 2407-3652 kg ha^{-1} (Saxena *et al*., 2016). Among the different agronomic practices, suitable planting method, planting geometry and optimum nutrition is essential for enhancing productivity in hybrid pigeonpea.

Materials and Methods

Experimental site

A field study entitled with "Hybrid pigeonpea (ICPH 2740) nodular activity and biological nitrogen fixation as influenced by Agronomic practices" was conducted at BW5 block ICRISAT development center (IDC). International Crops Research Institute for Semi-Arid Tropics (ICRISAT), Patancheru, Hyderabad, during *kharif,* 2021 and 2022. The rainfall received during *kharif*, 2021 and 2022 experimental periods was 998.4 mm in 55 rainy days and 1000 mm in 53.1 rainy days respectively. Experimental soil is moderately alkaline in reaction with pH of 8.1 and EC of 1.94 dSm^{-1}. The N, P_2O_5 and K_2O and organic carbon content of soil was, low (231.1 kg ha^{-1}), medium (29.8 kg ha^{-1}) and high (350.5 kg ha^{-1}) and low (0.42%), respectively.

Treatment and experimental design

The experiment was laid out in split-split plot design with two planting methods in main plots *i.e*., M_1: Dibbling, M_2: Transplanting, three crop geometries in sub-plots *i.e*., S_1: 100 cm × 100 cm (10,000 plants ha^{-1}), S_2: 120 cm × 120 cm (6,944 plants ha^{-1}), S_3: 150 cm × 60 cm (11,111 plants ha^{-1}) (Normal transplanting) and five nutrient management practices in sub-sub plots *i.e*., N_1: Control, N_2:100% soil test-based (STB) NPK (25:37.5:8.5 kg ha^{-1}), N_3: 100% STB NPK + vermicompost + PSB + Seed treatment (ST) with *Rhizobium*, N_4: 150 % STB NPK, N_5: 150% STB NPK + vermicompost + PSB + ST with *Rhizobium.* The pigeonpea hybrid used for this investigation was ICPH 2740. The seedlings for transplanting were raised in protrays and sown on 28th May in 2021 *kharif* and 30th May in 2022 *kharif*, respectively. Transplanting of 24 days and 25 days old seedlings were done on 20th June in 2021 *kharif* and 23th June in 2022 *kharif*, respectively, on the same day of dibbling in the main field. Vermicompost 5 t ha^{-1}, PSB (Phosphorus solubilizing bacteria 5 kg ha^{-1} added to vermicompost one week prior to application and seed treatment with *rhizobium* 500 g ha^{-1} seed 2 hours before sowing. Based on soil nutrient statusthe recommendation was made such that, if nutrient status is low, 25% more dose on recommended dose, if it is high, 25% less than RDF and recommended dose is based on medium nutrient status.

Data collection and analysis methods

The plant were uprooted carefully by digging around the plant by crowbar and the plants were washed under running tap in sieve and the nodules were counted and expressed in (No. $plant^{-1}$) at active nodulation stages *i.e.* 30, 45 and 60 DAS/DAT. Nitrogenase reduces acetylene (C_2H_2) to ethylene (C_2H_4) and the ethylene formed can be very sensitively detected by gas chromatography. In this method detached root with nodules enclosed in air tight glass jar (700 ml) and 10% gas inside replaced by pure acetylene. After 60 minutes incubation at room temperature, 0.5 ml of gas inside is sampled using a hypodermic syringe and concentration of ethylene gas is analysed by gas chromatography in acetylene reduction assay (ARA). The amount of ethylene formed will proportional to nitrogenase activity. ARA is expressed µmol C_2H_4 $plant^{-1}$ hr^{-1}. By using this nitrogenase activity we can calculate biological nitrogen fixation (kg ha^{-1}) by area under curve method (Hardy *et al*., 1968). 3 is the theoretical conversion factor of 3 moles of acetylene reduced/mole of dinitrogen reduced (Peoples *et al*., 2009).

Statistical analysis

Data on nodule number and nodular activity were subjected to analysis of variance procedures as outlined for spit-split plot design (Gomez and Gomez, 1984). Statistically significance was tested by F-value at p=0.05 (5%) level of probability and critical difference was worked out where ever the effect was significant. Treatment differences that were non-significant were denoted as NS.

Results and Discussion

Number of root nodules $plant^{-1}$

The effect of different agronomic practices on nodule number was studied during both years of experimentation (Table 1) and significant variation was found. Among the planting methods at 30 DAS/DAT significantly higher number of root nodule $plant^{-1}$ were recorded with transplanting (16.4) compared dibbling (12.4). Whereas, in planting geometry higher number of nodules $plant^{-1}$ were recorded with 120 × 120 cm (6944 plants ha^{-1}), which was statistically on par with 100 × 100 cm (10,000 plants ha^{-1}) at all the stages of crop and with respect to the nutrient management options, significantly higher number of root nodules were recorded with control at 30, 45 and 60 DAS/DAT and which was on par with 100 % STB NPK + 5 t ha^{-1} + vermicompost enriched with PSB + *Rhizobium* seed treatment. These treatments followed by application of 150% soil test based NPK 5 t ha^{-1} + vermicompost enriched with PSB + *Rhizobium* seed treatment and on par with 100% soil test based NPK. Similar trend was observed during both the years of experimentation. Lower nodule number was observed with 150 % soil test based NPK during both the years at all the stages. As pigeonpea is a leguminous crop it able to fix the atmospheric nitrogen. Wherever, nutrient demand is more, the fixation ability increases to meet the required demand. This could be the reason for the higher number of root nodules formation in control plot and also influenced by integrated nutrient approaches (Mishra and Prasad, 2010). During the both years of experimentation the interaction effect of planting method, plant geometry and nutrient management practices were found to be non-significant with respect to the number of root nodules $plant^{-1}$ at 30, 45 and 60 DAS/DAT. As pigeonpea is deep rooted legume it has ability to draw the fixed phosphorus from deeper layers of

soil by releasing certain root exudates and acids (Piscidic acid) but, the nitrogen requirement should be meet my biological nitrogen fixation (Singh *et al*., 2016). So, the recommended dose of nitrogen is very less, the required nitrogen demand of crop should be met by biological nitrogen fixation (Tigga and Singh, 2019), but this nodulation activity again depends upon the many external and management factors (Kumar and Dart, 1987). For successful nodulation certain agronomic practices at field level *i.e*. addition of rhizobium inoculum through seed treatment may help in the formation of effective nodule Patil and Padmani (2007); Mishra and Prasad (2010) and Rufini *et al*. (2016).

Nodular activity (Nitrogenase enzyme activity µmol C_2H_4 plant^{-1} hr^{-1})

As pigeonpea is leguminous crop nodule function and biological nitrogen fixation depends upon the nitrogenase enzyme activity. Nitrogenase activity was studied through acetylene reduction assay in *kharif* 2021 and 2022 (Table 2) and found that, significantly higher nitrogenase activity was recorded with transplanting at 30 DAT (36.8 µmol C_2H_4 plant^{-1} hr^{-1}) compared dibbling (23.9 µmol C_2H_4 plant^{-1} hr^{-1}). In plant geometry, higher nodule nitrogenase activity plant^{-1} was recorded with 120 × 120 cm (6944 plants ha^{-1}), which was statistically on par with 100 × 100 cm (10,000 plants ha^{-1}). Among the nutrient management practices, significantly higher nitrogenase activity was found with control, which was on par with integrated nutrient management approach (*i.e*. 100% soil test based NPK + vermicompost 5 t ha^{-1} + vermicompost enriched with PSB + *Rhizobium* seed treatment). Lowest nitrogenase activity was found with 150% STB NPK during both the years experimentation. From 30 to 60 DAS/DAT there increase in nodular activity in all the treatments but, from 60 to 75 DAS nitrogenase activity was decreased in all the treatments. Interaction effect of planting methods, plant geometry and nutrient management practices was found to be non-significant. The nodule number is indirect indicators of nitrogenase activity, but the enzyme nitrogenase activity which is actually involved in the biological nitrogen fixation (Bidlack *et al*., 2001) which is forerunner for meeting nitrogen requirement of the plant (Peoples *et al*., 2009; Barber *et al*., 1976).

Biological nitrogen fixation (kg ha^{-1})

In transplanting due to early establishment and vigorous growth of seedlings, at 30 DAT more nitrogen was fixed biologically compared with dibbling (Table 3). With respect to the plant geometry, 150 × 60 cm (11,111plants ha^{-1}) which was found on par with 100 × 100 cm (10, 000 plants ha^{-1}). Significantly lower nitrogen was fixed with 120 × 120 cm (6944 plants ha^{-1}) on hectare basis. Among the nutrient management practices, 100 % soil test based NPK +

Table 1: Number of root nodules plant^{-1} of hybrid pigeonpea as influenced by agronomic practices during *kharif*, season.

Treatment	30 DAS/DAT			45 DAS/DAT			60 DAS/DAT		
	2021	2022	Mean	2021	2022	Mean	2021	2022	Mean
Planting method									
M_1: Dibbling	11.1	13.6	12.4	25.8	25.0	25.4	28.3	32.6	30.5
M_2: Transplanting	14.7	18.1	16.4	36.3	39.7	38.0	23.3	23.3	23.3
SEm±	0.2	0.4	-	0.5	0.5	-	0.2	0.2	-
CD (P=0.05)	1.2	2.5	-	3.0	3.2	-	1.0	1.4	-
Plant geometry									
S_1: 100 × 100 cm	13.0	16.3	14.7	31.4	32.3	32.3	26.5	28.4	27.5
S_2: 120 × 120 cm	13.5	17.2	15.4	33.0	33.7	33.7	27.0	30.0	28.5
S_3: 150 × 60 cm	12.3	14.0	13.2	28.7	29.2	29.2	23.9	25.4	24.7
SEm±	0.2	0.3	-	0.4	0.5	-	0.4	0.5	-
CD (P=0.05)	0.6	1.0	-	1.4	1.5	-	1.2	1.7	-
Nutrient management									
N_1: Control	15.0	17.8	16.4	38.6	39.4	39.9	33.4	34.0	33.7
N_2 :100% STB RDF	12.2	15.3	13.8	28.1	27.8	26.8	22.4	24.8	23.6
N_3: 100% STB RDF + vermicompost + PSB + *Rhizobium*	14.1	17.0	15.6	36.0	36.4	35.9	30.4	33.5	32.0
N_4: 150% STB RDF	9.9	12.9	11.4	23.3	24.0	24.0	18.7	21.1	19.9
N_5: 150% STB RDF + vermicompost + PSB + *Rhizobium*	13.3	16.1	14.7	30.3	31.5	32.5	25.1	26.4	25.8
SEm±	0.5	0.5	-	1.0	0.9	-	1.0	0.7	-
CD (P=0.05)	1.4	1.5	-	2.9	2.5	-	3.1	2.1	-

All 2-way and 3-way Interactions are non-significant.

Table 2: Nitrogenase activity (µmol C_2H_4 $plant^{-1}$ hr^{-1}) of hybrid pigeonpea as influenced by agronomic practices during *kharif*, season.

Treatment	30 DAS/DAT			45 DAS/DAT			60 DAS/DAT			75 DAS/DAT		
	2021	2022	Mean	2021	2022	Mean	2021	2022	Mean	2021	2022	Mean
Planting method												
M_1: Dibbling	24.2	23.6	23.9	38.5	37.0	37.8	40.8	37.8	39.3	22.4	20.1	21.3
M_2: Transplanting	36.7	36.9	36.8	67.0	70.1	68.6	55.9	53.8	54.9	34.1	30.5	32.3
SEm±	0.8	1.0	-	1.2	0.9	-	1.6	2.1	-	1.1	0.9	-
CD (P=0.05)	4.9	6.1	-	7.2	5.5	-	9.7	12.7	-	6.7	5.4	-
Plant geometry												
S_1: 100 × 100 cm	30.4	30.1	30.3	52.7	54.2	53.5	48.0	46.6	47.3	29.1	28.1	28.6
S_2: 120 × 120 cm	31.0	30.5	30.8	53.6	55.8	54.7	49.7	47.1	48.4	32.0	32.4	32.2
S_3: 150 × 60 cm	30.0	29.2	29.6	51.9	50.5	51.2	47.4	43.8	45.6	25.0	24.1	24.6
SEm±	0.4	0.5	-	2.1	1.8	-	1.9	1.4	-	1.2	0.8	-
CD (P=0.05)	NS	NS	-	NS	NS	-	NS	NS	-	7.3	4.9	-
Nutrient management												
N_1: Control	41.1	42.8	42.0	76.0	72.7	74.4	72.9	64.2	68.6	56.1	51.6	53.9
N_2:100% STB RDF	26.3	24.9	25.6	44.2	43.9	44.1	37.6	32.3	35.0	21.4	14.7	18.1
N_3: 100% STB RDF + vermicompost + PSB + *Rhizobium*	37.5	36.7	37.1	66.9	70.4	68.7	62.6	60.5	61.6	40.6	36.4	38.5
N_4: 150% STB RDF	20.9	20.5	20.7	36.4	37.3	36.9	30.9	31.0	31.0	16.2	14.6	15.4
N_5: 150% STB RDF + vermicompost + PSB + *Rhizobium*	26.5	24.8	25.7	40.0	43.3	41.7	37.9	41.1	39.5	18.3	21.7	20.0
SEm±	0.4	0.7	-	2.1	1.6	-	1.9	1.1	-	0.9	1.1	-
CD (P=0.05)	1.3	1.9	-	6.1	4.5	-	5.2	3.4	-	6.1	6.6	-

All 2-way and 3-way Interactions are non-significant.

Table 3: Biological nitrogen fixation (kg ha^{-1}) of hybrid pigeonpea as influenced by agronomic practices during *kharif*, season.

Treatment	30 DAS/DAT			45 DAS/DAT			60 DAS/DAT			75 DAS/DAT		
	2021	2022	Mean	2021	2022	Mean	2021	2022	Mean	2021	2022	Mean
Planting method												
M_1: Dibbling	0.76	0.74	0.75	1.75	1.69	1.72	2.99	2.87	2.93	3.98	3.78	3.88
M_2: Transplanting	1.15	1.16	1.16	2.78	2.84	2.81	4.71	4.79	4.75	6.13	6.11	6.12
SEm±	0.02	0.03	-	0.05	0.03	-	0.10	0.08	-	0.13	0.11	-
CD (P=0.05)	0.14	0.22	-	0.30	0.21	-	0.61	0.54	-	0.81	0.67	-
Plant geometry												
S_1: 100 × 100 cm	1.02	1.01	1.02	2.42	2.43	2.43	4.11	4.12	4.12	5.40	5.38	5.39
S_2: 120 × 120 cm	0.72	0.71	0.72	1.71	1.72	1.72	2.92	2.92	2.92	3.87	3.85	3.86
S_3: 150 × 60 cm	1.12	1.09	1.11	2.65	2.58	2.62	4.50	4.34	4.42	5.85	5.61	5.73
SEm±	0.03	0.04	-	0.06	0.05	-	0.52	0.48	-	0.34	0.26	-
CD (P=0.05)	0.09	0.12	-	0.21	0.18	-	1.70	1.52	-	1.10	0.83	-
Nutrient management												
N_1: Control	1.29	1.34	1.32	3.13	3.16	3.15	5.47	5.31	5.39	7.50	7.13	7.32
N_2:100% STB RDF	0.83	0.78	0.81	1.93	1.86	1.90	3.22	3.06	3.14	4.15	3.80	3.98
N_3: 100% STB RDF + vermicompost + PSB + *Rhizobium*	1.18	1.15	1.17	2.82	2.84	2.83	4.85	4.89	4.87	6.47	6.41	6.44
N_4: 150% STB RDF	0.66	0.64	0.65	1.56	1.55	1.56	2.61	2.63	2.62	3.35	3.34	3.35
N_5: 150% STB RDF + vermicompost + PSB + *Rhizobium*	0.83	0.78	0.81	1.88	1.85	1.87	3.10	3.18	3.14	3.98	4.16	4.07
SEm±	0.04	0.05	-	0.04	0.05	-	0.12	0.09	-	0.41	0.38	-
CD (P=0.05)	0.12	0.14	-	0.11	.0.16	-	0.43	0.27	-	1.15	1.08	-

All 2-way and 3-way Interactions are non-significant.

vermicompost 5 t ha^{-1} + vermicompost enriched with PSB + *Rhizobium* seed treatment was recorded higher nitrogen fixation and was found on par with control. Lower nitrogen fixation was found with 150% soil test based NPK during *kharif* 2021, 2022. The average nitrogen was fixed about 6.12 kg ha^{-1} in transplanting, 5.73 kg ha^{-1} with 150 × 60 cm pant geometry (11,111 plants ha^{-1}) and 7.32 and 6.44 kg ha^{-1} with control and 100% soil test based NPK + vermicompost 5 t ha^{-1} + vermicompost enriched with PSB + *Rhizobium* seed treatment respectively. The higher root proliferation with transplanting, wider plant geometry leads to elevated root exudates secretions (Mark *et al*., 2009; Sekhon *et al*., 2018) might increase the microbial population in rhizosphere and proportionate biological nitrogen fixation (Tilak *et al*., 2006 and Dhaka *et al*., 2020).

Conclusion

Hybrid pigeonpea performed well under transplanting, 100 × 100 cm plant geometry and 100% soil test based NPK + vermicompost 5 t ha^{-1} + vermicompost enriched with PSB + *Rhizobium* seed treatment among nutrient management practices in terms of number of root nodules, nitrogen fixation which results required nutrient demand of the plant can be met for the growth and development of the crop.

Acknowledgement

Authors are very thankful to the technical and non-technical staff of International Crops Research Institute for Semi-arid Tropics, Patancheru, Hyderabad. And also special thanks to Indian Institute of Oil seeds Research for providing bus facility for smooth running of experiment.

Conflict of interest

This article publication is approved by all authors and tacitly or explicitly by the responsible authorities where the work was carried out and that, if accepted, it will not be published elsewhere in the same form, in English or in any other language.

References

Barber, L.E., Tjepkema, J.D., Russell, S.A. and. Evans, H.J. (1976). Acetylene reduction (nitrogen fixation) associated with corn inoculated with Spirillum. Applied and Environmental Microbiology. 32(1): 108-113.

Bidlack, J.E., Rao, S.C. and Demezas, D.H. (2001). Nodulation, nitrogenase activity and dry weight of chickpea and pigeon pea cultivars using different *Bradyrhizobium* strains. Journal of Plant Nutrition. 24(3): 549-560.

Dhaka, A.K., Kumar, S., Singh, B., Singh, K., Kumar, A. and Kumar, N. (2020). Nitrogen use efficiency, economic return and yield performance of Pigeonpea [*Cajanus cajan* (L.) Millsp.] as influenced by nipping and fertility levels. Legume Research-An International Journal. 43(1): 105-110. DOI:10.18805/LR-4062.

Directorate of Economics and Statistics. Government of India. (2022). https://www.ecostat.india.gov.in.

Gomez, K.A. and Gomez, A.A. (1984). Statistical Procedure for Agriculture Research. 2nd Edition. John Willey and Sons, New York. 680.

Hardy, R.W., Holsten, R.D., Jackson, E.K. and Burns, R. (1968). The acetylene-ethylene assay for N2 fixation: Laboratory and field evaluation. Plant physiology. 43(8): 1185-1207.

Kumar, R.J.V.D. and Dart, P.J. (1987). Nodulation, nitrogen fixation and nitrogen uptake in pigeonpea [*Cajanus cajan* (L.) Millsp] of different maturity groups. Plant and Soil. 99: 255-266.

Mark, B., Peoples, Murray, J., Unkovich, D. and Herridge, F. (2009). Measuring symbiotic nitrogen fixation by legumes. Agronomy Monograph. 52. 126-170.

Mishra, A. and Prasad, K. (2010). Synergism among *Rhizobium*, phosphate solubilizing bacteria and plant growth promoting *Rhizobacteria* in field pea. International Journal of Applied Agriculture Research. 5(1): 25-30.

Patil, A.B. and Padmani, D.R. (2007). Effect of INM practices on yield, quality and economics of Pigeonpea [*Cajanus cajan* (L.) Millsp.] under rainfed conditions. International Journal of Agriculture Sciences. 13(2): 202-204.

Peoples, M.B., Unkovich, M.J. and Herridge, D.F. (2009). Measuring symbiotic nitrogen fixation by legumes. Nitrogen Fixation in Crop Production. 52: 125-170.

Ramanjaneyulu, A.V., Reddy, K.I., Bhatt, P.S., Neelima, T.L. and Srinivas, A. (2017). Influence of pigeonpea varieties, N levels and planting methods on yield and economics under rainfed conditions. Legume Research-An International Journal. 40(5): 911-915. Doi: 10.18805/lr.v0i0.7291

Rufini, M., Oliveira, D.P., Trochmann, A., Soares, B.L. andrade, M.J.B.D. and Moreira F.M.D.S. (2016). *Bradyrhizobium* spp. strains in symbiosis with pigeon pea cv. Fava-Larga under greenhouse and field conditions. Revista Brasileira de Ciência do Solo. 40. https://doi.org/10.1590/18069657rbcs20160156 .

Saxena, K., Kumar, C.V., Hingane, A.J., Kumar, M.V., Vijaykumar, R., Saxena, R., Patil, S. and Varshney, R. (2016). Hybrid ICPH 2740 assures quantum jump in pigeonpea productivity in peninsular India. Journal of Food Legumes. 29(2):142-144.

Sekhon, F.S., Singh, T. and Saini, K.S. (2018). Productivity and nutrient uptake of pigeonpea (*Cajanus cajan*) in pigeonpea based intercropping systems as influenced by planting pattern and nutrients levels applied to intercrops. Indian Journal of Agricultural Sciences. 88(10): 1582-1586.

Singh, S., Kumari, N., Singh, C., Karmakar, S. and. Puran, A. (2016). Performance of hybrid pigeonpea performance of hybrid pigeonpea (*Cajanus cajan* L.) as influenced by nutrient and lime levels under rainfed condition of Jharkhand. Supplement on Agronomy. 11(1): 327-330.

Susithra, M., Balasubramanian, A., Gobi, R. and Sathiyamurthi, S. (2019). Yield maximization with nutrient management and nursery practices on transplanted red gram [*Cajanus cajan* (L) millsp.]. Plant Archives. 19(1): 323-326.

Tilak, K.V.B.R., Ranganayaki, N. and Manoharachari, C. (2006). Synergistic effects of plant growth promoting rhizobacteria and *Rhizobium* on nodulation and nitrogen fixation by pigeonpea (*Cajanus cajan*). European Journal of Soil Science. 57(1): 67-71.

Tiwari, K.P. and Namrata, J. (2020). Performance of transplanted pigeon pea [*Cajanus cajan* (L) Millsp.] under rainfed condition of Northern Hill Zone of Chhattisgarh, Madhya Pradesh. International Journal of Current Microbiology and Applied Sciences. 9(7): 3100-3106.

Pradeep, S., Ullasa, M.Y., Kumar Naik, A.H., Ganapathi. and Divya, M. (2018). Effect of different organic nutrient management practices on growth, yield and economics of Pigeonpea, [*Cajanus cajan* (L.) Millsp.] and soil properties. International Journal of Farm Sciences. 7(2): 10-14.

Mallikarjun, C, Hulihalli U.K. and Shantveerayya. (2015). Yield, yield parameters and economics of hybrid pigeonpea (cv.Icph-2671) as influenced by planting methods and geometry. International Journal Agriculture Science. 11(1): 19-23.

Tigga, R. and Singh, S. (2019). Technology transfer through field trials for increasing productivity and profitability of pigeon pea. Journal of Plant Development Sciences. 11(2): 97-100.