RESEARCH ARTICLE Legume Research- An International Journal, Volume 48 Special Issue: 115-120 (November 2025)

Weather Parameter's Impact on Natural Parasitization by *Campoletis chlorideae* Uchida in Chickpea Ecosystem - New Alluvial Zone of West Bengal

C. Meenambigai[1], Arunava Samanta[1], Snigdha Samanta[2], K. Sathees Kumar[3]

10.18805/LR-5187

ABSTRACT

Background: A major biotic constraint in chickpea (*Cicer arietinum* L.) production is gram pod borer, *Helicoverpa armigera* (Hübner). However, the population of this pest is reduced to some extent by a solitary endo-larval parasitoid, *Campoletis chlorideae* Uchida under field conditions. Climate change has a greater impact on natural enemies' effectiveness used in field-level pest management. Therefore, the present study was conducted to understand the impact of weather factors on *C. chlorideae* parasitization rate of *H. armigera* in chickpea grown in the new alluvial agro-climatic zone of West Bengal.

Methods: Chickpea was grown in a randomized complete block design (RCBD) with three replications during *Rabi* 2017-18, 2018-19 and 2019-2020, at Mondouri Farm, Bidhan Chandra Krishi Visawavidayala, Mohanpur, West Bengal. To know the most significant weather parameter which influences *C. chlorideae* parasitization or parasitoid incidence, *H. armigera* larvae (2nd instar) collected from the field experiment were reared until cocoon formation and parasitization % was correlated with weather parameters and subsequently subjected to stepwise regression analysis.

Result: The larval endo-parasitoid, *C. chlorideae* marked its first appearance from 3rd SMW after the appearance of *H. armigera* and its activity was high (68% to 72% parasitization) during the flowering to pod initiation stage and thereafter declined gradually. Among the weather variables, only temperature (maximum and minimum) had a consistent and significant negative correlation with *C. chlorideae* incidence and was responsible for nearly 50% *C. chlorideae* incidence. An increase in 1°C temperature results in a decrease of 4% parasitoid incidence. Thus, climate change in the near future will have a considerable influence on the overall survival, development and rate of parasitization of parasitoids.

Key words: Chickpea, *Campoletis chlorideae, Helicoverpa armigera*, Parasitization.

INTRODUCTION

Chickpea (*Cicer arietinum* L.) also known as Bengal gram or chana is one of the most important pulse crops of India and is considered as "king of pulses" (Bhatt and Patel, 2001). Though India contributes around 71% and 70% of the global chickpea area and production, respectively, the country lags in terms of productivity. Nearly around 60 insect species are known to infest chickpeas. Among that legume pod borer, *Helicoverpa armigera* (Hübner) (Lepidoptera: Noctuidae) is the most important biotic constraint and an insatiable feeder of chickpea flowers, pods and developing seeds (Reed *et al.,* 1987). Repeated usage of conventional insecticides led to high-level resistance development in *H. armigera* (Kranthi *et al.,* 2002). Despite the application of insecticides costing over US$500 million annually, *H. armigera* causes an estimated loss of US$325 million in chickpea which is uneconomical under subsistence farming (Sharma, 2005). Thus, biological control, an important Integrated Pest Management component, comes as a rescue. Chickpea crop harbours very few natural enemies such as *Campoletis chlorideae* Uchida, *Carcelia illota* Curran, *Banchopsis ruficornis* Cameron and *Eriborus* sp. (Srinivas and Jayaraj, 1989; Singh *et al.,* 1991). Among that, the endo-larval parasitoid *Campoletis chlorideae* Uchida (Ichneumonidae: Hymenoptera) is predominant in the chickpea ecosystem. The most preferred host of *C. chlorideae* is *H. armigera* and their parasitization ranges from 8.33% to 78% (Pawar *et al.,* 1989; Dhillon and Sharma, 2007; Gupta and Raj, 2003; Agnihotri *et al.,* 2011). It prefers to parasitize 4 to 5 days-old late 2nd instar *H. armigera* larvae (Dhembare, 1999). Survival threshold and thermal

[1]Department of Agricultural Entomology, Bidhan Chandra Krishi Viswavidyalaya, Mohanpur, Nadia-741 252, West Bengal, India.
[2]Department of Entomology, School of Agriculture and Allied Sciences, The Neotia University, Sarisha-743 368, West Bengal, India.
[3]Department of Agricultural Economics, SRM College of Agricultural Sciences, Chengalpattu-603 201, Tamil Nadu, India.

Corresponding Author: C. Meenambigai, Department of Agricultural Entomology, Bidhan Chandra Krishi Viswavidyalaya, Mohanpur, Nadia-741 252, West Bengal, India.
Email: meenaento@gmail.com

How to cite this article: Meenambigai, C.,Samanta, A., Samanta, S. and Sathees Kumar, K. (2025). Weather Parameter's Impact on Natural Parasitization by *Campoletis chlorideae* Uchida in Chickpea Ecosystem - New Alluvial Zone of West Bengal. *Legume Research.* **48 (Special Issue):** 115-120. doi: 10.18805/LR-5187.

Submitted: 07-06-2023 **Accepted:** 22-03-2024 **Online:** 12-06-2024

requirements influence the abundance and activity of natural enemies, which in turn influence biological control programs' success (Butler and Lopez, 1980; Chihrane *et al.,* 1993). As a result of climate change, the likely increase in temperature and relative humidity (RH) will have a great bearing on the effectiveness of natural enemies for pest management at the field level. For framing a future *H. armigera* bio-intensive management strategy in the chickpea ecosystem, quantification of *H. armigera* mortality by natural field parasitization of *C. chlorideae* and its activity in relation to changing abiotic factors are of major importance. Keeping this in view, the present study was undertaken to study the effect of the influence of weather parameters on the natural parasitization of gram pod borer by *C. chlorideae* in the new alluvial zone of West Bengal.

MATERIALS AND METHODS

The field experiment was conducted during three consecutive *Rabi* seasons of 2017-18, 2018-19 and 2019-2020, at Mondouri farm (22°56′ N, 88°32′E with an altitude of 9.75 m above the msl), Bidhan Chandra Krishi Visawavidayala (BCKV), Mohanpur, which comes under the new alluvial agro-climatic zone of West Bengal. Chickpea seeds (variety: Anuradha) were sown in a 20 m^2 plot with a spacing of 30 cm×10 cm (r-r × p-p). The experiment was laid out in randomized complete block design (RCBD) with three replications and the plots were separated by an alley of 1 m. The crop was grown by following all the recommended agronomic practices except plant protection practices and kept free from pesticide application. From the time of larva appearance in the field, every week thirty late 2nd instar *H. armigera* larvae were randomly collected, brought to the laboratory and reared individually on chickpea leaves in plastic vials. The food was changed regularly as and when required until cocoon formation. Every week, parasitization % by *C. chlorideae* was calculated based on the total number of *H. armigera* larvae collected and number of cocoons formed. Observations on daily data of weather parameters *viz.,* minimum and maximum temperatures (°C), morning and evening RH (%) and rainfall (mm) prevailed during three cropping seasons (December to April 2017-18, 2018-19 and 2019-20) were collected from Department of Agricultural Meteorology, BCKV and then daily data were converted to standard meteorological week (SMW) weather data. *C. chlorideae* mean larval parasitization % (dependent variable) was correlated with the weather parameters (independent variables) and a stepwise multiple regression analysis (both forward and backward) was carried out to get the regression equation for every season. All these statistical analyses were carried out using Minitab software (version 21).

RESULTS AND DISCUSSION

Natural parasitization (%) of *H. armigera* by *C. chlorideae*

During all three cropping seasons, a high parasitism level of *H. armigera* by its larval-endo parasitoid *C. chlorideae* was recorded and this parasitoid marked its 1st appearance (24% parasitization) on 3rd SMW after the appearance of *H. armigera* (Fig 1). During 2017-18 and 2018-19, the larval parasitization % by *C. chlorideae* gradually increased and reached its peak (64% and 72%, respectively) in 7th SMW (3rd week of February). From 8th SMW (4th week of February) there was a regular decline in parasitization from 20% and 40%, respectively, to 4% (11th SMW). While, during 2019-20 *C. chlorideae* parasitization % increased gradually and reached its peak of 68% parasitization in 6th SMW (2nd week of February). Thereafter, from 7th SMW (3rd week of February) a gradual decrease in parasitization of 60% to 4% (11th SMW). Thus, a maximum parasitization of 64% to 72% was recorded during 6th to 7th SMW *i.e.,* flowering to pod initiation stage. During all three years, parasitization became almost nil in 12th SMW (4th week of March) (Fig 1). Jagdish *et al.* (2016) reported the coincidence of parasitoid activity with the crop's flowering and pod formation stage. The maximum parasitization recorded in this study was in agreement with



Fig 1: Natural field *C. chlorideae* parasitization % of *H. armigera* during *Rabi* 2017-18, 2018-19 and 2019-20.

the findings of Ojha *et al.* (2017) and Divija and Agnihotri (2024) who recorded a maximum parasitization of 51.67% to 56.67% and 68% to 72%, respectively, in late sown crop. Ravi and Verma (1997) and Kaur *et al.* (2000) reported that in normal and late-sown crops, *H. armigera* parasitism by *C. chlorideae* was maximum in 6th and 7th SMW, respectively, which was followed by a gradual decline supports the present findings. During three years, the range of minimum - maximum temperatures and morning - evening RH of 9.7°C to 11.01°C - 27.94°C to 28.36°C and 93% to 96.71% - 43% to 50.37%, respectively, along with nil rainfall were found to be favourable for *C. chlorideae* population build-up. The minimum to a maximum threshold temperature and RH of 12°C to 35°C and 95%, respectively, were ideal for the survival, development and increased incidence of *C. chlorideae* (Kaur *et al.*, 2000; Dhillon and Sharma, 2008). Pillai *et al.* (2016) documented that the minimum - maximum temperature as well as morning - evening RH in the range of 8.3°C to 9.2°C - 23°C to 24°C and 89% to 90% - 44% to 54%, respectively and nil rainfall were found to be favourable for the increased *C. chlorideae* incidence in Terai region of Uttarakhand.

Association of *C. chlorideae* incidence with weather variables

A simple correlation was worked out between the weather parameters and *H. armigera* parasitization % by *C. chlorideae* during all three *Rabi* seasons (Fig 2). *C. chlorideae* parasitization % had a significant negative correlation with maximum temperature (r= -0.71* *p*= 0.02; r= -0.63* *p*= 0.04 and r= -0.73**, *p*= 0.01, respectively) and minimum temperature (r= -0.71*, *p*= 0.02, r= -0.73**, *p*= 0.01, r= -0.62*,



where * and ** indicates correlation coefficients significant at $p \leq 0.05$ and $p \leq 0.01$, respectively.

Fig 2: Correlogram showing the association of weather variables and parasitization % *C. chlorideae* during *Rabi* (a) 2017-18, (b) 2018-19 and (c) 2019-20.

p= 0.04), While, remaining independent variables such as evening RH (r= -0.16, *p*= 0.64; r= -0.58, *p*= 0.06 and r= -0.40, *p*= 0.23, respectively), morning RH (r= 0.46, *p*= 0.16; r= -0.47, *p*= 0.14 and r=-0.57, *p*= 0.07, respectively) and rainfall (not available; r= -0.07, *p*= 0.84 and r= -0.48, *p*= 0.14, respectively) showed non-significant association with *C. chlorideae* parsitization (Fig 2). Thus, among the weather variables only temperature (maximum and minimum) had a strong negative association with parasitoid incidence *i.e.,* as the temperature increases, the parasitoid incidence (calculated based on parasitization %) decreases sharply and other weather variables did not play any significant role in influencing parasitization %. The temperature had a negative association with *C. chlorideae* larval and pupal periods and the parasitoid activity was found to cease at a temperature above 35°C to 40°C (Gupta and Raj, 2003; Teggelli *et al.,* 2004; Dhillon and Sharma, 2008). Though Pillai *et al.* (2016) as well as Divija and Agnihorti, (2021) reported that the maximum and minimum temperatures had a significant negative association with *C. chlorideae* incidence, which supports the present result, they have also pointed out that there exists a positive correlation between morning/evening RH and parasitoid incidence. As per Singh *et al.* (2015) RH and rainfall did not play any precise function in *C. chlorideae* parasitization of *H. armigera* which is in agreement with the present findings.

Stepwise regression analysis of parasitoid incidence and weather variables

Based on all three years of weather data, it was confirmed that multicollinearity exists *i.e.,* a condition where explanatory/independent variables are interdependent or a high correlation is observed between independent variables (indicated inside the red colour circle) (Fig 2). Under multicollinearity conditions, one or two best explanatory variables with adequate information that appropriately explain the dependent variable were selected by dropping out other explanatory variables using stepwise regression (Farrar and Glauber, 1967). Based on both forward and backward regression analysis, during 2017-18 and 2019-20, maximum temperature was responsible for 50.28% and 52.71% variation in parasitoid incidence. While, during 2018-19 minimum temperature emerged as the best explanatory variable, responsible for 52.93% variation in parasitoid incidence. The effect of maximum temperature (2017-18 and 2019-20) and minimum temperature (2018-19) was statistically significant at 95% probability level which was evident from the t probability values (*p*= 0.05) (Table 1). Thus, only 50% *C. chlorideae* incidence was significantly influenced by temperature (maximum and minimum). Since the parasitoid incidence does not depend only on weather variables alone, some other factors such as the host density for parasitization and food availability for adults might have contributed to the remaining parasitoid population build-up. Bisane *et al.* (2013) reported that in pigeonpea *H. armigera* larva parasitization by *C. chlorideae* exhibited a density-dependent relationship. In contrast to the present result

Table 1: Estimates of the regression coefficient and t-probability for the explanatory variable selected using stepwise regression analysis of *C. chlorideae* incidence.

Year	Constant	Estimate	Standard error	t Statistics	t Probability	R square	Adjusted R square	Equation (*y=a+bx*)
2017-18	Constant	150.35	42.1	3.57	0.006	50.28%	44.75%	y (parasitoid incidence) = 150.37-4.16 (maximum temperature)*
	Maximum temperature	-4.16	1.38	-3.02	0.015			
2018-19	Constant	92.16	19.43	4.74	0.001	52.93%	47.70%	y (parasitoid incidence) = 92.15-4.78 (minimum temperature)**
	Minimum temperature	-4.78	1.50	-3.18	0.011			
2019-20	Constant	165.4	43.75	3.78	0.004	52.71%	47.46%	y (parasitoid incidence) =165.41-4.83 (maximum temperature)**
	Maximum temperature	-4.83	1.52	-3.17	0.011			

*, ** Significant at $p \leq 0.05$ and $p \leq 0.01$, respectively.

Gupta and Raj (2003) reported that RH and total rainfall had a significant effect of 74.15% on larval parasitism, both individually as well as in association with other abiotic factors. Pillai *et al.* (2016) revealed that various abiotic factors such as minimum and maximum temperatures, morning and evening RH, rainfall, sunshine hours and wind velocity were influencing 97.9% to 99.3% of parasitization. Keeping other factors constant, the model regression equation *(y=a+bx)* indicates that during 2017-18 and 2019-20, every 1°C decrease in maximum temperature resulted in 4.16% and 4.83% increase in the parasitoid population, respectively. During 2018-19 there was 4.78% increase in parasitoid incidence for every 1°C decrease in minimum temperature (Table 1).

CONCLUSION

The maximum and minimum temperatures in the range of 29.14°C to 33.14°C and 12.36°C to 14.69°C, along with the gradual build of *H. armigera* larvae during the flowering to pod initiation stage of the crop favoured *C. chlorideae* population to sustain in the environment. Though *H. armigera* larval incidence increased throughout the cropping season, the parasitoid population declined gradually as the temperature increased. Thus, only temperature (maximum and minimum) has the most significant influence as well as being responsible for nearly 50% *C. chlorideae* incidence or parasitization %. An increase in 1°C of maximum and minimum temperature results in a 4% decrease in parasitoid incidence. Thus, as a result of climate change, the temperature changes may have a considerable influence on the survival, development and parasitization of *C. chlorideae.*

ACKNOWLEDGEMENT

The authors thank the Department of Agricultural Entomology, BCKV for their support in carrying out the experiments. We gratefully acknowledge the Department of Science and Technology (DST), Ministry of Science and Technology, Government of India for their financial support provided under the INSPIRE Fellowship scheme.

Conflict of interest

The authors declare that they have no conflict of interest.

REFERENCES

Agnihotri, M., Gairola, S.C. and Anju, B. (2011). Seasonal incidence of *Campoletis chloridae* Uchida, a larval parasitoid of *Helicoverpa armigera* (Hübner) in chickpea. *Journal of Insect Science.* **24:** 362-366.

Bhatt, N.J. and Patel, R.K. (2001). Screening of chickpea cultivars for their resistance to gram pod borer, *Helicoverpa armigera*. *Indian Journal of Entomology.* **63:** 277-280.

Bisane, K.D., Wadaskar, R.M., and Deotale, R.O. (2013). Tritrophic interaction of *Helicoverpa armigera* (Hubner) in major pulses ecosystem. *Legume Research*. **36:** 116-124.

Butler, G.D. and Jr, Lopez, J.D. (1980). *Trichogramma pretiosum*: development in two hosts in relation to constant and fluctuating temperatures. *Annals of Entomological Society of America.* **73:** 671-673.

Chihrane, J., Lauge´, G. and Hawlitzky, N. (1993) Effects of high temperature shocks on the development and biology of *Trichogramma brassicae* (Hym.: Trichogrammatidae). *Entomophaga.* **38:** 185-192.

Dhembare, A.J. (1999). Preference of host *Helicoverpa armigera* by larval parasite *Campoletis chlorideae* Uchida. *Journal of Experimental Zoology.* **2:** 27-28.

Dhillon, M.K. and Sharma, H.C. (2007). Survival and development of *Campoletis chlorideae* on various insect and crop hosts: Implications for Bt-transgenic crops. *Journal of Applied Entomology.* **131:** 179-185.

Dhillon, M.K. and Sharma, H.C. (2008). Influence of temperature and *Helicoverpa armigera* food on survival and development of the parasitoid, *Campoletis chlorideae*. *Indian Journal Plant Protection*. **36:** 240-244.

Divija, S.D. and Agnihotri, M. (2024). Impact of Abiotic Factors and Dates of Sowing on *Helicoverpa armigera* (Hübner) Parasitoid *Campoletis chloridae* (Uchida) under Chickpea Cropping System in Tarai Region of Uttarakhand. *Legume Research*. **47(3):** 501-504. doi: 10.18805/LR-4500.

Farrar, D.E. and Glauber, R.R. (1967). Multicollinearity in regression analysis: The problem revisited. *The Review of Economics and Statistics.* **49:** 92-107.

Gupta, R.K. and Raj, D. (2003). Natural parasitism by *Campoletis chlorideae* Uchida, a promising parasitoid of *Helicoverpa armigera* (Hubner) on chickpea. *Journal of Biological Control.* **17:** 9-12.

Jagdish, J., Agnihotri, M. and Chakravarty, S. (2016). Biology and relative parasitization of larval endoparasitoid *Campoletis cholrideae* Uchida on *Heliocoverpa armigera* Hübner under sole and chickpea-coriander intercropping system. *Journal of Biological Control*. **30:** 84-90.

Kaur, S., Brar, K.S., Sekhon, B.S., Joshi, N., Shenhmar, M. and Singh, J. (2000). Role played by *Campoletis chlorideae* Uchida in natural mortality of *Helicoverpa armigera* (Hübner) on chickpea in Punjab. *Journal of Biological Control.* **14:** 51-54.

Kranthi, K.R., Jadhav, D.R., Kranthi, S., Wanjari, R.R., Ali, S.S. and Russell, D.A. (2002). Insecticide resistance in five major insect pests of cotton in India. *Crop Protection.* **21:** 449-460.

Ojha, P.K., Kumari, R. and Chaudhary, R.S. (2017). Impact of abiotic and biotic factors on population dynamics of *Helicoverpa armigera* Hubner (Noctuidae: Lepidoptera) in chickpea. *Journal of Entomology and Zoology Studies.* **5:** 636-642.

Pawar, C.S., Bhatnagar, V.S. and Jadhav, D.R. (1989). *Campoletis chlorideae* Uchida (Hymenoptera: Ichneumonidae) as a parasite of *Helicoverpa armigera* (Hub.) (Lepidoptera: Noctuidae) in southwest India. *Proceedings: Animal Sciences.* **98:** 259-265.

Pillai, A.K., Selvaraj, S. and Agnihotri, M. (2016). Seasonal abundance of *Campoletis chlorideae* Uchida (Hymenoptera: Ichneumonidae), A larval parasitoid of *Helicoverpa armigera* (Hubner) in chickpea. *Legume Research.* **39:** 643-647. doi: 10.18805/lr.v39i4.11260.

Ravi, G. and Verma, S. (1997). Seasonal incidence of chickpea pod borer *Helicoverpa armigera* and its larval parasitoids on chickpea crop. *Indian Journal of Entomology.* **59:** 359-361.

Reed, W., Cardona, C., Sithanantham, S. and Lateef, S.S. (1987). The Chickpea Insect Pests and Their Control. The Chickpea. CAB International, Wallington, UK. pp. 283-318.

Sharma, H.C. (2005). Strategies for *Heliothis/Helicoverpa* Management: Emerging Trends and Strategies for Future Research. Oxford and IBH Publishing, New Delhi, India. pp: 465-474.

Singh, D., Singh, S.K. and Vennila, S. (2015). Weather parameters influence population and larval parasitization of *Helicoverpa armigera* (Hübner) in chickpea ecosystem. *Legume Research*. **38:** 402-406. doi: 10.5958/0976-0571.2015.00111.3

Singh, Y., Singh, S.P. and Mathur, K.K. (1991). Parasitization of *Helicoverpa armigera* on Bengal gram in Delhi. *Indian Journal of Entomology.* **53:** 128-133.

Srinivas, P.R. and Jayaraj, S. (1989). Record of natural enemies of *Heliothis armigera* from Coimbatore district, Tamil Nadu. *Journal of Biological Control.* **3:** 71-72.

Teggelli, R.G., Rao, K.J. and Kulkarni, K.A. (2004). Effects of temperature and relative humidity and their interaction on the development of *Campoletis chlorideae* Uchida. *Karnataka Journal of Agricultural Sciences*. **17:** 476-478.