RESEARCH ARTICLE Legume Research- An International Journal, Volume 48 Issue 5: 855-860 (May 2025)

Land Configuration and Nutrient Management on Yield and Nutrient Uptake of Green Gram under Sodic Soil

S. Anandhakrishnaveni[1], N. Karikalan[2], S. Sheeba Joyce Roseleen[3], A.S. Mailappa[4]

10.18805/LR-5456

ABSTRACT

Background: Pulses, endowed with unique ability of nitrogen fixation constitute an important component of crop diversification and resource conservation in farming systems. Green gram [*Vigna radiate*], a drought resistant crop, is one of the thirteen food legumes grown in India and the third most important pulse crop after chickpea and pigeon pea It is a protein rich staple food which contains about 25 per cent protein and plays a crucial role in enhancing soil fertility through biological nitrogen fixation. Sodic soils typically exhibit poor nutrient availability as essential cations are displaced by sodium ions. They significantly influence green gram productivity, posing challenges to its sustainability, with the potential for sodification extending into subsoil layers, impeding plant root development and requiring remedial action. Appropriate land configuration and nutrient management could mitigating the challenges posed by sodic soils for pulse crops. Understanding the effects of land configuration and nutrient management on its growth, yield andeconomic viability is vital for sustainable agriculture. Keeping this in view, the field experiment was conducted to study the effect of different land configuration and nutrient management for their impact on plant height, dry matter production, grain yield, haulm yield, net income and benefit cost ratio with nutrient uptake under sodic condition.

Methods: The land configuration factor consists of the three treatments *viz.,* flat bed, ridges and furrows and broad bed furrow system. The second factor nutrient management comprising of five treatments like 100% RDF + 1% DAP, 100% RDF + 1% MAP, 75% RDF + 2% MAP, 100% RDF + 1% MKP, 75% RDF + 2% MKP. The RDF for green gram was 25 kg N + 50 kg P_2O_5 + 25 kg K_2O per hectare.

Result: Higher plant height, dry matter production, grain and haulm yield, protein yield, economic and nutrient uptake was recorded in broad bed furrow over farmer's practice of flatbed method. N_4 - 100% RDF + 1% MKP foliar spray twice at (plant height, dry matter production, grain and haulm yield, protein yield, economic and nutrient uptake) were observed than farmer's practice. Hence, Broad bed furrow and nutrient management practice of 100% RDF + 1% MKP has the potential to enhance the productivity green gram under sodic condition.

Key words: Broad bed furrow, Grain yield, Green gram, Nutrient uptake, Protein yield.

INTRODUCTION

Green gram [*Vigna radiata* (L), one of the important pulses in India, has 4.5 million hectares area under cultivation, with a total production of 2.5 million tonnes and productivity of 548 kg/ha, contributing 10% to the total production (Anon., 2021). Green gram has significant amount of protein and is easier to digest than other pulses. Though pulse crops are nutritionally richer than cereals, they have comparatively lower productivity. In order to increase productivity, various crop management practices need to be optimized.

The problem of water logging in sodic soil, even for a short period, proves detrimental to the green gram, particularly during the early growth period (Karikalan *et al.,* 2020). The high levels of sodium in sodic soil negatively impact soil structure and nutrient availability. Sodic soils typically exhibit the presence of excess exchangeable sodium which adversely affect the physical and nutritional properties of the soil (Bahadhur *et al*., 2012). Land configuration plays a crucial role in influencing the root and nodule growth in green gram (Sweta Rath *et al.,* 2023). The cultivation of greengram faces many problems like water logging and inadequate aeration, which affect the growth and yield in flatbed conditions adversely. Further the major causes of poor productivity and soil fertility are imbalanced application of nutrients (Porpavai and Nagarajan, 2022). Water is a critical input for sustained crop production and is becoming limited day by day. Hence, the need of the hour is to evaluate effective resource

[1]Department of Agronomy, Anbil Dharmalingam Agricultural College and Research Institute Navalur Kuttapattu Tiruchirappalli-620 027, Tamil Nadu, India.

[2]Department of Agronomy, Dhanalakshmi Srinivasan Agricultural College, Perambalur-621 212, Tamil Nadu, India.

[3]Department of Agricuiture Entomology, Horticultural College and Research Institute for Women, Tamil Nadu Agricultural University, Navalur Kuttapattu, Tiruchirappalli-620 027, Tamil Nadu, India.

[4]Department of Natural Resource Management, College of Horticulture and Forestry, Central Agricultural University, Pasighat-791 102, Arunachal Pradesh, India.

Corresponding Author: S. Sheeba Joyce Roseleen, Department of Agricuiture Entomology, Horticultural College and Research Institute for Women, Tamil Nadu Agricultural University, Navalur Kuttapattu, Tiruchirappalli-620 027, Tamil Nadu, India.
Email: sheeba@tnau.ac.in

How to cite this article: Anandhakrishnaveni, S., Karikalan, N., Roseleen, S.S.J. and Mailappa, A.S. (2025). Land Configuration and Nutrient Management on Yield and Nutrient Uptake of Green Gram under Sodic Soil. Legume Research. 48(5): 855-860. doi: 10.18805/LR-5456.

Submitted: 06-12-2024 **Accepted:** 30-12-2024 **Online**: 25-01-2025

conservation practices, such as land configurations and scientific scheduling of irrigation to achieve higher productivity, profitability and resource use efficiency (Aakash Meher *et al*., 2024). Appropriate land configuration and nutrient management could mitigate the challenges posed by sodic soils for pulse crops. For example, the adoption of raised beds or broad bed furrows (BBF) improves soil structure, enhances water drainage, aids in root development andallows better aeration in sodic soils (Dhimmar, 2003 and Akbar *et al.,* 2007). Optimum nutrient management involves adjusting external nutrient application to address sodicity-induced deficiencies, ensuring pulse crops access essential nutrients adequately. The application of essential plant nutrients through foliar spray in sodic soil prevents the interaction of applied nutrients with soil sodium, thereby enhancing nutrient use efficiency and facilitating the rapid correction of nutrient deficiencies in crop plants (Vijayakumar *et al.,* 2022). Given the aforementioned facts, a study was conducted with the aim of identifying appropriate land configuration and nutrient management practices for green gram under sodic soil during the rainy season.

Materials and Methods

The field experiment was conducted at Anbil Dharmalingam Agriculture College and Research Institute, Trichy during *kharif* season of 2020 and 2021.The experimental field featured a sandy clay loam soil texture with an alkaline (pH = 8.1; EC = 4.6 dSm) reaction. The study was designed using Factorial Randomised Block Design with three replications. Fifteen treatment combinations were tested, including three levels of land configuration (factor A) *viz.,* L_1 - flat bed, L_2 - ridges and furrows and L_3 - broad bed furrow system (BBF) and five levels of nutrient management (factor B), namely N_1 - farmer's practice of 100% recommended dose of fertilizer (RDF) + 2 % Diammonium phosphate (DAP), N_2 - 100% RDF + 1 % Monoammonium Phosphate (MAP), N_3 - 75% RDF + 2% MAP, N_4 - 100% RDF + 1% Monopotassium Phosphate (MKP), N_5 - 75% RDF + 2% MKP. The RDF for greengram was 25 kg N + 50 kg P_2O_5 + 25 kg K_2O per hectare and applied as 55 kg of urea, 315 kg of single super phosphate and 42.5 kg of murate of potash. The selected green gram variety for the study was VBN (Gg) 2.

The spray solutions of DAP, MKP and MAP were prepared by dissolving 10 g and 20 g of fertilizers in one litre of water to achieve 1% and 2% concentration, respectively. MKP and MAP fertilizers are highly water-soluble salts that completely dissolve in water, but DAP granules were dissolved in a small quantity of water and allowed to settle overnight. The supernatant solution was taken for spraying after dilution with remaining quantity of water. The plant height and dry matter production were recorded for five plants at the harvesting stage andthe averages were reported. Data on growth, yield attributes and yield were recorded plot-wise from each replication.

Representative samples of seed were taken from each treatment and dried in oven at 65°C temperature for 24 hours and powdered by mechanical grinder. Then the nitrogen content of seed was determined by micro Kjheldahl's method. The protein content in seed was calculated by multiplying nitrogen content of seed (%) with the conversion factor of 6.25 (Bhuiya and Chowdhary 1974). The protein yield in seed was calculated by using the following formula:

$$\text{Protein yield}\left(\frac{\text{kg}}{\text{ha}}\right) = \frac{\text{Protein content in seed (\%)} \times \text{Seed yield (kg/ha)}}{100}$$

Statistical analysis

The pooled data was subjected to statistical scrutiny by Panse and Sukhatme (1985). The ANOVA and critical difference at five per cent level of significance were calculated.

Results and Discussion

Land configuration

Among the land configuration methods tested, Broad Bed Furrow (BBF) system recorded the highest plant height (49.9 cm) andit remained on par with the ridges and furrow method (48.1 cm) at harvest stage. On contrary, the lowest plant height (43.3 cm) was recorded in flat bed method. The plant height under BBF was 15 times (6.6 cm) higher than that of the flat method (Table 1). This might be attributed to the effective maintenance of suitable moisture regimes and the adequate availability of plant nutrients, thereby contributing to the improved growth and development of the plant under both methods. Similar positive effect of broad bed method of sowing on growth has registered by Dhimmar (2003), Sangakkara (2004) and Kumar and Singh, (2014). Correspondingly, dry matter production exhibited a significant increase with the BBF method (2874 kg ha^{-1}), followed by ridges and furrows method (2616 kg ha^{-1}) at harvest stages. Similar to plant height, the lowest dry matter production (1986 kg ha^{-1}) was recorded in flat bed method. Under BBF method, the dry matter production increased by 59.8% over the flat bed method. This heightened dry matter production in BBF is attributed to the overall increased growth of plant in terms of plant height and number of branches per plant. similar findings have also been reported by Lewade (2017) and Sodavadiya (2017). Consistent with our study results, Singh *et al.* (2020) reported higher plant height and dry matter production under the BBF and the ridges and furrow method compared to the flat bed method. Similar report was given by Sainath *et al*. (2023).

The seed yield (854.6 kg ha^{1}) and haulm yield (1952.8 kg ha^{-1}) of green gram under BBF increased by 58.2% and 36.5%, respectively over flatbed method Fig 1. Additionally, this method also resulted in a higher net return of ` 29,955 ha^{-1} and a BCR value of 2.0 (Table 1). Following closely, the ridge and furrow method yielded 40% higher grain yield and 25.4% higher haulm yield than flatbed method.

Similarly, the highest net return and B: C ratio was recorded in BBF, followed by ridges and furrow, while the lowest was recorded in the flat bed method. BBF recorded a net return of ¹ 23,244 ha^{-1} with a BCR of 1.8. The net return and B: C ratio recorded in the BBF method were 139% and 33.3% higher than the flat bed method, respectively.

The interaction between land configuration and nutrient management had a significant impact. The findings indicate that the combination of L_3N_4 positively enhances greengram grain yield performance by improving growth parameters.

Regarding protein yield (Table 2), once again, BBF recorded a higher protein yield (184.89 kg ha^{-1}), which is 12.5% and 62.5% higher than the ridges and furrow and flat bed methods, respectively. Due to its higher seed yield and protein content, BBF significantly outperformed both the ridge and furrow method and the flat bed method in protein yield. The results obtained were consistent with those reported by Subbulakshmi *et al.* (2024), who observed higher grain yield, net return anda better benefit-to-cost ratio (B:C ratio) in the best bet fertilization (BBF) treatment under conditions of moisture stress. These results were supported with the findings of Shete *et al.* (2010) in green gram.

With respect to nutrient uptake (Table 2), significantly higher NPK uptake (52.2, 11.1, 39.0 kg N, P, K/ha) was observed under the BBF method, which was comparable to that of the ridges and furrow at harvest. Conversely, lower NPK uptake (49.1, 10.4, 37.1 kg N, P, K/ha) was recorded with flatbed method. This discrepancy may be attributed to the deeper penetration of root and suitable soil environment provided by BBF method over the flat bed method. Additionally, the better aeration, microbial activity, optimum moisture and nutrients for crop growth under sodic condition likely contributed to high nutrient recovery by seed and haulm in the BBF method of sowing, reflecting in the overall higher uptake of nutrients. These results are in close conformity with Muralidaran and Solaimalai (2010).

Table 1: Effect of land configuration and nutrient management on growth, yield and economics of green gram.

Treatment	Plant height (cm)	Dry matter Production (kg ha^{-1})	Grain yield (kg ha^{-1})	Haulm yield (kg ha^{-1})	Net income (Rs.)	B:C Ratio
Land configuration						
L_1 - Flat Bed	43.32	1986	540.2	1430.2	12511	1.5
L_2 - Ridges and Furrows	48.12	2616	761.3	1794.4	23244	1.8
L_3 - Broad Bed Furrow	49.91	2874	854.6	1952.8	29955	2.0
SEd	1.11	74	20.31	63.54	-	-
C.D (P=0.05)	2.27	153	41.61	130.16	-	-
Nutrient management						
N_1 - 100%RDF + 2% DAP	47.73	2493	730.1	1721.89	22944	1.8
N_2 - 100%RDF +1% MAP	49.35	2764	792.3	1912.56	26110	1.9
N_3 - 75 %RDF + 2% MAP	42.92	2086	600.8	1438.33	15207	1.6
N_4 - 100%RDF +1% MKP	49.80	2860	801.4	1996.22	26794	2.1
N_5 - 75 %RDF + 2% MKP	45.81	2257	651.7	1560.00	18460	1.7
SEd	1.43	96	26.22	82.03	-	-
C.D (P=0.05)	2.29	197	53.72	168.04	-	-
Interaction between L × N						
L_1N_1	43.22	2123	590.2	1510.4	15544	1.6
L_1N_1	44.85	2312	610.3	1612.6	17582	1.7
L_1N_1	41.82	1943	540.1	1380.1	12001	1.4
L_1N_1	45.31	2456	620.5	1650.2	18312	1.7
L_1N_1	42.87	2096	550.4	1465.8	13018	1.5
L_2N_2	48.34	2545	750.3	1805.9	24211	1.9
L_2N_2	49.93	2820	780.1	1935.8	26515	2
L_2N_3	45.23	2180	690.2	1510.2	17250	1.7
L_2N_4	50.85	2903	810.6	2011.8	27831	2.2
L_2N_5	46.78	2368	710.2	1632.4	19162	1.8
L_3N_1	49.63	2810	850.2	1950.4	29177	2.1
L_3N_2	53.28	3160	910.5	2290.2	34232	2.3
L_3N_3	46.72	2135	672.2	1506.7	16370	1.8
L_3N_4	53.24	3222	850.1	2326.7	34239	2.4
L_3N_5	48.78	2311	695.7	1581.8	18599	1.9
SEd	1.43	96	26.2	82	-	-
CD (P=0.05)	2.29	197	53.7	168	-	-

Nutrient management

The significantly highest plant height (49.8 cm) and plant dry matter (2860 kg ha^{-1}) at harvest were recorded with 100% RDF + 1% MKP at harvest, which remained on par with 100% RDF + 1% MAP, followed by the farmer's practices of 100% RDF + 2% DAP (Table 1). This result could be attributed to the application of full recommended fertilizer dose at basal, along with an additional application of K as MKP through foliar spray which might aid in the better uptake of nutrients by lowering the concentration of Na^+ salts in plants, particularly under sodic condition. Similar results was observed by Mohammadi *et al.* (2008), Pessarakli and Kopec (2009) and Tsegay and Gebreslassie (2014).

The application of 100% RDF + 1% MKP produced significantly higher seed yield (797 kg ha^{-1}) and haulm yield (1996 kg ha^{-1}) (Fig 1.), which remained on par with 100% RDF + 1% MAP and 100% RDF + 2% DAP. This might be due to optimum supply of P and K, enabling the green gram crop to increase the absorption of other essential nutrients and assimilate adequate photosynthetic products under sodic conditions. This cumulative effect resulted in the plants producing more pods with a greater number of seeds, higher test weight and ultimately maximum seed and haulm yield. This results are supported by Gawai and Pawar (2007) in chickpea and Geetha and Velayutham, (2009).

In terms of economics, the highest net return of ₹ 26794 ha^{-1} and a BCR of 2.1 were obtained under the treatment 100% RDF + 1% MKP, followed by 100% RDF + 1% MAP with net return of ₹ 26110 ha^{-1} and BCR of 1.9 (Table 1). This is due to higher seed and haulm yields of green gram in these treatment in comparison to the farmer's practice. The result corroborates with the findings of Gupta *et al* (2011) and Kulkarni *et al.* (2016). Treatment 100% RDF +

Table 2: Effect of land configuration and nutrient management on quality and nutrient uptake of greengram.

Treatment	Protein yield (kg ha^{-1})	Nutrient uptake (kg ha^{-1})		
		N	P	K
Land configuration				
L_1 - Flat Bed	113.72	49.13	10.40	37.11
L_2 - Ridges and Furrows	164.32	51.43	10.91	38.81
L_3 - Broad Bed Furrow	184.89	52.23	11.07	38.97
SEd	6.03	1.03	0.16	0.72
C.D (P=0.05)	12.36	2.11	0.33	1.48
Nutrient management				
N_1 - 100%RDF + 2% DAP	156.21	51.20	10.98	38.74
N_2 - 100%RDF +1% MAP	178.23	52.50	11.71	39.77
N_3 - 75 %RDF + 2% MAP	121.87	48.23	9.68	36.08
N_4 - 100%RDF +1% MKP	182.43	52.93	11.78	40.08
N_5 - 75 %RDF + 2% MKP	132.81	49.80	9.97	36.81
SEd	7.79	1.32	0.21	0.93
C.D (P=0.05)	15.95	2.71	0.42	1.91
Interaction between L × N				
L_1N_1	127.42	49.60	10.80	37.62
L_1N_1	143.32	50.88	11.24	38.12
L_1N_1	98.61	47.21	9.53	35.42
L_1N_1	148.18	50.94	11.32	38.34
L_1N_1	110.08	48.12	9.73	36.05
L_2N_2	168.41	51.61	10.97	38.89
L_2N_2	192.68	52.91	11.72	39.90
L_2N_3	137.83	49.24	9.75	36.19
L_2N_4	199.84	53.40	11.81	40.42
L_2N_5	145.85	50.01	9.94	37.02
L_3N_1	172.81	52.38	11.16	39.71
L_3N_2	198.69	53.71	12.18	41.29
L_3N_3	129.18	48.25	9.77	36.64
L_3N_4	199.26	54.45	12.22	41.47
L_3N_5	142.51	51.26	10.25	37.37
SEd	7.79	1.32	0.21	0.93
CD (P=0.05)	15.95	2.71	0.42	1.91

N: Nitrogen; P: Phosphorus; K: Potassium.




Fig 1: Effect of land configuration and nutrient management on grain yield and haulm yield.

1% MKP recorded a higher protein yield (182.43 kg ha^{-1}) and it remains on par with treatment 100% RDF + 1% MAP, followed by 100% RDF + 2% DAP. This might be due to fact that an adequate supply of P and K through foliar spray ensures optimum nitrogen supply by promoting root nodulation process and reducing Na^+ salt concentration in plants. This combined effect increased the nitrogen content in seed, leading to higher protein content. Nitrogen serves as a structural element of several co-enzymes which involved in protein synthesis. The higher protein content and seed yield ultimately result in increased protein yield under this treatment. Choudhary *et al.* (2013) with respect to protein yield of green gram.

Significantly maximum NPK uptake (52.9, 11.8, 40.1 kg N, P, K/ha) was observed with 100% RDF + 1% MKP and it remain on par with 100% RDF + 1% MAP and farmer's practice of 100% RDF + 2% DAP. This is due to proper nutrient management, more nodules formation, resulting in higher uptake of N, P and K in green gram compared with other treatment under sodic conditions. These results are in close conformity with of Veerabhadrappa and Yeledhalli (2010).

A significant difference was observed in the interaction between land configuration and nutrient management concerning protein yield and nutrient uptake. The combination of L_3N_4-BBF sowing with 100% RDF + 1% MKP-significantly recorded higher protein yield and nutrient uptake.

CONCLUSION

The results revealed that, among the land configuration methods, the broad bed furrow (BBF) method of sowing demonstrated a favourable impact on greengram, influencing factors such as plant population, growth and yield attributes, ultimately leading to a 12% increase in seed yield (854.6 kg ha^{-1}) and haulm yield (1952.8 kg ha^{-1}) along with significantly higher NPK uptake compared to the flat bed method. Among the nutrient management practices, the application of 100% RDF + 1% MKP produced significantly higher seed yield (797 kg ha^{-1}), haulm yield (1996 kg ha^{-1}) and higher protein yield (182.43 kg ha^{-1}). In terms of economics, the highest net return of ` 26794 ha^{-1} and a BCR of 2.1 were obtained under the treatment of 100% RDF + 1% MKP. Thus, the study highlights the potential of sowing green gram on Broad Bed Furrows (BBF) combined with the application of 100% RDF with 1% MKP foliar spray as effective strategies for maximizing productivity and profitability under sodic soil conditions, these practices could significantly contribute to sustainable agriculture by enhancing resource use efficiency and economic viability.

Conflict of interest

All authors declare that they have no conflicts of interest.

REFERENCES

Akbar, G., Hussain, Z. and Yasin, M.J. (2007). Problems and potentials of permanent raised bed cropping systems in Pakistan. Pakistan Journal of Water Resources. 11: 11-23.

Anonymous, (2021). Green Gram Outlook Report- January- May. Bhuiya, Z.H. and Chowdhary S.V. (1974). Effect of N, P, K and S on quality of groundnut. Indian Journal of Agricultural Science. 44: 751-754.

Bahadur Lal, D., Tiwari Mishra, D.J. and Gupta B.R. (2012). Effect of integrated nutrient management on yield, microbial population and changes in soil properties under rice-wheat cropping system in sodic soil. Journal of the Indian Society of Soil Science. 60: 326-329.

Bhuiya, Z.H. and Chowdhary, S.V. (1974). Effect of N, P, K and S on quality of groundnut. Indian Journal of Agricultural Science. 44: 751-754.

Choudhary, H.R., Sharma, O.P., Singh, R.K., Singh, K., Kumar, R. and Yadav, L. (2013). Influence of organic manures and chemical fertilizer on nutrient uptake, yield and profitability of mungbean [*Vigna radiata* (L.) Wilczek]. Madras Agric Journal. 100: 747-750.

Dhimmar, S.K. (2003). Response of *kharif* cowpea [*Vigna unguiculata*. (L.) Walp.] to land configuration and bio-fertilizers under south Gujarat conditions. Agronomy Department, NM College of Agriculture Gujarat Agricultural University.

Gawai, P.P andPawar V.S. (2007). Nutrient balance under INMS in sorghum-chickpea cropping sequence. Indian Journal of Agricultural Research. 41: 137-141.

Geetha, P. and Velayutham, A. (2009). Refinement of nutrient management techniques for growth, yield and nutrient uptake of rice fallow black gram. Madras Agric. Journal. 96: 163-166.

Gupta, S.C., Kumar, S. and Khandwe, R. (2011). Effect of bio-fertilizers and foliar spray of urea on symbiotic traits, nitrogen uptake and productivity of chickpea. Food Legumes. 24: 155-157.

Karikalan, N., Anandha Krishnaveni, S., Avudaithai, S. and Nithila, S. (2020). Effect of land configuration and nutrient management on yield and economics of greengram under sodic soil. International Journal of Current Microbiology and Applied Sciences. 9: 2125-2129.

Kulkarni, S., Upperi, S.N. and Jadhav, R.L. (2016). Green gram productivity enhancement through foliar spray of nutrients. Legume Research. 39: 814-816. doi: 10.18805/LR-4109.

Kumar, B. and Singh G.R (2014). Response of land configurations, IW/CPE ratios and integrated nutrient supply systems on growth function, yield and water use efficiency of French bean (*Phaseolus vulgaris* L.). International Journal of Agriculture, Environment and Biotechnology. 7: 825-831.

Lal, B., Tiwari, DD., Mishra, J. and. Gupta B.R. (2012). Effect of integrated nutrient management on yield, microbial population and changes in soil properties under rice-wheat cropping system in sodic soil. Journal of the Indian Society of Soil Science. 60: 326-329.

Lewade (2017). Effect of land configurations and foliar sprays on soybean [*Glycine max* (L.) Merrill] + pigeonpea [*Cajanus cajan* (L.) Mill sp.] intercropping system under rainfed condition. Vasantrao Naik Marathwada Krishi Vidyapeeth, Parbhani.

Meher, A., Sujatha, H.T., Korav, S., Bisarya, D., Adishesha, K. (2024). Resource use efficiency and economics of summer greengram as influenced by land configuration and irrigation regimes in North-western part of India. Agricultural Science Digest. doi: 10.18805/ag.D-5836.

Mohammad, P. and Kopec, D.M. (2009). Screening various ryegrass cultivars for salt stress tolerance. J. Food Agric. Environ. 7: 4.

Mohammadi, H., Poustini, K. and Ahmadi, A. (2008). Root nitrogen remobilization and ion status of two alfalfa (*Medicago sativa* L.) cultivars in response to salinity stress. Journal of Agronomy and Crop Science. 194: 126-134.

Muralidaran, C. and Solaimalai, A. (2010). Studies on land cofiguration methods, row spacing and time of nitrogen application on nutrient uptake and yield of cotton in dryland condition of westrern zone of Tamil Nadu. Karnataka. Journal of Agricultural Sciences. 18: 1-3.

Panse, V.G. and Sukhatme, P.V. (1985). Statistical Methods for Agricultural Workers. ICAR, New Delhi.

Porpavai, S., Nagarajan, M. (2022). Effect of land configuration and nutrient management methods on growth and yield of blackgram (*Vigna mungo*). Agricultural Science Digest. 42(1): 88-90. doi: 10.18805/ag.D-5245.

Rath, S., Mishra, G., Gulati, J.M.L., Mohapatra, A.K. (2023). Root growth, nutrient uptake, nodulation and yield of summer mung [*Vigna radiata* (L.) Wilczek] as Influenced by land configuration and varying nutrient management practices under Coastal Plains of Odisha Indian Journal of Agricultural Research. 57(4): 461-467. doi: 10.18805/IJARe.A-6084.

Sainath, K., Singh, B., Bokado, K. and Narender Roa, K. (2023). Growth, Yield and water use efficiency of Greengram (*Vigna radiata* L.) as influenced by irrigation scheduling and land configuration. Eco. Env. and Cons. 29 (October Suppl. Issue): S274- S278.

Sangakkara, U.R. (2004). Effect of tillage and moisture levels on growth, yield and nodulation of common bean (*Phaseolus vulgaris*) and mungbean (*Phaseolus radiatus*) in the dry season. Indian Journal of Agronomy. 49: 60-63.

Shete, P.G, Baviskar V.S. and. Adhav, S.L. (2010). Effect of land configuration, inorganic fertilizers and levels of FYM on quality and nutrient status of *rabi* green gram. International Journal of Agricultural Sciences. 6: 546-548.

Singh, G., Virk, H.K. and Khanna, V. (2020). Effect of land configuration and weed management on productivity of Greengram (*Vigna radiata*). The Indian Journal of Agricultural Sciences. 947-951.

Sodavadiya, H.B. (2017). Effect of land configuration, irrigation and INM on growth and yield of Indian bean (VAR. GNIB-21). Department of Agronomy Nm College of Agriculture Navsari.

Subbulakshmi, S., Selvarani, A., Sanjivkumar, A.V. and Baskar, K.L. (2024). Land configurations and drought mitigation practices to improve productivity and profitability of blackgram (*Vigna mungo*) under moisture stress conditions. The Indian Journal of Agricultural Sciences. 94: 96-99.

Tsegay, B.A. and Berhane, G.G. (2014). The effect of salinity (NaCl) on germination and early seedling growth of *Lathyrus sativus* and *Pisum sativum* var. *abyssinicum*. African Journal of Plant Science. 8: 225-231.

Veerabhadrappa, B.H. and Yeledhalli, N.A. (2010). Effect of soil and foliar nutrition on partitioning of nutrients at different growth stages of groundnut. Karnataka Journal of Agricultural Sciences. 18: 3-7.

Vijayakumar, S., Gobinath, R., Gompa, S.J. Surekha, K., Kumar, R.M. and Sundaram, R.M. (2022). Management of micro-nutrient deficiency and toxicity in rice. Indian Farming. 72: 11-14.